

山形県月山の樹林帯の積雪上に現れる雪氷藻類のブルームとその要因

Environmental factors associated with snow algal bloom in the deciduous forest of Mt. Gassan, Yamagata prefecture, Japan

*渡邊 茜¹、竹内 望¹

*Akane Watanabe¹, Nozomu Takeuchi¹

1. 千葉大学大学院理学研究科

1. Graduate School of Science, Chiba University

Snow algae are photosynthetic microbes inhabiting alpine and polar snow fields. They bloom on melting snow and change snow color from white to red or green. Colored snow appears widely in mountainous regions in Japan. However, the conditions for the algal bloom are still not understood well. In this study, we aim to describe the temporal change of snow algae and physical and chemical conditions of surface snow in Mt. Gassan, Yamagata prefecture, Japan. Study site is located in the Japanese beech forest at an elevation of 750 m a.s.l. Field studies were carried out three times from April to May of 2016. We collected samples of surface snow and snow pit down to the ground surface at the study site. In laboratory, we measured chlorophyll-a concentrations, EC, pH, and soluble chemical compositions in the samples.

Field observations revealed that algal green snow appeared patchy after the late April. The algal patches were frequently observed in the snow surface under trees compared with the open sunny surface. The chlorophyll-a concentration in the surface snow gradually increased during the study period. The analysis of major soluble ions revealed that the phosphate concentration in the surface snow under the trees increased up to 51.4 $\mu\text{eq/L}$ while it in the open snow surface kept low value during the study period. Results suggest that the phosphate is supplied with rain water from canopy of the trees to the snow surface and that it causes the patchy algal bloom on the snow surface.

キーワード：雪氷藻類、栄養塩、落葉樹林

Keywords: snow algae, nutrients, deciduous forest

札幌で8冬期間に観測された積雪不純物がアルベドに与える効果

Effect of snow impurities on albedo observed during 8 winter seasons in Sapporo

*広沢 陽一郎¹、青木 輝夫^{2,3}、庭野 匡思³、的場 澄人⁴、兒玉 裕二⁵

*Yoichiro Hirozawa¹, Teruo Aoki^{2,3}, Masashi Niwano³, Sumito Matoba⁴, Yuji Kodama⁵

1. 岡山大学 理学部、2. 岡山大学 大学院自然科学研究科、3. 気象研究所、4. 北海道大学 低温科学研究所、5. 国立極地研究所

1. Okayama University Faculty of Science, 2. Okayama University, Graduate School of Natural Science and Technology, 3. Meteorological Research Institute, 4. Institute of Low Temperature Science, Hokkaido University, 5. National Institute of Polar Research

近年、積雪面積や積雪期間は主に北極域において減少している。雪氷は一般にアルベドが高いため、地球温暖化に伴う雪氷の融解はアルベドを低下させる。その結果、地表面による太陽放射の吸収の増加し、地球温暖化をさらに加速させる。代表的な豪雪地帯である札幌において、積雪アルベドは積雪粒径と積雪不純物濃度に強く依存していることが報告されている (Aoki et al., 2003, 2007)。本研究では、札幌の積雪アルベドにおける積雪不純物濃度の影響を、Aoki et al. (2011) によって開発された積雪アルベド物理モデル (PBSAM) を用いて調べた。観測場所は北海道大学低温科学研究所の気象観測露場 (43°04'56"N, 141°20'30"E, 15 m a.s.l.) で、観測期間は2007-2015年の8冬期間である。積雪断面観測及び放射観測データをPBSAMに入力して得られる広波長帯域アルベドの理論計算値と観測値を比較した。さらに、積雪不純物によるアルベドの変化に関する感度実験も行った。

解析期間の各年で時系列の観測値とモデル計算値を比較すると、積雪粒径や積雪不純物濃度の変化に対応したアルベド変動の計算値は観測値とよく一致していることが分かった。全解析期間において、短波長域 (SW) におけるアルベド観測値と計算値の比較から得られる決定係数 (R^2) と二乗平均平方根誤差 (RMSE) はそれぞれ0.831と0.045で、これらの結果からPBSAMの高い精度が確認された。次に、ブラックカーボン (BC) と鉱物性ダストからなる積雪不純物の有無による、可視域、近赤外域、短波長域のアルベド変化に関する感度実験を行った。全解析期間における積雪不純物 (BC+ダスト) によるアルベド変化は可視域で-0.085、近赤外域で-0.016、短波長域で-0.053であった。また、短波長域のアルベド変化へのBCとダストそれぞれからの寄与を調べた結果、全解析期間の短波長域のアルベド変化はBCのみにより-0.043、ダストのみにより-0.009であった。さらに、涵養期と融雪期間の短波長域のアルベド変化の比 (融雪期/涵養期) はBCが4.1倍、ダストが12.0倍、BC+ダストが4.9倍であった。

参考文献

Aoki et al., 2003: *J. Geophys. Res.*, 108(D19), 4616, doi:10.1029/2003JD003506.

Aoki et al., 2007: *Ann. Glaciol.*, 46, 375–381, doi:10.3189/172756407782871747.

Aoki et al., 2011: *J. Geophys. Res.*, 116, D11114, doi:10.1029/2010JD015507.

Observational study on spatial development structure of blowing snow

*杉浦 幸之助¹、大井 聖也¹、戸沢 達矢¹

*Konosuke Sugiura¹, Seiya Ooi¹, Tatsuya Tozawa¹

1. 富山大学

1. University of Toyama

Reducing visibility and forming snow drifts by blowing snow are still social natural disaster and a major problem. Although a large number of studies have been made on the vertical development of blowing snow, little is known about the horizontal development of blowing snow especially focused on near the snow surface. For a better understanding of spatial development structure of blowing snow, observations about the horizontal development of blowing snow near the surface have been carried out. The results obtained by the cold wind-tunnel experiments showed if the wind became strong, the development to a horizontal direction became weaker than that to a vertical direction. In addition, the results of field observations in blowing snow at Sapporo showed blowing-snow flux remarkably increased when the wind blew from the direction of a long fetch. It is necessary to carry out the spatial observation above the blowing snow layer using such as a kite in the field in future while investigating the horizontal development of blowing snow experimentally in detail.

キーワード：吹雪、空間発達

Keywords: blowing snow, spatial development

Assimilation of all-sky GCOM-W/AMSR2 brightness temperature using a strongly coupled atmosphere-land data assimilation system in snowy Siberia

*鈴木 和良¹、Zupanski Milija²、Zupanski Dusanka³、寺崎 康児⁴、三好 建正⁴

*kazuyoshi suzuki¹, Milija Zupanski², Dusanka Zupanski³, Koji Terasaki⁴, Takemasa Miyoshi⁴

1. 国立研究開発法人海洋研究開発機構、2. コロラド州立大学、3. Spire Global Inc.、4. 国立研究開発法人理化学研究所
1. Japan Agency for Marine-Earth Science and Technology, 2. Colorado State Univ., 3. Spire Global Inc., 4. RIKEN

Coupled numerical models address the interaction between processes in the atmosphere, ocean, land surface, biosphere, chemistry, cryosphere, and hydrology. Including the interaction between such processes can potentially extend the predictability and eventually help in reducing the uncertainty of the prediction. Coupled data assimilation is a branch of data assimilation that deals with coupled modeling systems. There are two kinds of coupled data assimilation systems such as weakly and strongly coupled data assimilation. Recently we developed a strongly coupled atmosphere-land data assimilation system (Suzuki et al., 2017). In this article the fundamentals of bias correction for the all-sky GCOM-W/AMSR2 brightness temperature using coupled data assimilation are described. Through a series of data assimilation experiments, we analyze the effectiveness of bias correction coefficients and predictors. Through this study, we analyze the impact of all-sky brightness temperature in reanalysis. Finally, applying coupled data assimilation can visualize more details of coupled atmosphere-land interaction.

Reference

Suzuki, K., Zupanski, M. and Zupanski, D. (2017), A case study involving single observation experiments performed over snowy Siberia using a coupled atmosphere-land modelling system. *Atmos. Sci. Lett.* doi:10.1002/asl.730

キーワード：結合データ同化、大気積雪相互作用、降雪

Keywords: Coupled data assimilation, Atmosphere-Snow interaction, Snowfall

The method for identification of the cryoconite distribution by using satellite image

*安本 有唯¹、山家 大知¹、Bilesan Alireza²、今井 正堯⁶、川俣 大志^{1,3}、成瀬 延康⁴、竹内 望⁵、高橋 幸弘^{1,6}

*Ally YASUMOTO¹, Daichi Yamaga¹, Alireza Bilesan², Masataka Imai⁶, Hiroshi Kawamata^{1,3}, Nobuyasu Naruse⁴, Nozomu Takeuchi⁵, Yukihiko Takahashi^{1,6}

1. 北海道大学グローバルサイエンスキャンパス、2. 北海道大学大学院情報科学研究科、3. 北海道大学高等教育推進機構、4. 滋賀医科大学、5. 千葉大学大学院理学研究科、6. 北海道大学大学院理学研究院

1. Global Science Campus, Hokkaido University, 2. Graduate School of Information science and technology, Hokkaido University, 3. Institute for the Advancement of Higher Education, Hokkaido University, 4. Shiga University of Medical Science, 5. Graduate School of Science, Chiba University, 6. Graduate School of Science, Hokkaido University

Cryoconite, which is very small dark substance on glacier containing cyanobacteria and growing tangling with minerals, can absorb sunlight much, causing the faster melting than usual white snow. The valuable field survey enables us to study the cryoconite, albeit applying in the small area and with long interval. And a few remote sensing study uncover the glacier darkening and regression. Nevertheless no one knows how to point out its distribution over a wide area.

Here, according to luminance difference of between the glacier and the cryoconite, we propose new method with using multispectral bands of Landsat8 (resolution 30m), band2(450-515nm, B2) and band5(850-880nm, B5) to distinguish those two by making a new formula” $(B5-B2)/(B5+B2)$ ” (R). In the range of B2, the cryoconite's reflectance is about 10 %; otherwise the glacier has high percentage. In the range of B5, the cryoconite's one is about 20%; in contrast the glacier's almost half. The satellite images we use in 2016 July 30th, are analyzed since the cryoconite appeared widely and well in the period of 2016 late July to the beginning of August. We were successful in remove the shadow on this image by comparing R and RGB image. If a pixel in R is brighter than the other images, the place of the pixel should be cryoconite or sand.

The calculated index(R), in their spectra from previous study, applied that both of the cryoconite and the sand take a positive value while the glacier take a negative value in the pixel. The criterion will be shown in the presentation.

キーワード：クリオコナイト、リモートセンシング、氷河

Keywords: cryoconite, remote sensing, glacier

ひまわり8号熱赤外域バンドを用いたオホーツク海海氷面温度と薄氷域の抽出

Retrieval of ice surface temperature and thin ice area using thermal infrared bands of Himawari-8/AHI

*堀 雅裕¹、青木 輝夫²、谷川 朋範³、庭野 匡思³

*Masahiro Hori¹, Teruo Aoki², Tomonori Tanikawa³, Masashi Niwano³

1. 宇宙航空研究開発機構地球観測研究センター、2. 岡山大学、3. 気象研究所

1. Earth Observation Research Center, Japan Aerospace Exploration Agency, 2. Graduate School of Natural Science and Technology, Okayama University, 3. Meteorological Research Institute

Ice surface temperature (IST) has been an important observation target from space not only for calculating radiation budget but also for estimating the production of thin ice thickness in the cryosphere. In particular, the latter is important for assessing the amount of dense water with high salinity produced under newly formed thin ice. In this study we developed an algorithm for estimating IST and emissivity simultaneously using a semi-empirical emissivity model which can simulate the dependence of spectral emissivity on the surface snow/ice type and exitance angle. In this analysis we neglected the effect of water vapor absorption in the atmosphere and applied the algorithm to the data of AHI sensor onboard the Japanese geostationary satellite Himawari-8. Channel 13 (center wavelength: 10.4 μm) and 15 (12.4 μm) were used for the retrieval. The results show that emissivity as well as IST seemed to be successfully retrieved over the Okhotsk ice areas (but not validated with in-situ data). From the retrieved emissivity image, the area of thin sea ice such as nilas were easily determined. In addition, from the comparison with the case using a fixed emissivity for all snow and ice type, possible error in the IST retrieval with the fixed emissivity could be estimated to be up to a few Kelvin, which is due to the low emissivity of thin sea ice and the large viewing zenith angle of AHI around 60 degrees when observing the area of Okhotsk Sea. The same approach can also be applied to the data of polar orbit satellite such as the coming Japanese satellite mission “Global Change Observation Mission-Climate” (GCOM-C) to be launched in 2017.

キーワード：氷面温度、氷タイプ、射出率、リモートセンシング、AHI、SGLI

Keywords: Ice surface temperature, Ice type, Emissivity, Remote Sensing, AHI, SGLI

Inter-annual modulation of seasonal glacial velocity changes in the Eastern Karakorum detected by ALOS-1/2 data

*Muhammad Usman¹, Masato Furuya¹

1. Space Geodesy Research Laboratory, Department of Earth and Planetary Sciences, Hokkaido University, N10W8, Kita-ku, Sapporo 060-0810, JAPAN

Whereas the ice sheets all over the world are receding, the glaciers in Karakoram are either stagnant or advancing, which is known as 'Karakoram anomaly'. The surging dynamics and mass balance have been extensively studied in this area. However, in the Eastern Karakorum Range, the spatial and temporal changes in glacial velocity have been so far poorly understood. We have analyzed nearly all the available ALOS-1/2 data in this area and have examined the inter-annual modulation of five glaciers. The glaciers with size >30km, i.e. Siachen, Baltoro and Eastern tributary of Kundos, are mostly showing a considerable velocity change in their various parts, accompanying clear seasonal changes both in ALOS-1/2 data. However, this change mostly depends upon the individual glacier and is variable in space and time. On the other hand, the smaller glaciers (<30km), i.e. Singkhu, Gasherbrum and Western tributary of Kundos glaciers, are showing a slowdown in ALOS-2 data. Analysis of the local air surface temperature data at five observatories indicates that during the same season, the temperature trend in the study area is uneven and probably varies significantly between different glaciers. It can result in localized warming/cooling that can affect the availability of melt-water for an individual glacier. The excess surface melt-water at each individual glacier may undergo a variety of en/sub-glacial hydraulic and hydrological processes that are further different at each glacier. Thus, it will result in a complex velocity behavior in this region.

Keywords: Eastern Karakorum, Pixel offset, Glacier velocity, Inter-annual modulation

Measurement of mass balance with high altitude, and thermal property of debris-covered area at the Trambau Glacier, Nepal

*砂子 宗次郎¹、藤田 耕史¹、坂井 亜規子¹

*Sojiro Sunako¹, Koji Fujita¹, Akiko Sakai¹

1. 名古屋大学大学院環境学研究科

1. Graduate School of Environmental Studies, Nagoya University.

Shrinkage of Himalayan glaciers is unabated and thus contributes to sea level rise. In this region, in-situ measurements of mass balance of large glaciers have been conducted at few glaciers due to the difficult accessibility to their accumulation area. Moreover, many large glacier tongues are covered with debris, which makes the ablation process complicate. Thus, it is required to carry out in-situ measurement at debris-covered glaciers with accessible accumulation area.

We carried out in-situ measurements at Trambau glacier in the Nepal Himalaya pre- and post-monsoon seasons in 2016. We installed stakes network from ablation to accumulation area, and obtained direct mass balance data. An automatic weather station was set beside the glacier to obtain basic meteorological data during observation period. In order to establish a model for ablation at debris-covered ice, we also measured thermal conductivity and water content in debris layer, and set temperature sensors at different depth in the debris layer. Observed mass balance ranges from -2.62 m to +0.12 m w.e. a⁻¹ during the period. The maximum ablation is found at 5280 m a.s.l., which is the lower bound of debris-free area, and a liner relationship is found between mass balance and elevation ($r = 0.94$, $p < 0.01$). In contrast, no significant correlation is found between mass balance and elevation in debris-covered area whereas a coefficient correlation between mass balance and thermal resistance, which is a proxy of debris thickness, is 0.71 ($p < 0.05$).

キーワード：氷河、ヒマラヤ、質量収支、デブリカバー

Keywords: glacier, Himalaya, mass balance, debris cover

山岳氷河域研究のための全球スケール無償DEMの定量評価

Quantitative evaluation of global-scale free DEMs for mountain glaciology

*永井 裕人¹、田殿 武雄¹

*Hiroto Nagai¹, Takeo Tadono¹

1. 国立研究開発法人宇宙航空研究開発機構

1. Japan Aerospace Exploration Agency

航空機や人工衛星を用いた写真測量の原理による地形データの作成は、数値標高モデル（DEM: Digital Elevation Model）として近年まで様々な方法で実施され、数多くのデータセットが整備されてきた。特に人工衛星を用いたDEM作成は雪氷山岳地域において均一な地形情報を得る上で非常に重要であり、氷河研究の進展への影響も大きい。本発表では近年整備が進められてきた無償公開の全球スケールDEMを日本国内の起伏の異なる地形およびヒマラヤ氷河域で比較し、生じる差異を検証し、山岳氷河研究での有効性を議論する。

様々なDEMデータセットのうち、最近になって画素サイズ30 mのASTER GDEM, SRTM1, ALOS World 3D-30m (AW3D30) が整備された。これらの標高データについて国土地理院・基盤地図情報から三角点・測量点 (CP: Check Point) の標高値を取得し、精度評価を北アルプスの氷河地形、富士山裾野、利根川流域の沖積平野にて実施した。The

その結果、どの場所においてもAW3D30が地理院CPに近い値を示すことが分かった。DEM間でもっともばらつきが大きいのが北アルプスであり、ASTER GDEM, SRTM1について13-18 mほどCPより低く、ばらつき (CP値とDEM値の差分の標準偏差) は15-20 mとなった。起伏が緩やかな地形ほど、DEMによる精度の差異は小さくなり、利根川流域のほぼフラットな水田地帯ではSRTM1とAW3D30がほぼ等しい精度となった ($+0.1 \pm 3.1$ m)。ASTER GDEMにおいてもCP標高値との差の平均および標準偏差はともに5 m以下であった。このようにDEMが異なれば、地形によって精度が大きく異なることが示された。AW3D30が北アルプスの急峻地形においても最も高精度であるが、より平坦な地形ではSRTM1の精度も大きく改善されることがわかった。

DEM作成ではステレオペア画像間で対応点の取得が十分にできない部分で欠損が生じる。欠損そのものは有益ではないが、どこにどのように欠損が生じるかは、研究に最も適したDEMを選ぶ際に重要な知見となる。そこでネパールヒマラヤにおいて、上記三種DEMの欠損がどのように生じているかを検証した。AW3D30の欠損はヒマラヤ主脈周辺で多く見られた。これはALOS衛星の光学画像によって取得されたため、山頂付近・氷河涵養域では積雪によって上手く対応点をマッチングできないことが原因と考えられる。SRTM1では、より低標高で斜度45°付近に最も欠損が集中していることが分かった。軌道横方向斜め下にレーダを照射して地形計測を行うため、急峻な山に妨げられるシャドーイングやレイオーバー (近くの谷より遠くの山の方が先に電波を反射することに起因する受信順序の反転) が原因と考えられる。一方、最も長期にわたる観測データから作成されたASTER GDEMでは対象地域内に欠損は見られなかった。精度は高くないが、集水域の作成など、抜け目のないデータが必要となる際に有用なDEMであると言える。

キーワード : SRTM, GDEM, AW3D

Keywords: SRTM, GDEM, AW3D

InitMIP-Greenland experiments with the ice sheet model SICOPOLIS

*Ralf Greve¹, Reinhard Calov²

1. Institute of Low Temperature Science, Hokkaido University, 2. Potsdam Institute for Climate Impact Research (PIK Potsdam)

The Ice Sheet Model Intercomparison Project for CMIP6 (ISMIP6) brings together a consortium of international ice sheet and climate models to explore the contribution from the Greenland and Antarctic ice sheets to future sea level rise. For such projections, initialisations are required that provide initial states of the respective ice sheet. Therefore, as one of the first initiatives within ISMIP6, InitMIP-Greenland was launched in order to explore this issue for the Greenland ice sheet across a variety of models and initialisation techniques. Two different initialisation techniques are common, namely spin-up methods (paleoclimatic simulations until the present) and assimilation methods (assimilation of observations of the present-day ice sheet). We contribute to InitMIP-Greenland with the ice sheet model SICOPOLIS and two different spin-up techniques, (1) a SeaRISE-legacy spin-up over 125 ka with essentially fixed topography, and (2) a new spin-up over 135 ka with freely evolving topography. New methods applied for spin-up (2) are monthly-mean (rather than mean annual) input data for the present-day precipitation, a sub-grid-scale ice discharge parameterisation and an iterative correction of the present-day precipitation based on the misfit between the simulated and observed present-day ice thickness. The agreement between simulated and observed ice topography is naturally better for the fixed-topography case (1) than for the freely evolving case (2). Both spin-ups produce a realistic distribution of the surface velocity, including the major ice streams and outlet glaciers (at 5 km horizontal resolution). InitMIP-Greenland also comprises two future climate scenarios, ctrl (present-day climate over 100 a) and asmb (prescribed schematic surface mass balance anomaly over 100 a due to global warming), both to be run with freely evolving ice topography. The response of the ice sheet (mass loss) to the asmb forcing is, in absolute terms, ~50% larger for spin-up (2) than for spin-up (1), and relative to the respective control run ctrl even ~85% larger. This demonstrates impressively that, even with the same ice sheet model, different initialisation methods can lead to a major spread of results of future climate experiments.

Keywords: Greenland, Ice sheet, Climate change, Sea level rise, Modelling


The Effects of H₂SO₄ on the Flow and Fabric of Polycrystalline Ice

*Kevin Hammonds¹, Ian Baker²

1. Montana State University, 2. Dartmouth College

It is well established that the Earth's large continental ice sheets contain a variety of naturally occurring impurities, both soluble and insoluble. Understanding how these impurities affect the rheology, intrinsic thermodynamic properties, and ultimate fate of these ice sheets is much less understood. Previous work has shown that H₂SO₄ dramatically reduces the strength and increases the ductility of single crystal ice, but its effects on polycrystalline ice are unknown. In order to investigate the effects that trace amounts of H₂SO₄ have on the flow and ductility of polycrystalline ice a series of mechanical tests were conducted at -6°C, -10°C, -12°C, and -20°C using laboratory-prepared ice with a mean grain diameter of 1 mm and doped with 1-10 ppm of H₂SO₄. Parallel tests were performed on identical, but undoped polycrystalline ice. Mechanical testing included uniaxial tensile creep tests at a constant load of 38 kg (0.75 MPa initial stress) and uniaxial compression tests at constant strain rates ranging from $1 \times 10^{-6} \text{ s}^{-1}$ to $1 \times 10^{-4} \text{ s}^{-1}$. The tensile tests showed that H₂SO₄-doped specimens exhibited faster creep rates than undoped ice, while the compression tests demonstrated that H₂SO₄-doped specimens exhibit a significantly lower peak stress than undoped ice. Post-mortem microstructural analyses were performed using cross-polarized light thin section imaging, X-ray computed microtomography, Raman spectroscopy, and electron backscatter diffraction. These analyses showed that H₂SO₄-doped specimens had a much larger grain size at strains 15%, and an earlier onset of micro-cracking at lower strain rates than the undoped ice. Further, a liquid-like phase containing H₂SO₄ appears to be present at the grain boundaries of the H₂SO₄ doped ice.

Keywords: Ice, Sulfuric , Acid, Microstructure, Rheology



Estimation of glacier velocities at Pio XI in the Southern Patagonia Icefield detected by ALOS-2/PALSAR-2

*波多 俊太郎¹、古屋 正人¹

*Shuntaro Hata¹, Masato Furuya¹

1. 北海道大学理学部地球惑星科学科

1. Hokkaido University School of Science Department of Earth and Planetary Sciences

The Southern Patagonia Icefield(SPI) is the largest temperature ice mass in the southern hemisphere. Recently, it is reported that majority glaciers in the SPI had undergone significant retreat and thinning. On the contrary to this trend, only Pio XI glacier advanced in the SPI in the last century. Pio XI glacier, the largest glacier in the SPI, has two termini. The north tongue calves into the Greve lake, whereas south tongue into the Eyre Fjord. There are two interesting reports on the behaviours of Pio XI. One is about whether Pio XI is surge-type glacier, another is about switch the flow path of Pio XI.

In order to understand the dynamics of Pio XI glacier, this paper reports the temporal variations of flow velocities and terminus positions by using Advanced Land Observation Satellite-2/Phased Array-type L-band Synthetic Aperture Radar-2(ALOS-2/PALSAR-2)data acquired in 2015.

In 2015, Pio XI flows < 4m/d from February to June. However in the term from August to September, Pio XI revealed significant acceleration. In 2015, the primary flow of Pio XI is terminate into Greve lake from north tongue.

キーワード : 氷河、パタゴニア

Keywords: Glacier, Patagonia

Uncertainties in ice-sheet simulation due to a variation in the numerical schemes of the ice transport equations

*齋藤 冬樹¹、阿部 彩子^{2,1}

*Fuyuki SAITO¹, Ayako Abe-Ouchi^{2,1}

1. 国立研究開発法人海洋研究開発機構、2. 東京大学大気海洋研究所

1. Japan Agency for Marine-Earth Science and Technology, 2. Atmosphere Ocean Research Institute, University of Tokyo

Generally the evolution of ice-sheet thickness in an ice-sheet model is formulated using the divergence term of horizontal ice flux and the mass balance terms at the upper and lower surfaces, derived from the continuity equation with the assumption of incompressible fluid (i.e., a non-linear transport equation). There are many variation of the formulation, which differs in numerical aspects such as stability, accuracy, numerical diffusivity, conservation, complexity, computational costs and so on.

Since ice-sheet thickness near grounding line is relatively steep, simulated evolution of thickness over this region is expected to be much influenced by numerical diffusion and/or oscillation caused by characteristics of numerical schemes to represent the transport equation. Often the evolution near the grounding line is a dominant aspect for large-scale ice-sheet evolution, the uncertainties due to the numerical characteristics should be evaluated.

In this study implementation of a variation of Constrained Interpolation Profile (CIP) in a numerical ice-sheet model ICIES is reported. Simulation under configuration of past ice-sheet model intercomparison experiments (e.g., EISMINT, ISMIP) is reported, comparing to those using typical schemes such as an upwind scheme and/or diffusion-type scheme.

キーワード：氷床、数値スキーム

Keywords: ice-sheet, numerical schemes

Lansat 8 による赤雪分布の再評価

Revaluation with Landsat 8 of red snow mapping in ice field

*畠山 栞¹、長谷川 佳佑¹、田代 達也¹、李 肖陽²、Roshanianafard Ali³、川俣 大志^{1,4}、成瀬 延康⁵、竹内 望⁶、高橋 幸弘^{1,7}

*Shiori Hatakeyama¹, Keisuke Hasegawa¹, Tatsuya Tashiro¹, Xiaoyang Li², Ali Roshanianafard³, Hiroshi Kawamata^{1,4}, Nobuyasu Naruse⁵, Nozomu Takeuchi⁶, Yukihiro Takahashi^{1,7}


1. 北海道大学グローバルサイエンスキャンパス、2. 北海道大学大学院環境科学院、3. 北海道大学大学院農学院、4. 北海道大学高等教育推進機構、5. 滋賀医科大学、6. 千葉大学大学院理学研究科、7. 北海道大学大学院理学研究院

1. Global Science Campus, Hokkaido University, 2. Graduate School of Environmental Science, Hokkaido University, 3. Graduate School of Agriculture, Hokkaido University, 4. Institute for the Advancement of Higher Education, Hokkaido University, 5. Shiga University of Medical Science, 6. Graduate School of Science, Chiba University, 7. Graduate School of Science, Hokkaido University

The melting of glaciers has various effects on the environment such as sea level rise and glacial lake collapse. The global warming is commonly believed to cause the melting, while some scientists insist that the breeding of snow creatures on the glacier contributes mainly. The colorful phenomenon such as red snow, where microorganisms breed on the surface of the glaciers, makes the absorption of heat from sunlight increase, resulting in the faster melting rate of the glacier than usual. However, the actual amount of melting in the glacier is unknown. In order to elucidate it, the distribution of the coloring phenomenon on the glacier should be investigated. Therefore, we aimed at observing glaciers using remote sensing which can observe coloring phenomenon extensively and easily. Using remote sensing the red snow has been observed twice (2006, 2015). However, around 19 percent area in the ice field was not observed due to the saturation of the conventional 8-bit image sensor of the satellite. In this study Landsat 8 with the saturation-free 16-bit image sensors (band 2 and 4) is adopted and Harding ice field in June- August 2015 was selected for our observation with the information from previous studies. As shown in figure, we achieve to observe the whole area without the sensor saturating and the red snow appears in the larger area than before. The detail evaluation will be shown in the presentation.

キーワード：ランドサット、赤雪、氷原、リモートセンシング

Keywords: Landsat, Red snow, Ice field, Remote sensing



ALOS1/2を用いたInSAR解析による永久凍土地帯の地盤変動検出：Herschel島（カナダ）とBatagaika crater（シベリア）の事例

GROUND DEFORMATION MAPPING BY ALOS1/2 INSAR: CASE STUDIES AT HERSCHEL ISLAND, CANADA, AND BATAGAIKA CRATER, SIBERIA

*柳谷 一輝¹、古屋 正人¹

*Kazuki Yanagiya¹, Masato Furuya¹

1. 北海道大学大学院理学院

1. Graduate School of Science, Hokkaido University

永久凍土は北半球において四分の一の面積を占め、その融解や凍結により地盤の沈下や隆起が生じる。これらの地盤変動は人口建築物の損壊や変形をもたらすものとして、ロシアなどの北極海沿岸地域において古くから地域に密着した問題として研究されてきた。しかし、定量的な観測データは皆無であり、またその時空間分布などは殆ど調査されていない。一方、近年地球温暖化問題が提起され、温室効果ガスの推移について多くの研究がされている中で、広大な凍土に貯蔵されている有機炭素の重要性が指摘されている。北半球のうち永久凍土地帯が占める面積は広大であり、実際に凍土の融解により形成されるサーモカルスト湖ではメタンガスの放出が確認されている。永久凍土の融解は地球環境を考える上で無視できない変動であり、高緯度地域に限られた問題ではない。一方で永久凍土そのものの観測を広域で行うことは非常に難しい。その代わりに、永久凍土地帯の地盤の沈下や隆起の広域モニターは技術的には可能であり、それができれば凍土の融解過程の監視に重要な貢献ができる可能性がある。

本研究では合成開口レーダーを用いたリモートセンシングによる永久凍土地帯の地盤変動観測、特に融解による地盤沈下の検出を試みた。JAXAによって打ち上げられた2つのSAR衛星ALOSとALOS2のデータを用いて、2007年から2016年の最近の地盤変動を観測した。観測対象としては変動速度が比較的速いスランプ地形に着目し、カナダのハーシェル島とロシアのバタガイカクレーターにおいて地盤沈下を検出した。ハーシェル島では海岸部の沈降と後退を先行研究では調べられていないALOS2による近年の変動を観測した。一方永久凍土起源のクレーターとして知られるバタガイカクレーターでは、内陸部にできたメガスランプで変動のシグナルを検出したが、解析に用いたDEMによる影響が結果に生じている。

キーワード：ALOS、InSAR、永久凍土、サーモカルスト、リモートセンシング

Keywords: ALOS, InSAR, permafrost, thermokarst, remote sensing