

## Underestimation characteristics of TRMM 2A25 V7 near surface rain over and around the Meghalaya Plateau

\*徹 寺尾<sup>1</sup>、村田 文絵<sup>2</sup>、山根 悠介<sup>3</sup>、木口 雅司<sup>4</sup>、福島 あずさ<sup>5</sup>、田上 雅浩<sup>6</sup>、林 泰一<sup>7</sup>

\*Toru Terao<sup>1</sup>, Fumie Murata<sup>2</sup>, Yusuke Yamane<sup>3</sup>, Masashi Kiguchi<sup>4</sup>, Azusa Fukushima<sup>5</sup>, Masahiro Tanoue<sup>6</sup>, Taiichi Hayashi<sup>7</sup>

1. 香川大学教育学部、2. 高知大学理学部、3. 常葉大学教育学部、4. 東京大学生産研、5. 神戸学院大学人文学部、6. 東京大学大学院工学系研究科、7. 京都大学東南アジア研究所

1. Faculty of Education, Kagawa University, 2. Faculty of Science, Kochi University, 3. Faculty of Education, Tokoha Univesity, 4. Insitute of Industrial Science, The University of Tokyo, 5. Faculty of Humanity, Kobe Gakuin University, 6. School of Engineering, The University of Tokyo, 7. Center for Southeast Asian Research, Kyoto University

Utilizing our original tipping bucket raingauge network over Bangladesh and northeast India (Fig. 1), we have detected underestimation of the near surface rain in TRMM 2A25 V7 dataset over and around the Meghalaya Plateau. This underestimation was prominent especially in the monsoon season. Such underestimation of TRMM PR sensor was detected in other mountaneous areas also (Prat and Barros 2010; Wilson and Barros 2014; Terao et al. 2017).

In the present study, we further analyzed the characteristics of underestimation utilizing TRMM 2A25 V7 and raingauge dataset. In TRMM 2A25 dataset, rain type is defined for each ray to distinguish stratiform and convective rainfalls. We evaluated the contribution to the underestimation from different types of rainfall (Figs. 2 and 3) for two regions, Meghalaya and Sylhet-Barak areas, with different orographic situation. In these figures, we evaluated the averaged contribution ratio and their 95 % confidence intervals for each rain type. These confidence intervals were evaluated by the boot-strap method. The Meghalaya area is the hill area in India, and Sylhet-Barak area is the northeastern part of the Bengal Plain, which consists of both Sylhet Division in Bangladesh and Barak Basin in Assam, India.

Figure 2 shows that the underestimation has been highly contributed by stratiform rain over the Meghalaya Plateau. Averaged for the stratiform rain cases, rainfall intensity detected by raingauges was greater than  $6 \text{ mm h}^{-1}$ , with more than 50 % negative bias ratio. Most notable result was the high contribution of the no-rain detected cases. We detected tipping of raingauge more than 5 % out of TRMM 2A25 no-rain detected cases. For other areas, this ratio was much less than 0.5 %. Over the Sylhet-Barak area, underestimation was explained mainly only by the convective rain (Fig. 3). Thus, although the areas of underestimation in TRMM PR sensor were geographically adjacent to each other, the cause of the underestimation was largely different.

For near nadir cases, the clutter free bottom height (CFB height) of the ray tends to be lower. Therefore, we checked the impact of the angle of ray to check its impact on the underestimation (not shown), but we found no clear tendency.

キーワード : TRMM、降雨レーダー、過小評価、インド亜大陸北東部

Keywords: TRMM, PR, underestimation, northeastern Indian subcontinent




Fig. 1: Raingauge network




Fig. 2




Fig. 3

## チャオプラヤ流域の土地利用変化が陸域水収支に及ぼす影響

### Impact of land-use change on terrestrial water balance in the Chao Phraya River Basin

\*高田 久美子<sup>1</sup>、花崎 直太<sup>1</sup>

\*Kumiko TAKATA<sup>1</sup>, Naota Hanasaki<sup>1</sup>

1. 国立環境研究所 地球環境研究センター

1. Center for Global Environmental Research, National Institute for Environmental Studies

チャオプラヤ川上流域の山岳地帯では、近年、森林伐採が進んでおり、これによる洪水の増加が懸念されている。これに対し、植林を進めることによって土壌の保水能力を高め、流出応答をゆっくりにする効果で洪水を抑制できるのではないかと期待がある。一方で、実際に植林を進めようとした場合、陸域水収支や水資源・水利用の観点に加えて、収入源となる耕作地の減少によって社会経済的な影響なども与える可能性があり、慎重な影響評価が必要である。これまで当地域では、ダムなど人間活動による水利用を考慮した全球水資源モデルH08 (Hanasaki et al., 2008ab) による水資源と陸域水収支の評価が行われているが、陸面水文過程にはバケツモデルが用いられており、土地利用変化(植生変化)を陽に扱うことができなかった。そこで本研究では、植生を陽に扱っている陸面水文過程モデルMATSIRO (Takata et al., 2003; Nitta et al., 2014) を用いて、チャオプラヤ川流域における植生変化が陸域水収支に及ぼす影響について調べる。具体的に、MATSIROに地上気象データを与えて、植生分布を変えた数値実験を行い、土地利用変化の影響を評価する。予備実験として、水平解像度約100kmで、1979年1月1日から2007年12月31日の全球地上気象データ (Kim et al., 2009) を与え、現在の植生分布(図)を与えた実験(CTL)と、自然植生(常緑広葉樹林、混合林、草地)を耕作地に変えた実験(CROP)を行い、チャオプラヤ流域での河川流量の変化を調べた。

流域北西部のBhumibolダムや流域北東部のSirikitダムに流入する河川流量の観測値と数値実験の河川流量を比較したところ、流域北西部(Bhumibolダム)では7月から10月に流量が増大する季節変化は定性的に再現されていたものの、5月-6月の降雨に対応した流出が表現されなかった。また、流域北東部(Sirikitダム)では流出の起こる期間が8月に集中し、7月から徐々に増加して9月に徐々に減少する様子が再現されなかった。さらに、年間の河川流量はどちらも観測の数分の1に過小評価されていた。土地利用変化に対する応答について、常緑広葉樹林から耕作地に転換したチャオプラヤ流域北西部では、森林のほうが耕地よりも河川流出の開始時期が遅く、また年間の河川流量が小さくなり、森林による洪水の緩和を示唆する結果となった。一方で、混合林から耕作地に転換したチャオプラヤ流域北東部では、森林と耕地の場合で河川流出の開始時期はほとんど変わらず、年間の河川流量は森林のほうが大きかった。

今後、チャオプラヤ流域における水平解像度約10kmの気象データ (Kotsuki et al., 2013) を用いた高解像の数値実験を行い、観測された河川流量変化や年間流量の再現性を向上するとともに、植生タイプによって耕地化した時に応答が異なる原因や、土地利用変化が陸域水収支と河川流量に及ぼす影響についてメカニズム解明と定量評価を進める。

謝辞：東京大学の新田友子博士と理化学研究所の小槻峻司博士には、MATSIROオフライン実験の実行に当たってご協力いただいた。本研究は科学技術振興機構(JST)と国際協力機構(JICA)による地球規模課題対応国際科学技術協力プログラム (SATREPS) の支援を受けて実施している。

#### 参考文献：

- Hanasaki, N., et al., Hydrology and Earth System Sciences, 12, 1007-1025, 2008a.
- Hanasaki, N., et al., Hydrology and Earth System Sciences, 12, 1027-1037, 2008b.
- Takata, K., et al., Global and Planetary Change, 38, 209-222, 2003.
- Nitta, T., et al., J. Clim., 27(9), pp3318-3330, 2014.

Kim, H., et al., Res. Lett., 36, L17402, 2009.

Kotsuki, S., et al., Hydrolog. Res. Lett., 7, 7984, 2013.

キーワード：土地利用変化、陸域水収支、チャオプラヤ流域

Keywords: Land-use change, Terrestrial water balance, Chao Phraya river basin




Figure Land-use distribution used in the preliminary experiment. Black lines indicate river route of the Chao Phraya river. Boxes (orange and green) are the grid points compared with observed river discharge.

## An analysis of the atmospheric circulation around the Tibetan Plateau revealed by the stable isotope in precipitation—A case study of GEWEX-GAME/Tibet in 1998

黒崎 豊<sup>1</sup>、\*谷田貝 亜紀代<sup>2</sup>、杉本 敦子<sup>3</sup>

Yutaka Kurosaki<sup>1</sup>, \*Akiyo Yatagai<sup>2</sup>, Atsuko Sugimoto<sup>3</sup>

1. 弘前大学 理工学部、2. 弘前大学 大学院 理工学研究科、3. 北海道大学 北極域研究センター

1. Faculty Science and Technology, Hirosaki University, 2. Graduate School of Science and Technology, Hirosaki University, 3. Arctic Research Center, Hokkaido University

Data of stable isotopes obtained from glaciers and tree rings on the Tibetan Plateau are useful in restricting the paleoclimate. It is important to meteorologically analyze stable isotopes in precipitation over the Tibetan Plateau, which are affected by complicated atmospheric circulation processes, because the ratio of stable isotopes in precipitation based on the transport process is affected by atmospheric circulation. However, this approach has not yet been well implemented.

Data of temporally and spatially stable isotopes in precipitation were obtained over the Tibetan Plateau and Nepal during a field campaign of the Global Energy and Water Experiment Asian Monsoon Experiment/Tibet in 1998. The data reveal a relationship between stable isotopes in precipitation over the Tibetan Plateau and active/break variations of the Indian monsoon.

During a break phase, low  $\delta^{18}\text{O}$  values and low d-excess values were observed at all observational sites. Transportation in this phase was an upslope process in which an air parcel gains altitude near the Himalayas. This trend can be explained by air parcels crossing over the Himalayas.

During an active phase, a characteristic trend of stable isotopes in precipitation over Tibetan Plateau was observed. Low  $\delta^{18}\text{O}$  and low d-excess values were observed around the south of the Tibetan Plateau (hereinafter called region 1) while high  $\delta^{18}\text{O}$  and high d-excess values were observed around the north of the Tibetan Plateau (hereinafter called region 2). The phase of region 1 coincided with the break phase, and transport might be an upslope process. However, the phase of region 2 was different because of the inland effect. To interpret the high  $\delta^{18}\text{O}$  values, we used the forward trajectory from convective cloud over central India, and examined the top height of convective cloud around region 2 over the Tibetan Plateau using measurements made by the precipitation radar onboard the Tropical Rainfall Measuring Mission satellite. Results showed that air parcels at an altitude exceeding 8,000 m in convective cloud around central India were transported to the Tibetan Plateau, and high  $\delta^{18}\text{O}$  values between 8,000 and 10,000 m in convective cloud around central India might be associated with precipitation around region 2 over the Tibetan Plateau.

To interpret the characteristics of stable isotopes in precipitation around the Tibetan Plateau, it is important to consider the active/break phase and trajectory of air parcels of the Indian monsoon. Clarifying the vertical distribution of stable isotopes in precipitation in convective cloud can improve our knowledge of the paleoclimate and help determine an isotope model in future work.

キーワード：降水安定同位体、チベット高原、大気循環

Keywords: stable isotope in precipitation, the Tibetan Plateau, atmospheric circulation

## Heavy rain prediction applying satellite-based cloud data assimilation over land

\*Seto Rie<sup>1</sup>、Koike Toshio<sup>2</sup>

\*Rie Seto<sup>1</sup>, Toshio Koike<sup>2</sup>

1. 東京工業大学、2. 水災害・リスクマネジメント国際センター

1. Tokyo Institute of Technology, 2. International Centre for Water Hazard and Risk Management (ICHARM)

For accurate flood prediction, warning systems, and optimized dam control, information of positional relationship between rain areas and river basins is crucial. This requires very fine precision in the prediction of rainfall areas. Assimilation of satellite-based microwave observation of cloud has great potential to improve precipitation areas because it can directly obtain information on rainfall locations as well as amount of cloud. However, it is difficult to observe clouds over land using satellite microwave remote sensing, because land emissivity is much stronger and more heterogeneous than that of cloud. To overcome this challenge, appropriate representation of heterogeneous land emissivity is needed. Thus, We developed a Coupled Atmosphere and Land Data Assimilation System with the Weather Research and Forecasting model (CALDAS-WRF), which can assimilate soil moisture, cloud water content over land, and heat and moisture within clouds simultaneously. Results of application of CALDAS-WRF to heavy rain events show that the system effectively assimilated cloud signals and produced very accurate cloud and precipitation distributions with appropriate intensity. Also, the local atmospheric fields are modified appropriately around the area of assimilated clouds. Furthermore, by using operationally analyzed dynamical and moisture fields as initial and boundary conditions, the system improved prediction of precipitation duration. The results demonstrate the method's promise in dramatically improving predictions of heavy rain and consequent flooding.

キーワード：雲データ同化、リモートセンシング、豪雨予測

Keywords: cloud assimilation, remote sensing, heavy rain prediction

## Regional seasonal marches of precipitation and their long-term variations in India for 1901-2013

\*井上 知栄<sup>1</sup>、松本 淳<sup>1,2</sup>

\*Tomoshige Inoue<sup>1</sup>, Jun Matsumoto<sup>1,2</sup>

1. 首都大学東京 都市環境学部 地理環境コース、2. 海洋研究開発機構 大気海洋相互作用研究分野

1. Department of Geography, Tokyo Metropolitan University, 2. Department of Coupled Ocean-Atmosphere-Land Processes Research (DCOP), JAMSTEC

Regional characteristics of climatological seasonal marches of precipitation and their long-term variations have been examined for the period 1901-2013 in India using a high resolution ( $0.25^{\circ} \times 0.25^{\circ}$ ) daily gridded precipitation dataset provided by the Indian Meteorological Department.

Cluster analysis (Ward's method) was applied for the 30-year climatological 5-day precipitation (1981-2010) at each grid box, and nine regions were divided. Then changes of seasonal precipitation characteristics, including onset, peak, and retreat of rainy season were examined for the 113-year period from 1901-2013 in a regional basis. As a result, for example, in the west coast area where typical monsoonal seasonal changes are observed, a prominent precipitation peak appeared in July prior to 1940, while precipitation in the subsequent rainy season in August has increased during the 20th century, and the degree of concentration of precipitation in July has decreased after the 1940s.

キーワード：インド、降水量、気候変動

Keywords: India, precipitation, climatic variation

## Predictable and unpredictable monsoons

## Predictable and unpredictable monsoons

\*山中 大学<sup>1</sup>

\*Manabu D. Yamanaka<sup>1</sup>

1. 海洋研究開発機構大気海洋相互作用研究分野／神戸大学名誉教授

1. Department of Coupled Ocean-Atmosphere-Land Processes Research, Japan Agency for Marine-Earth Science and Technology; Professor Emeritus of Kobe University

MAHASRI renewed GAME both in scientific and non-scientific aspects of monsoon-related issues. The infrastructures for hydro-climatological observations have been improved by economic growth and public needs. Now we can distinguish the predictable and unpredictable parts of monsoons, and must do it. The monsoonal or rainy-seasonal cycles are originated from the insolation varying astronomically with season and latitude, and are amplified geographically mainly with the solid (land) - liquid (sea) heat capacity contrast on the Earth's surface. The former astronomical process is completely predictable as described in a classical agricultural calendar which might have been changed gradually by the recent global warming. In addition, most of geographical variabilities of Asia monsoon have been revealed by GAME-MAHASRI periods. At first we must perfect to archive them, to let everybody know them, and to apply them for agricultural application, disaster prevention, and so on. Those parts might be still supported by so-called developed countries including outside of the monsoonal Asia, but are not so many in my view. At last we will recognize the truly unpredictable parts, which may be varying dynamically or even indeterministically due to nonlinear or multivariate processes. These remainder parts are so truly pioneering/challenging and/or domestic that any advanced countries can never support them. In this meaning now we are entering into indeed a qualitatively new era.

キーワード : monsoon、science and society、international collaboration

Keywords: monsoon, science and society, international collaboration

# Dynamics of changing impacts of tropical Indo-Pacific variability on Indian and Australian rainfall

\*Ziguang Li<sup>1</sup>, Xiaopei Lin<sup>1</sup>, wenju Cai<sup>2</sup>

1. Ocean Univ. of China, 2. CSIRO

A positive Indian Ocean Dipole (IOD) and a warm phase of the El Niño-Southern Oscillation (ENSO) reduce rainfall over the Indian subcontinent and southern Australia. However, since the 1980s, El Niño's influence has been decreasing, accompanied by a strengthening in the IOD's influence on southern Australia but a reversal in the IOD's influence on the Indian subcontinent. The dynamics are not fully understood. Here we show that a post-1980 weakening in the ENSO-IOD coherence plays a key role. During the pre-1980 high coherence, ENSO drives both the IOD and regional rainfall, and the IOD's influence cannot manifest itself. During the post-1980 weak coherence, a positive IOD leads to increased Indian rainfall, offsetting the impact from El Niño. Likewise, the post-1980 weak ENSO-IOD coherence means that El Niño's pathway for influencing southern Australia cannot fully operate, and as positive IOD becomes more independent and more frequent during this period, its influence on southern Australia rainfall strengthens. There is no evidence to support that greenhouse warming plays a part in these decadal fluctuations.

Keywords: Monsoon, ENSO, IOD

## Dominant synoptic disturbance in the extreme rainfall at Cherrapunji, northeast India, based on 104 years of rainfall data (1902-2005)

\*Fumie Murata<sup>1</sup>, Toru Terao<sup>2</sup>, Hatsuki Fujinami<sup>3</sup>, Taiichi Hayashi<sup>4</sup>, Haruhisa Asada<sup>5</sup>, Jun Matsumoto<sup>6</sup>, Hiambok Jones Syiemlieh<sup>7</sup>

1. Faculty of Science, Kochi University, 2. Faculty of Education, Kagawa University, 3. Institute for Space-Earth Environmental Research, Nagoya University, 4. Center for Southeast Asian Studies, Kyoto University, 5. Nara Women's University, 6. Tokyo Metropolitan University, 7. Department of Geography, North-Eastern Hill University

The characteristics of active rainfall spells (ARSs) at Cherrapunji, northeast India, where extreme rainfall is experienced, and their relationships with large-scale dynamics were studied using daily rainfall data from 1902 to 2005 and Japanese 55-year reanalysis from 1958 to 2005. Extreme daily rainfalls occur in association with ARSs. The extremely large rainfall amounts in the monsoon season are decided by a cumulative rainfall during ARSs. ARSs start when anomalous anticyclonic circulation (AAC) at 850-hPa propagates westward from the South China Sea and western North Pacific, and covers the northern Bay of Bengal. The AAC propagates further westward and suppresses convection over central India during ARSs at Cherrapunji, and continues for 3 to 14 days. Consequently, a northward shift of the monsoon trough during the 'break' in the Indian core region occurs. The westerly wind, which prevails in the northern portion of the AAC, transports moisture toward northeast India and enhances moisture convergence over northeast India with southerly moisture transport from the Bay of Bengal, and greatly intensifies the orographic rainfall. In the upper troposphere, the Tibetan high tends to extend southward with the onset of ARSs. A linear relationship can be seen between the length and total rainfall of an ARS. Longer ARSs tends to result in greater total rainfall. AACs with a greater zonal scale tend to produce longer and more intense ARSs. This study provide a certain evidence for the effect of AACs in the western North Pacific on the Indian summer monsoon.

Keywords: extreme rainfall , Indian summer monsoon, intraseasonal variability, orographic rainfall

# Characteristics of the Rainfall over Luzon during the Summer Monsoon of the Philippines

\*Lyndon Mark Payanay Olaguera<sup>1</sup>, Jun Matsumoto<sup>1</sup>

1. Tokyo Metropolitan University

The changes in rainfall over Luzon Island during the summer monsoon of the Philippines are investigated using averaged pentad rainfall data from 1981-2010. A monsoon break (P32 to P34; Jun. 5 to Jun. 19) after the climatological onset (P29, May 21 to May 25) was identified. The break is associated with the southwestward extension of the subtropical high during the seasonal evolution of the Western North Pacific Monsoon (WNPM). The break is obvious in stations located over the north and central Luzon. The average rainfall distribution reveals the impact of intra-seasonal oscillations in the summer monsoon rainfall of the Philippines.

Keywords: summer monsoon, Philippines, monsoon break, Intra-seasonal Oscillation




Figure 3 Rainfall distribution from a). CMAP data set averaged from 119E to 121E, and b). averaged across the nine PAGASA stations. Shaded contours are for rainfall greater than 5 mm/day. The dashed line is the rainy seasonal mean (12.52mm/day) averaged from P29 (onset) to P67 (withdrawal).


## Records of the East Asian Monsoon from Beibu Gulf since Last Deglaciation

\*Dong Xu<sup>1</sup>, Liming Ye<sup>1</sup>, Xibin Han<sup>1</sup>

1. Second Institute of Oceanography, SOA, China

The East Asian Monsoon is one of the most active components in the global climate system. Many researchers have found that the intensity of East Asian Summer Monsoon (EASM) quickly increased in the early Holocene, characterized by strengthened precipitation and humid climate, while gradually decreased during the middle and late Holocene, characterized by decreased precipitation and dry climate. But due to the difficulty of constructing substitute indexes, the evolution of East Asian Winter Monsoon (EAWM) through Holocene was debatable. In this paper we studied the East Asian Monsoon records in core B106 recovered from the Beibu Gulf (the Gulf of Tonkin) in northwestern South China Sea. The 300cm long core was located in 108°36'02"E, 19°54'04"N with a water depth of 62 m, and the grain size, chemical and clay mineral component, and AMS<sup>14</sup>C age of this core was analyzed. The results show that before 13ka BP, the location of B106 was in continental sedimentation environment, with sedimentation source mainly from south China paleo-rivers, and the sediments were characterized by coarser grain size and high  $\Sigma \text{HREE}/(\Sigma \text{LREE} + \Sigma \text{MREE})$ ,  $\text{TiO}_2/\text{Al}_2\text{O}_3$  and C/N. From about 13 ka BP to 7ka BP, B106 located in marine environment with increasing water depth and fading influence of south China paleo-rivers. Since the formation of loop current in Beibu Gulf at about 7ka BP, the sediments of core B106 came from mixed sources but with little contribution of Red River or Pearl River. The fluctuation of CIA, Rb/Sr, (I/S+I)/K and the content and mean grain size of less than 16  $\mu\text{m}$  grain size population of core B106 were influenced by the evolution of East Asian monsoon. By synthesizing these indexes, the warm and humid climate in early Holocene was revealed, and also the cold event around 8.2, 5.4 and 4.8ka BP in middle Holocene, the drought climate during 3.6-1.6ka BP in late Holocene were revealed, reflecting a regional response to global climate change. But more attention should be paid in using CIA and mean grain size of fine environmental sensitive population as the substitution indexes of EAWM intensity, as they may be influenced by EASM too. In middle and late Holocene, the content of less than 16  $\mu\text{m}$  grain size population of core B106 can reflect the strength of EAWM to some extent, while the mean grain size of less than 16  $\mu\text{m}$  population was insensitive to EAWM and the increasing of which was closely linked with the weakening of EASM. When extracting grain size index to study the evolution of East Asian Monsoon, the restriction of sedimentation source and dynamics should be defined firstly.

Keywords: Beibu Gulf, East Asian Monsoon, Last Deglaciation



(a) CIA records of core B106(this study), S20 (Xu et al., 2010) and MZ01 (Liu et al., 2010a); (b) Rb/Sr of B106; (c) clay mineral proxy-(illite-smectite interstratified clay minerals + illite) vs. kaoline of B106 (Wu et al., 2011); (d) content and mean grain-size (e) of fine grain-size population (<16µm) of B106; (f) fine grain-size population (1-2.2µm) content of core KNG5 (Huang et al., 2011); (g) content and mean grain size (h) of fine grain-size population (<28µm) of core PC-6 (Xiao et al., 2006); (i) records of the stalagmite  $\delta^{18}\text{O}$  from Dongge Cave, South China (Dykoski et al., 2005); (j) relative abundance ratio of two diatom species (*A. granulata* vs. *C. stelligera*) recorded in Huguang Marr Lake (Wang et al., 2012); (k) the temperature differences between sea surface (based on alkenone) and thermocline (based on thermocline-dwelling planktonic foraminifera *Pulleniatina obliquiloculata*) recorded by core MD05-2904 (Steinke et al., 2011).

## Characteristics on the seasonal march of rainfall at Manila for the late 19<sup>th</sup> century - the early 20<sup>th</sup> century

\*赤坂 郁美<sup>1</sup>、財城 真寿美<sup>2</sup>、久保田 尚之<sup>3</sup>、松本 淳<sup>4</sup>

\*Ikumi Akasaka<sup>1</sup>, Masumi Zaiki<sup>2</sup>, Hisayuki Kubota<sup>3</sup>, Jun Matsumoto<sup>4</sup>

1. 専修大学、2. 成蹊大学、3. 東京大学、4. 首都大学東京

1. Senshu Univ., 2. Seikei Univ., 3. Tokyo Univ., 4. Tokyo Metropolitan Univ.

Rainfall is one of the most important climatic elements in Monsoon Asia, including the Philippines, because the seasonal change of rainfall is larger than that of temperature and is closely related to the water resources. Therefore, we aim to clarify longer-term variability in seasonal march of rainfall and its causes in the Philippines. To achieve our purpose, as the first step, we have collected and digitized the older meteorological observation records of the Philippines before the late 20<sup>th</sup> century under the data rescue projects in Japan. As the meteorological observations in the Philippines had been conducted by Spanish Jesuits for the late 19<sup>th</sup> century and by U.S. administration for the early 20<sup>th</sup> century, those data were found in the different places (e.g. UK, Spain and Japan). By connecting those data, we made the historical rainfall dataset in the Philippines for the period.

From the dataset, we used daily rainfall data at Manila where has the longest records of the observation in the Philippines. Based on daily rainfall at Manila from 1868 to 1940, we calculated pentad rainfall to study the seasonal change excluding daily rainfall variations. There are no data in 1875, 1877 and 1889. Manila has distinct dry season for February-April and wet season for May-October. Thus, to investigate the long-term changes in the seasonal march of rainfall at Manila, we determined the onset and withdrawal pentads of the rainy season: the onset (withdrawal) pentad corresponds to the first pentad when the pentad rainfall exceeds (falls below) 25mm since April. As results, the inter-annual variability in the onset of the rainy season since 1914 was small and the delayed withdrawal frequently appeared compared to the period before 1914. The durations of the rainy seasons for 1914-1940 were longer than those for 1950-2012. We will discuss these characteristics on the seasonal march of rainfall at Manila for the late 19<sup>th</sup> century-the early 20<sup>th</sup> century and its relation to the long-term variability in the Asian summer monsoon.

キーワード：マニラ、降水量、季節進行、長期変化、データレスキュー

Keywords: Manila, rainfall, seasonal march, long-term change, data rescue

## Projections of the duration of low-precipitation season in the Chao Phraya river basin based on the output from CMIP5 GCMs

\*渡部 哲史<sup>1</sup>、内海 信幸<sup>2</sup>、平林 由希子<sup>2</sup>、鼎 信次郎<sup>3</sup>

\*Satoshi Watanabe<sup>1</sup>, Nobuyuki Utsumi<sup>2</sup>, Yukiko Hirabayashi<sup>2</sup>, Shinjiro Kanae<sup>3</sup>

1. 東京大学大学院工学系研究科、2. 東京大学生産技術研究所、3. 東京工業大学環境・社会理工学院

1. School of Engineering, the University of Tokyo, 2. Institute of Industrial Science, the University of Tokyo, 3. School of Environment and Society, Tokyo Institute of Technology

The duration of low-precipitation season under climate change was projected in the Chao Phraya river basin based on the output from all 31 CMIP5 GCMs which is available for both historical (1951-1981) and RCP8.5 (2070-2100) emission scenario. We estimated the length of the continuous days in each year that total precipitation during preceding 30days below a threshold which defines low-precipitation season, 15, 30, 45, and 60 mm/30days in this study (the annual average is 82.4 mm/30days). The result indicates that the top 10 percentile of long duration becomes much longer under climate change while the average duration slightly decrease. This tendency is valid for each thresholds. In the case of 15 mm/30days threshold, the occurrence of 10<sup>th</sup> longest duration in historical period (1951-80 in this study), which corresponds almost 0.1% (ones in ten years), becomes 3.79 times as frequent under climate change from the estimation using 31 GCMs. The range of changing ratio estimated without highest and lowest 2 GCMs, which corresponds almost 90% confidence level, is 1.00 to 6.33. The fact suggests that the severe low-precipitation will happen more often under climate change.

The result of projection is significantly different between with and without applying bias correction method. For the average duration, an increase trend calculated without bias correction changes to a decrease trend after bias correction. It is well known that precipitation simulated by GCMs generally have considerable bias, thus it is common to correct bias before the application. This is true for the projections of the duration of low-precipitation season. To the best of our knowledge, there is no specific correction method for this purpose. Hence, we developed a method that correct the duration of low-precipitation season directly by changing threshold of precipitation for GCMs so that a duration of GCM low-precipitation season calculated by corrected threshold is agree to that of observation by original threshold in historical period. The developed method is different from common bias correction method in terms of the characteristic that not a precipitation itself but a threshold is corrected. This approach used in the correction of low-precipitation amount, which correct low-precipitation below a threshold as 0 considering the characteristic of GCM that there are significantly larger number of low-precipitation than observation.

It is important to understand the change of the duration of low-precipitation season because not only it has some impact on hydrology but also it affects the accuracy of bias correction for the amount of precipitation especially for pre- and post-monsoon season because many of bias correction methods adopt the approach that low-precipitation and others are separately corrected. Due to this reason, it is known that the error of bias correction generally large in these seasons. The results of this study can contribute to the improvement of bias correction as well as understanding the characteristics of the projections of precipitation among GCMs in monsoon regions.

キーワード：気候変動、降水、チャオプラヤ川流域

Keywords: Climate change, Precipitation, Chao Phraya river basin

## Uncertainty from climate forcing of glacier projection for High Mountain Asia

\*渡辺 恵<sup>1</sup>、渡部 哲史<sup>3</sup>、平林 由希子<sup>3</sup>、吉川 沙耶花<sup>2</sup>、鼎 信次郎<sup>2</sup>

\*Watanabe Megumi<sup>1</sup>, Satoshi Watanabe<sup>3</sup>, Yukiko Hirabayashi<sup>3</sup>, Sayaka Yoshikawa<sup>2</sup>, Shinjiro Kanae<sup>2</sup>

1. 東京工業大学情報理工学研究科、2. 東京工業大学環境・社会理工学院、3. 東京大学大学院工学系研究科

1. Graduate School of Information Science and Engineering, Tokyo Institute of Technology, 2. School of Environment and Society, Tokyo Institute of Technology, 3. School of Engineering, the University of Tokyo

Current model-based projections in glacier runoff are affected by considerable uncertainties. One of the largest uncertainties originates from climate forcing especially precipitation data. Underestimation of input precipitation data due to poor gauge network in mountainous regions is serious problem. Climate model driven information on climate change is also needed for future projection of glacier but often deemed unreliable. Those hamper effort to simulate the glacier runoff peak timing and magnitude. This research aims at an assessment of the major uncertainties from climate forcing in the modeling of future glacier runoff.

The glacier runoffs were calculated by a glacier model (HYOGA2) forced by two precipitation data and observed temperature data over historical period. Future glacier runoff was projected forced by eight climate models under the RCP4.5 and RCP8.5. The glacier model was calibrated using two precipitation data-sets. Bias correction of climate models were also done by comparing against two precipitation data-sets. The uncertainty of glacier runoff projection from input precipitation datasets and climate model spread will be discussed.

キーワード：氷河、降水、GCM

Keywords: Glacier, Precipitation, GCM

## Seasonal Responses of Pacific Japan Teleconnection on Indo-China Peninsula

\*Menaka Revel<sup>1</sup>, Utsumi Nobuyuki<sup>2</sup>, Sayaka Yoshikawa<sup>1</sup>, Shinjiro Kanae<sup>1</sup>

1. Department of Civil and Environmental Engineering, Tokyo Institute of Technology, 2. Institute of Industrial Science, The University of Tokyo

Summer monsoon precipitation provides a valuable support to the livelihood of the people live in Southeast Asia where the population density is very high. But monsoon precipitation shows high variation in seasonal and yearly time scales affecting daily life of the people in the regions such Indo-China Peninsula where most of the countries depend on agricultural economy. Tropical cyclones (TCs) and westward-propagation disturbances (WDs) are some of the major summer precipitation providing weather systems of this region. Pacific-Japan (PJ) teleconnection is a meridional teleconnection pattern which dominate the summer in Western North Pacific (WNP). TC locations were identified by TC best track dataset and WDs were detected subjectively using reanalysis dataset. PJ pattern timeseries is represented as 1<sup>st</sup> principle component seasonal average of relative vorticity at 850 hPa. High interannual and interseasonal variation of occurrence frequency was observed in Indo-China Peninsula which is not understood well enough. TC occurrence and genesis showed increasing trend in PJ pattern whereas it is negative for WDs. TCs and WDs have positive correlation with landfall in Indo-China Peninsula.

Keywords: Pacific Japan teleconnection, westward-propagating disturbance

# Influence of global warming on Eurasian snow cover teleconnection to the Indian monsoon rainfall using a large ensemble AGCM experiment

\*M.M. Gavin Dayanga Madakumbura<sup>1,2</sup>, Satoshi Watanabe<sup>2</sup>, Masahiro Tanoue<sup>2</sup>, Yukiko Imada<sup>3</sup>, Yukiko Hirabayashi<sup>2</sup>

1. Department of Civil Engineering, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, Japan. , 2. Institute of Engineering Innovation, The University of Tokyo, 2-11-16 Yayoi, Bunkyo-ku, Tokyo 113-8656, Japan. , 3. Meteorological Research Institute, Japan Meteorological Agency, 1-1 Nagamine, Tsukuba, Ibaraki 305-0052, Japan.

Relationship with Eurasian snow cover (ESC) and Indian summer monsoon rainfall (ISMR) is an extensively discussed aspect in terms of monsoon forecasting. Previous studies have found a strong negative correlation with the winter snow cover of western Eurasian region and ISMR. A weakening of this negative correlation has been observed recently but the reasons behind this are inconclusive, especially due to the limited observational record of snow amount. Influence of global warming on the change in snow-monsoon teleconnection has been widely discussed and suggested to have impacts on typical monsoon behavior such as the inverse relationship with ENSO. Therefore, this study was carried out to investigate the influence of anthropogenic global warming on the snow-monsoon relationship using a large ensemble experiment with and without human influence on the climate, using MRI-AGCM.

Carefully conducted correlation and composite analysis showed that the global warming has a possible weakening effect on the ESC-ISMR inverse relationship. This impact seems to be inflicted upon the ISMR with a modulation in the summer walker circulation anomaly over the South/South-East Asian region. Based on the correlation analysis, the impact of the global warming was shown to be less than the change observed from the observation-based analysis. Therefore, recent (after 1990) and past (before 1990) time slices were analyzed using correlation and composite based methods to observe any apparent deviations. Both ensemble simulations with and without human influence showed a similar decrease of the negative relationship with a westward shift of the rising anomalies associated with Indian ocean walker circulation during recent heavy snow years. This result was consistent with observations, suggesting a low-frequency variation of the circulation patterns associated with the ESC-ISMR relation due to the stochastic nature of the processes occurs from the natural variability, independent from global warming. Based on these results, we speculate that although an apparent weakening impact from global warming has occurred, more dominant low-frequency variability might be the reason for the significant reduction of ESC-ISMR correlation during recent decades.

Keywords: Indian monsoon, Eurasian snow cover, Teleconnections, Global warming