

東シナ海陸棚起源のマアジ (*Trachurus japonicus*) 0歳魚の太平洋側沿岸域への輸送・来遊実態の解明

Transport of age-0 jack mackerel (*Trachurus japonicus*) from the East China Sea to coastal areas along the Kuroshio.

*石川 和雄¹、伊藤 幸彦¹、渡邊 千夏子²、清水 学³、渡慶次 力⁴、西口 政治⁴、橋田 大輔⁵、大河 俊之⁶、原田 慈雄⁷、岡田 誠⁸、鈴木 勇己⁹、高村 正造¹⁰、福元 亨介¹¹

*Ishikawa Kazuo¹, Sachihiko Itoh¹, Chikako Watanabe², Manabu Shimizu³, Tsutomu Tokeshi⁴, Nishiguchi Masaharu⁴, Daisuke Hashida⁵, Toshiyuki Ohkawa⁶, Shigeo Harada⁷, Makoto Okada⁸, Yuki Suzuki⁹, Syozo Takamura¹⁰, Ryousuke Fukumoto¹¹

1. 東京大学大気海洋研究所、2. 国立研究開発法人 水産研究・教育機構 中央水産研究所、3. 国立研究開発法人 水産研究・教育機構、4. 宮崎県水産試験場、5. 愛媛県水産研究センター、6. 高知県水産試験場、7. 和歌山県水産試験所、8. 三重県水産試験場、9. 静岡県水産技術研究所 伊豆分場、10. 神奈川県水産技術センター 相模湾試験場、11. 鹿児島県水産技術開発センター

1. Atmosphere and Ocean Research Institute, The University of Tokyo, 2. National Research Institute of Fisheries Science, Japan Fisheries Research and Education Agency, 3. Japan Fisheries Research and Education Agency, 4. Miyazaki Prefectural Fisheries Research Institute, 5. Fisheries Research Center, Ehime Research Institute of Agriculture, Forestry and Fisheries, 6. Kochi Prefectural Fisheries Experimental Station, 7. Wakayama Prefectural Fisheries Experimental Station, 8. Mie Prefecture Fisheries Research Institute, 9. Izu Branch, Shizuoka Prefectural Research Institute of Fishery, 10. Sagami Bay Experimental Station, Kanagawa Prefectural Fisheries Research Institute, 11. Kagoshima Prefectural Fisheries Technology and Development Center

Jack mackerel (*Trachurus japonicus*) inhabiting coastal areas in temperate western North Pacific is an important fisheries resource in East Asian countries. Ichthyoplankton sampling surveys indicate that large spawning grounds are formed in the south of East China Sea from February to March, which is thought to be the main source of the resources in the Pacific coastal waters of Japan. However, transport and recruitment processes from the East China Sea to the coastal waters of Japan have yet to be quantified. In the present study, transport of age-0 jack mackerel by the Kuroshio and the recruitment to the southern coastal areas of Japan were investigated by analyzing fisheries landing records and temperature data from real time monitoring buoy systems. Mean seasonal fluctuation of the landing of age-0 jack mackerel compiled from 10-years records showed multiple peaks in the western part (Miyazaki and Kochi prefectures), first in mid spring and second in mid fall, and single peak in the eastern part (Mie, Shizuoka and Kanagawa prefectures). Time lags of the first peaks are not linearly related to the distance between the fishing grounds, and the eastward propagation of the landing is clearly observed only from Ehime to Kochi. The timing of the peaks in the eastern part in Mie (first peak), Shizuoka (single), Kanagawa (single) prefectures were detected in mid fall, later than the timing expected for those from the East China Sea. These results suggest that larvae and juveniles from the East China Sea are not predominant source and local population reproduced in each area is also important. In the presentation, temperature fluctuations around the landing pulses and their physical characteristics are also discussed.

キーワード：マアジ、黒潮、卵稚仔輸送、加入過程

Keywords: Jack mackerel, Kuroshio, larval transport, recruitment process

低次生態系-魚モデルを結合させた西部北太平洋領域海洋モデルの開発 Development of a physical-biogeochemical-fish coupled model for the western North Pacific

*Yokoi Takaaki¹、伊藤 進一¹、Dussin Raphael²、Curchitser Enrique²

*Takaaki Yokoi¹, Shin-ichi Ito¹, Raphael Dussin², Enrique Curchitser²

1. 東京大学大気海洋研究所、2. Rutgers University

1. Atmosphere and Ocean Research Institute University of Tokyo, 2. Rutgers University

Japanese food culture is highly depends on seafood and marine capture fishery is one of the important food supply for Japanese. However, the catch of many fish species in Japan showed multi-decadal fluctuations. Species alternation of sardine, anchovy, and chub mackerel is one of the distinctive phenomena. Until now, the mechanisms of fish stock fluctuations are still unclear, although many studies have been conducted. The authors are focusing on a 18.6-years tidal oscillation. Tides generate strong mixing interacting with bottom topography. Therefore, we focused on the Kuril Islands, Izu Ridge, and Nasei Islands as tidal mixing hotspots. As fish species live in tidal mixing hotspot areas in the western North Pacific, the authors are focusing on Jack mackerel, chub mackerel, and Pacific cod. To investigate the influence of the 18.6-years tidal oscillation on those species, we are planning to apply the Regional Ocean Modeling System (ROMS) coupled with North Pacific Ecosystem Model for Understanding Regional Oceanography For Including Saury and Herring (NEMURO.FISH) for chub mackerel in the western North Pacific (hereafter: this model is referred as the ROMS-NEMURO.FISH). The ROMS-NEMURO.FISH was driven by the atmospheric forcing of Common Ocean Reference Experiment version 2 (CORE2). Also, the Simple Ocean Data Assimilation (SODA) was used as the initial and the boundary conditions for the ROMS-NEMURO.FISH. In this presentation, as a first step, we briefly report the climatology and interannual fluctuation of physical fields and will discuss the model reproducibility to evaluate the model physical field from the point that the model can be applied for the fish stock fluctuations.

キーワード：海洋物理-生物地球化学-魚類結合モデル、ROMS、NEMURO.FISH、マサバ

Keywords: physical-biogeochemical-fish coupled model, ROMS, NEMURO.FISH, chub mackerel

A challenge to investigate environmental factors which determine spawning migration variability of Pacific saury

*伊藤 進一¹、笥 茂穂²、奥西 武²、巢山 哲²、安倍 大介²、亀田 卓彦²

*Shin-ichi Ito¹, Shigeho Kakehi², Takeshi Okunishi², Satoshi Suyama², Daisuke Ambe², Takahiko Kameda²

1. 東京大学大気海洋研究所、2. 水産研究・教育機構

1. Atmosphere and Ocean Research Institute, The University of Tokyo, 2. Fisheries Research and Education Agency

Spawning migration is one of the most difficult behavior to represent by a numerical model. For example, Pacific saury make a wide spawning migration from the subarctic to subtropical regions, while the detailed mechanism of the spawning migration has not been elucidated. Based on the synoptic survey of Pacific saury as realistic initial condition, an Euler-type model of Pacific saury was applied to investigate environmental factors which determine spawning migration variability. As realistic boundary conditions, satellite derived environmental conditions were used as forcing; sea surface temperature (SST), prey density estimated from surface chl-a concentration and surface current velocity fields. Growth of Pacific saury was calculated based on a fish bioenergetics model (NEMURO.FISH). A fitness algorithm was applied for feeding migration in which the fish are assumed to be moving towards a place with optimal growth condition. A larvae fitness algorithm was applied for spawning migration in which the spawning fish moves to a place of the optimal growth of larvae. For spawning migration, westward migration was added to reproduce realistic spawning grounds around Japan Islands. Strength of the westward migration during spawning migration was adjusted to realize observed variability of saury migration to fishing grounds. The adjusted westward migration variability was compared to the environmental factors.

キーワード：海洋生態系モデル、魚類成長-回遊モデル、サンマ、気候変動

Keywords: marine ecosystem model, fish growth-migration model, Pacific saury, climate variability

温暖化に対する海洋炭素循環の応答の再検討

Revisiting response of oceanic carbon cycle to global warming

*山本 彬友¹、阿部 彩子¹、山中 康裕²

*Akitomo Yamamoto¹, Ayako Abe-Ouchi¹, Yasuhiro Yamanaka²

1. 東京大学大気海洋研究所、2. 北海道大学大学院地球環境科学研究院統合環境科学部門

1. Atmosphere and Ocean Research Institute, The University of Tokyo, 2. Faculty of Environmental Earth Science, Hokkaido University

Climate warming tends to reduce oceanic uptake of CO₂ from the atmosphere thereby accelerating the rate of CO₂ accumulation in the atmosphere and global warming. Most previous studies have shown that the physical effects, decrease in CO₂ solubility and increased stratification, are major contributors to this reduction in oceanic CO₂ uptake [Sarmiento et al., 1998; Plattner et al., 2001]. In those previous studies, changes in the biological pump associated with ocean circulation change were regarded as a second-order process even though biological effects on the natural carbon cycle can be very important. However, the contributions of both physical and biological effects to the reduction in oceanic CO₂ uptake are not evaluated directly in the recent generation of coupled atmosphere–ocean general circulation models (AOGCMs) and ocean biogeochemical models. To address this we reevaluate the individual mechanisms contributing to the reduction in oceanic CO₂ uptake using a series of multi-centennial global warming simulation conducted with AOGCM and an offline ocean biogeochemical model. The uptake reduction of 13% due to global warming at 140 years is consistent with the same simulation using models in the CMIP5 [Arora et al., 2013]. Sensitivity studies show that changes in the biological pump and gas solubility are the dominant processes for this reduction in oceanic carbon uptake, which is opposite to most of the previous studies: changes in ocean circulation and solubility are the dominant processes. Decrease in new production caused by lower nutrient supply and enhanced remineralization from seawater warming increase dissolved inorganic carbon at the surface, thereby substantially preventing oceanic CO₂ uptake. The weaker Atlantic meridional overturning circulation reduces oceanic CO₂ uptake, while weaker equatorial upwelling and increased mixing due to enhanced westerly winds in the Southern Hemisphere enhance CO₂ uptake. As these effects cancel each other out, the effect of circulation change becomes a second-order process. Our results demonstrate that the biological pump plays a significant role in not only natural carbon cycle but also anthropogenic carbon cycle.

キーワード：地球温暖化、炭素循環、海洋物質循環モデル

Keywords: Global warming, Carbon cycle, Ocean biogeochemical model

CMIP5モデルを用いた温暖化が海洋基礎生産へ及ぼす影響の解析

Analysis of future changes in ocean primary production using CMIP5 models

*中村 有希¹、岡 顕¹

*Nakamura Yuki¹, Akira Oka¹

1. 東京大学

1. University of Tokyo

温暖化により海洋では成層化、酸性化、脱酸素化などの変化が生じ、海洋生態系へも影響が及ぶことが懸念されている。モデルを用いた複数の先行研究によって、将来的に植物プランクトンによる全球の海洋基礎生産（Net Primary Production ; NPP）と海洋輸出生産（Export Production ; EP）は減少することが予測されており、その主な原因は成層化による栄養塩供給の減少であると考えられている(Fu et al.,2015ほか)。しかし、CMIP5モデル間の結果を比較した研究によると、EPについては全てのモデルが減少を示す一方で、NPPについてはほとんど変化しないモデルもみられた(Bopp et al.,2013)。本研究の目的は、10個のCMIP5モデルを用いて、温暖化進行時にNPPの変化がモデルにより大きく異なる要因を明らかにすることである。

全球的なNPPとEPの変化には熱帯の寄与が大きいことがわかったため、熱帯に着目して研究を行った。NPP、EPに関してCMIP5モデル間の相互比較を行うことにより、NPPの変化がモデルによって異なる要因は、1. 成層化、2. 再無機化の温度依存性、の程度の違いによって説明できることがわかった。温暖化した際にNPPがほとんど変化しない例外的なモデル(GFDL-ESM2G)に関しては、温度に依存する再無機化のパラメータが過大評価されている可能性が考えられる。つまり温暖化で再無機化が活発化したことによって、EPは減少しているにも関わらずNPPは変化しないと考えられる。Nino.3 SST Indexとの回帰から、ENSOに伴うNPP変化を調べたところ、衛星データによる評価および他のモデルではエルニーニョ時にNPPが低下するのに対し、温暖化時にNPPがほとんど変化しなかったモデルについては、上昇することがわかった。ENSOなどの自然変動に伴うNPPの応答を各モデルで評価することは、より正確な将来予測へとつながることが示唆される。

今後は温暖化により海洋基礎生産が炭素循環へ与える影響の変化を明らかにすることを目的とし、海洋生態系モデルを用いて感度実験を行う予定である。

キーワード：海洋基礎生産、海洋輸出生産、CMIP5、温暖化

Keywords: Primary Production, Export Production, CMIP5, global warming

Preliminary experiment of ocean acidification in the western North Pacific

*橋岡 豪人¹、杉江 恒二¹、重光 雅仁¹

*Taketo Hashioka¹, Koji Sugie¹, Masahito Shigemitsu¹

1. 国立研究開発法人 海洋研究開発機構

1. Japan Agency for Marine-Science and Technology

Responses of marine phytoplankton to ocean acidification is reported from an on-board manipulation experiment in the western North Pacific (March 9, 2015; 41.9N, 146E). In this experiment, it is suggested that an increase in nano-/pico-phytoplankton and a decrease in diatoms under high pCO₂ conditions (1420 and 2850 μ atm; 1420 μ atm corresponds to Y2150 of AR5 RCP8.5 scenario), respectively. In this study, we parameterized the responses of phytoplankton to acidification, and conducted a simplified ocean acidification experiment using a 1D ecosystem model, MEM (Shigemitsu et al., 2012) at the Station A04 (42.15N, 145.07E). The results showed a potential of delay in spring bloom with decrease in the magnitude, and an importance of acidification on not only calcifiers but also on general phytoplankton.

キーワード：海洋酸性化、生態系モデル、西部北太平洋

Keywords: Ocean Acidification, Ecosystem Model, western north Pacific

Modelling the impact of riverine and atmospheric nitrogen inputs on the marine biogeochemistry

*相田 真希¹、渡辺 路生²、羽島 知洋³

*Maki Noguchi Aita¹, Michio Watanabe², Tomohiro Hajima³

1. 国立研究開発法人海洋研究開発機構 地球環境観測研究開発センター、2. 国立研究開発法人海洋研究開発機構 気候変動リスク情報創生プロジェクトチーム、3. 国立研究開発法人海洋研究開発機構 統合的気候変動予測研究分野

1. Research and Development Center for Global Change, Japan Agency for Marine-Earth Science and Technology, 2. Project Team for Risk Information on Climate Change, Japan Agency for Marine-Earth Science and Technology, 3. Department of Integrated Climate Change Projection Research, Japan Agency for Marine-Earth Science and Technology

Nitrogen in the ocean plays a key role in primary production, and biological activities, such as assimilation of phytoplankton and decomposition of organic matter are well studied. However, nitrogen cycle of the basin scale is still unclear because of the quite long time scale of the processes such as denitrification, sediment processes, and advection caused by meridional overturning circulation. Furthermore, the impact of the atmospheric and riverine nitrogen input has not been clarified. Human activities will change the nitrogen cycle in the ocean shifting riverine and atmospheric nitrogen inputs, so that it is required to study the nitrogen balance in the whole ocean. To evaluate the nitrogen cycle in the ocean, we created a marine biogeochemical model including processes such as nitrogen fixation, denitrification, atmospheric and riverine nitrogen input, then embedded it into an Earth System Model: MIROC-ESM 5.9 developed by Program for Risk Information on Climate Change, JAMSTEC. We spun up MIROC-ESM for 800 years and analyzed the results of the last 10 simulated years. The model results showed that riverine nitrogen and atmospheric nitrogen inputs changes the production of phytoplankton about 20% and 10%, respectively, especially in the subtropical region of the eastern North Pacific and equatorial region of the western Atlantic Ocean. Riverine nitrogen input changed nitrate concentration up to $\pm 2.5 \text{ } \mu\text{mol L}^{-1}$ in the high latitudes such as the subarctic and polar regions (excluding the coastal regions). Our results showed the importance of riverine and atmospheric nitrogen inputs to the basin-scale nitrogen cycle.

キーワード：窒素循環、河川窒素流入、大気窒素化合物沈着、海洋物質循環、地球システムモデル

Keywords: Nitrogen cycle, riverine nitrogen input, atmospheric nitrogen deposition, marine biogeochemistry, Earth System Model

Biophysical Controls on Vertical Fluxes of Dissolved and Particulate Carbon, Nitrogen and Phosphorus in the Northern South China Sea

*Jia-Jang Hung¹, Z.-Y. Lin¹, S.-H. Peng¹

1. Department of Oceanography, National Sun Yat Sen University, Kaohsiung, Taiwan

Vertical fluxes of particulate organic matter and diffusion flux of dissolved organic matters (DOM) are the major part of biological pump in the ocean. In this study, we employed shipboard observations and sediment-trap deployments to collect water samples and sinking materials for hydrographic and chemical analyses in the northern South China Sea (NSCS). In addition to seasonal variability, effects of eddy and internal waves on vertical fluxes of carbon (C), nitrogen (N) and phosphorus (P) are also explored. Vertical fluxes of C, N and P in the NSCS were respectively estimated to be $66.8 \pm 1.29 \text{ mg C m}^{-2} \text{ d}^{-1}$, $12.8 \pm 0.38 \text{ mg N m}^{-2} \text{ d}^{-1}$ and $0.99 \pm 0.07 \text{ mg P m}^{-2} \text{ d}^{-1}$ in the spring season, about $64.3 \pm 1.47 \text{ mg C m}^{-2} \text{ d}^{-1}$, $12.1 \pm 0.47 \text{ mg N m}^{-2} \text{ d}^{-1}$ and $0.93 \pm 0.04 \text{ mg P m}^{-2} \text{ d}^{-1}$ in the summer season, and about $155 \pm 15.9 \text{ mg C m}^{-2} \text{ d}^{-1}$, $21.2 \pm 1.68 \text{ mg N m}^{-2} \text{ d}^{-1}$ and $1.79 \pm 0.19 \text{ mg P m}^{-2} \text{ d}^{-1}$ in an internal-waves induced event. Meanwhile, the organic carbon flux derived from the modified Vertical Generalized Production Model (VGPM) was $183 \pm 17 \text{ mg C m}^{-2} \text{ d}^{-1}$ in a winter anticyclonic eddy in NSCS. Positive correlations were significant between the chlorophyll inventory and DIN inventory, and also significant between vertical fluxes of CNP and DIN inventory in the euphotic zone. Thus, vertical fluxes of CNP were apparently driven by primary production that was determined by the availability of DIN. The diffusion flux of DOM increased generally with depth and was enhanced obviously in the internal-waves condition. The proportions of CNP vertical fluxes to the biological pump in NSCS are 66.5%, 74.3% and 71.3%, respectively, in regular spring, about $63.8 \pm 9.18\%$, $76.6 \pm 0.58\%$ and $75.1 \pm 2.02\%$, respectively, in regular summer, and about 62.1%, 73.5% and 64.7%, respectively, in an internal-waves induced event. The vertical flux accounts for the highest proportion of the biological pump. However, the proportion of DOM flux is insignificant in the biological pump.

Keywords: Vertical fluxes, Biological pump, Carbon, Nitrogen and Phosphorus, Northern South China Sea

Impact of mesoscale eddies on spring bloom initiation in the Japan Sea

*Eligio de Raus Maure¹, Joji Ishizaka², Chiho Sukigara¹, Yoshihisa Mino², Hidenori Aiki², Takeshi Matsuno³, Hiroyuki Tomita²

1. Graduate School of Environmental Studies, Nagoya University, 2. Institute for Space-Earth Environmental Research, Nagoya University, 3. Research Institute for Applied Mechanics, Kyushu University

Mesoscale eddies play an important role in the ocean primary production and biogeochemical processes and induce large spatial and temporal variability of near surface chlorophyll-a concentration (CHL). In particular, they are known to impact on the temporal variation of mixed layer depth (MLD), which is important for biological processes such as phytoplankton spring bloom phenomenon, a recurring phenomenon observed in temperate seas that results in large increase in phytoplankton abundance in springtime. In this study we investigated the influence of mesoscale eddies on timing of spring bloom initiation in the Yamato Basin region (133-139° E and 35-39.5° N), Japan Sea, in a period spanning 2002-2011. We identified mesoscale eddies based on geometric characteristic of satellite estimated geostrophic velocity around eddies (Nencioli et al., 2010). Eddy region was defined based sea level anomaly (SLA) data as the outmost closed contour of SLA enclosing identified centre (Chelton et al., 2011). CHL from winter (January) to early summer (June) was used to capture the spring bloom event. We also used in-situ profiles of temperature-salinity data within eddy region to estimate the MLD on individual profiles as the density change of 0.03 kg m^{-3} from surface reference level (10 m). Results showed that bloom was initiated early in cyclonic eddies (CEs) with shallow MLD (mostly < 100 m) compared to anticyclonic eddies (AEs) in which it was initiated despite the deeper MLD (> 100 m). From the examination of net heat flux (Q_0) within eddies we found that the onset of spring bloom in CEs occurred while large heat loss to the atmosphere was being observed, whereas in AEs it was observed close to the commencement of positive Q_0 . This suggested that in AEs relaxation of turbulent mixing was important for bloom to start, whereas in CEs, because of shallower MLD, improved light condition for phytoplankton growth within the turbulent layer was reached earlier, thus triggering the bloom.

Keywords: Mesoscale eddies, mixed layer depth, phytoplankton, spring bloom

Global size distribution of phytoplankton communities from space

*平田 貴文¹、増田 良帆¹、鈴木 光次¹、山中 康裕¹

*Takafumi Hirata¹, Yoshio Masuda¹, Koji Suzuki¹, Yasuhiro Yamanaka¹

1. 北海道大学地球環境科学研究院

1. Faculty of Environmental Earth Science, Hokkaido University

We developed a remote sensing methodology to estimate size distribution of various pigment-based phytoplankton groups such as diatoms, peridinin-containing dinoflagellates, haptophytes, cyanobacteria etc. Our estimation was compared with a local in situ observation to show an agreement between them. According to our state-of-the-art remote sensing methodology, global size structure of the entire phytoplankton community could be divided into three classes to the first approximation, agreeing well with a conventional classification based on historical in situ observations. However, in contrast to historical size classifications (Sieburth et al, 1978), i.e. pico-phytoplankton ($< 2 \mu\text{m}$), nano-phytoplankton $2\text{-}20 \mu\text{m}$, micro-phytoplankton ($> 20 \mu\text{m}$), we propose new size boundaries for these classes based on global satellite observation: pico-phytoplankton ($< 1 \mu\text{m}$), nano-phytoplankton $1\text{-}10 \mu\text{m}$, micro-phytoplankton ($> 10 \mu\text{m}$). Size-diversity index of a given phytoplankton group, defined by a difference between logarithmic maximum and minimum sizes of the group, was largest for haptophytes than diatoms. The maximum size-diversity of a given phytoplankton group was not necessarily correlated to its dominance in chlorophyll abundance either. Our results are expected to cast light upon global marine biodiversity and marine ecosystem analysis.

キーワード：植物プランクトン、生物多様性、サイズ、海色、生態系、海洋

Keywords: phytoplankton, biodiversity, size, ocean color, ecosystem, ocean

Converting electron transfer rates of phytoplankton into daily carbon fixation rates in coastal waters of Ariake Bay and the East China Sea

*Yuanli Zhu¹, Joji Ishizaka², Sarat Tripathy³, Takeshi Matsuno⁴, David Suggett⁵

1. Graduate School of Environmental Studies, Nagoya University, Japan, 2. Institute for Space-Earth Environmental Research, Nagoya University, Japan, 3. ESSO-National Centre for Antarctic and Ocean Research, Ministry of Earth Sciences, India, 4. Research Institute for Applied Mechanics, Kyushu University, Japan, 5. Climate Change Cluster, University of Technology Sydney, Australia

In the last 20 years, introduction of the active fluorescence technique, fast repetition rate fluorometer (FRRf), has opened up capability to measure instantaneous electron flow rate through phytoplankton photosystem II (electron transfer rate, ETR). However, in most of the early studies, the conversion factor (K_c) from ETR to primary production was assumed to be constant. Recent studies have shown that K_c can vary over a large range in nature and that understanding of the variability of K_c in relation to other physical and biogeochemical parameters is the crucial for accurately estimating primary production with FRRf. In this study, K_c was determined from parallel measurements of ETR and daily net primary production (NPP) and modeled as a function of key environments and phytoplankton community structure for the first time in Ariake bay and the East China Sea. In Ariake Bay study, we firstly confirmed that K_c varied considerably in nature and then demonstrated the strong correlation ($R^2 = 0.94$) between daily photosynthetically active radiation (PAR) and K_c ; the novel and simple PAR-dependent relationship used for deriving K_c opens the possibility for directly FRRf based NPP estimating. In the East China Sea study, we confirmed that PAR was still the main controlling factor of K_c ($R^2 = 0.72$); moreover, this factor appeared secondarily influenced by dominant phytoplankton taxa present. Results showed that the correlation between K_c and PAR was improved ($R^2 = 0.78-0.86$) by considering two clusters of taxonomical groups, large phytoplankton ($>20 \mu\text{m}$) and small phytoplankton ($<20 \mu\text{m}$). Overall, we have developed a novel algorithm for estimating K_c from PAR and phytoplankton community composition and this algorithm can be applied to future FRRf-based high resolution studies of primary production.

Keywords: ETR, Primary productivity, Quantum requirement for carbon fixation, Phytoplankton composition, Fast repetitive rate fluorometer