

The Impact of Ocean Surface Currents on Global Eddy Kinetic Energy via the Wind Stress Formulation

*Zhitao Yu¹, Edward Joe Metzger¹

1. US NRL

A more complete wind stress formulation takes into account the ocean surface currents, while the conventional wind stress popularly used in ocean circulation models is only a function of 10-m winds. A pair of 12.5-year (July 2002 –December 2014) HYbrid Coordinate Ocean Model (HYCOM) simulations that only differ in the wind stress formulation are used to study the impact of ocean surface currents on global Eddy Kinetic Energy (EKE). The model results (2004-2014) show that including ocean surface currents in surface wind stress formulation reduces global EKE by more than 40%. To understand the mechanisms behind the large difference, we calculate the global EKE budget using the standard Reynolds averaging procedure. The direct impact of surface wind stress on EKE is through surface wind work. The indirect impact is through changes in the mean circulation that affect the shear production, buoyancy work, and bottom friction. Model Results indicate that the reduction of global EKE is primarily due to the buoyancy work and shear production. Even though the surface eddy wind work is one order larger than buoyancy work, and two orders larger than shear production, it is mainly balanced by bottom friction and thus not the main contribution for the EKE differences.

Keywords: HYCOM, eddy kinetic energy, wind work, baroclinic instability, wind stress

Long-term change and variation of salinity in the western North Pacific subtropical gyre revealed by 50-year long observations along 137°E

*岡 英太郎¹、桂 将太¹、井上 博敬²、小嶋 惇^{2,3}、北本 萌子^{2,3}、中野 俊也^{2,3}、須賀 利雄^{4,5}

*Eitarou Oka¹, Shota Katsura¹, Hiroyuki Inoue², Atsushi Kojima^{2,3}, Moeko Kitamoto^{2,3}, Toshiya Nakano^{2,3}, Toshio Suga^{4,5}

1. 東京大学大気海洋研究所、2. 気象庁地球環境・海洋部、3. 気象研究所海洋・地球化学研究部、4. 東北大学理学研究科、5. 海洋研究開発機構地球環境観測研究開発センター

1. Atmosphere and Ocean Research Institute, The University of Tokyo, 2. Global Environment and Marine Department, Japan Meteorological Agency, 3. Oceanography and Geochemistry Research Department, Meteorological Research Institute, 4. Department of Geophysics, Graduate School of Science, Tohoku University, 5. Research and Development Center for Global Change, Japan Agency for Marine-Earth Science and Technology

The 137°E repeat hydrographic section for 50 winters during 1967–2016 has been analyzed to examine interannual to interdecadal variations and long-term changes of salinity and temperature in the surface and intermediate layers of the western North Pacific, with a particular focus on freshening in the subtropical gyre. Rapid freshening on both isobars and isopycnals began in mid-1990' s and persisted for the last 20 years in the upper main thermocline/halocline in the western subtropical gyre. In addition, significant decadal variability of salinity existed in the Subtropical Mode Water (STMW), as previously reported for the shallower layers. An analysis of the 144°E repeat hydrographic section during 1984–2013 supplemented by Argo profiling float data in 2014 and 2015 revealed that the freshening trend and decadal variability observed at 137°E originated in the winter mixed layer in the Kuroshio Extension (KE) region and was transmitted southwestward to 137°E 1–2 years later in association with the subduction and advection of STMW. The mechanism of these changes and variations in the source region was further investigated. In addition to the surface freshwater flux in the KE region pointed out by previous studies, the decadal KE variability in association with the Pacific Decadal Oscillation likely contributes to the decadal salinity variability through water exchange between the subtropics and the subarctic across the KE. Interdecadal change in both the surface freshwater flux and the KE state, however, failed to explain the rapid freshening for the last 20 years.

キーワード：西部北太平洋亜熱帯循環、主水温・塩分躍層、長期変化、十年規模変動、観測定線

Keywords: Western North Pacific subtropical gyre, Main thermocline/halocline, Long-term change, Decadal variability, Repeat hydrographic section

Interannual and inter-decadal variability of the North Equatorial Countercurrent in the Western Pacific


*Xiao Chen¹, Bo Qiu

1. Hohai University

Abstract

Interannual and longer timescale variations of the North Equatorial Countercurrent (NECC) in the western Pacific are investigated using the multi-decade (1960-2014) hindcast by the Ocean general circulation model for the Earth Simulator (OFES). The OFES-simulated sea level and upper ocean circulation changes show favorable comparisons with available tide gauge data and repeat hydrographic surveys along the 137°E meridian. An empirical orthogonal function (EOF) analysis reveals that the low-frequency NECC variability is dominated by two distinct modes. The first mode fluctuates interannually and shows strengthening and southward migration of the NECC concurrent with the development of El Niño events. Unlike the extra-tropical western Pacific Ocean circulation variability controlled by wind forcing west of the dateline, the interannual NECC variations are forced by equatorial wind forcing cumulative across the entire Pacific basin. The second mode of the NECC variability has a inter-decadal timescale and is characterized by NECC's progressive weakening in strength, migrating poleward, and broadening in width over the past 50 years. These long-term changes in NECC are caused by the corresponding changes in the trade wind system that weakened and expanded poleward in the past half a century across the Pacific basin.

Keywords: Interannual and Inter-decadal Variability, Western Pacific, Northern Equatorial Countercurrent



Seasonal transport variation over the entire Kuroshio region

*張 振龍¹、中村 啓彦¹

*ZHENLONG ZHANG¹, Hirohiko Nakamura¹

1. 鹿児島大学

1. Kagoshima University

As the observations of the Kuroshio were conducted regionally by different groups, studies for the seasonal variation and its mechanism were also proceeded separately. By using the data collected by satellites and assimilation model's outputs, we focus on the seasonal transport variation over the entire Kuroshio region. Our main purpose is to show the temporal and spatial feature of the seasonal transport variation and give an insight for its driving mechanism.

The results from Aviso data reveal that the surface transport variation reaches maximum in July for most of the Kuroshio region, but its minimum appears in different months. Near Luzon Island and in the East China Sea, the minimum values appear in October or November. For the area of Taiwan, most of the minimum values are found in January or February. Time when the minimum values occur also shows propagation signals: it shifts from September to November in Luzon Strait, while moves gradually from November to June south of Japan. Another feature is that the amplitudes of the surface transport variation are large near Luzon Island, Taiwan, and south of Japan, where there is a land boundary to the west of the Kuroshio. The results from EOF analysis, which is conducted by using an assimilation dataset estimated from the Multivariate Ocean Variation Estimation system, also indicate that the transport variation tends to dominate in the areas where the western land boundary exists.

Czeschel et al. (2012) suggested that fast barotropic waves propagate the signals due to wind-induced coastal upwelling (downwelling) southward along the simple slope off the North American coast and contribute largely to the annual cycle of the Florida current transport. While for the North Pacific western boundary, the barotropic waves are considered to propagate along the complicated coastline with large topographic modulation. This might be a possible cause that leads to the regional features of seasonal transport variation revealed in this study.

キーワード：黒潮、流量の季節変動

Keywords: Kuroshio, seasonal transport variation

Mooring Measurement of the Abyssal Circulations in the Western Pacific Ocean

*Jianing Wang¹, Fan Wang¹

1. Institute of Oceanology, Chinese Academy of Sciences

A scientific observing network in the western tropical Pacific has initially been established by the Institute of Oceanology, Chinese Academy of Sciences (IOCAS). Using fifteen moorings that gives unprecedented measurements in the intermediate and abyssal layers, we present multi-timescale variations of the deep ocean circulations prior to and during 2015 El Niño event. The deep ocean velocities increase equatorward with high standard deviation and nearly zero mean. The deep ocean currents mainly flow in meridional direction in the central Philippine Basin, and are dominated by a series of alternating westward and eastward zonal jets in the Caroline Basin. The currents in the deep channel connecting the East and West Mariana Basins mainly flow southeastward. Seasonal variation is present in the deep jets in the Caroline Basin, associating with vertical propagating annual Rossby wave. The high-frequency flow bands are dominated by diurnal, and semi-diurnal tidal currents, and near-inertial currents. The rough topography has a strong influence on the abyssal circulations, including the intensifications in velocity and internal tidal energy, and the formation of upwelling flow.

Keywords: Abyssal Circulation, Western Pacific, Mooring Measurement

西部熱帯太平洋における中層海洋変動

Ocean variability at intermediate depth in the western equatorial Pacific

*柏野 祐二¹、長谷川 拓也¹、Syamusdin Fadli²

*Yuji Kashino¹, Takuya Hasegawa¹, Fadli Syamusdin²

1. 海洋研究開発機構、2. インドネシア技術評価応用庁

1. JAMSTEC Japan Agency for Marine-Earth Science and Technology, 2. BPPT

東経156度以西の西部熱帯太平洋における中層海洋（水深300m～750m）の変動をトライトンブイや気象庁のCTD観測データなどを用いて調べた。変動は西岸近くで大きく、特に季節変動が顕著であった。季節内変動も無視できず、特にニューギニア沿岸潜流が赤道潜流に反転する北緯2度東経138度付近で大きい。そこでは、その反転が不安定でニューギニア渦という時計回りの渦が形成されており、その渦が周りの渦と結合・分離を繰り返している。経年変動もENSOスケールのみならず、準10年変動スケールで750m深においても見られるが、海域全体で同じ符号の相関が中層海洋の温度変動とこれらの経年変動の間に見られるのではないことから、単純に暖水プールの東西移動によってもたらされているのではなく、赤道波に伴うサーモクラインの形状の変化によってもたらされていると考えられる。また、10年以上の減少トレンドも塩分変化に見られ、これは北太平洋中層水や南極中層水の経年変化に伴っていると考えられる。

キーワード：中層海洋変動、西部熱帯太平洋、トライトンブイ、中層水

Keywords: Intermediate ocean variability, Western equatorial Pacific, TRITON buoy, Intermediate waters

Seasonal variability of salinity in the Indian Ocean in response to riverine freshwater forcing using ROMS

*K. K. Sandeep¹, Vimlesh Pant¹

1. Indian Institute of Technology Delhi

Indian Ocean exhibits large spatial and temporal variability in the sea surface salinity distribution. The unique geographical features causes the salinity distribution over the Arabian sea and Bay of Bengal quite different. The Arabian sea being connected to highly saline waters from Red Sea and Persian Gulf remains salty when compared to the Bay of Bengal. Major rivers such as Ganges, Brahmaputra, Mahanadi, Godavari, Krishna, and Irrawaddy debouch an excess amount of freshwater into the Bay. In addition to this, the annual precipitation over Bay of Bengal always remains higher than evaporation when compared to Arabian Sea. Due to these factors the Bay remains fresher than Arabian sea throughout the year and also exhibits large seasonal variability in the circulation patterns. The seasonal variability of sea surface salinity in the Indian ocean is studied by using a ocean general circulation model 'Regional Ocean Modeling System' (ROMS). The model domain extends from 30° S-30°N, 30°E-120°E with quarter degree resolution in the horizontal and 40 sigma levels. The model is initialized with annual mean climatology of tracer fields from World Ocean Atlas 2013 (WOA13) and forced with daily climatological winds from Quikscat and ASCAT and other atmospheric forcing fields from TropFlux. It is found that there is significant contribution to freshening from riverine freshwater discharge in coastal regions near river mouths. Over the Indian Ocean, northern boundaries of Bay of Bengal found to have largest impact of river discharges on SSS simulations during monsoon (June-September) season when the peak discharge from major rivers Ganges, Brahmaputra, and Irrawaddy accumulate freshwater plume along coastal regions in Bay of Bengal. The impact of continental freshwater discharge on the seasonal salinity pattern are investigated. Satellite and in-situ observations are also used to understand the response of riverine freshwater forcing in controlling the spatial and seasonal variations of salinity in various sectors of the Indian Ocean.

Keywords: Indian Ocean, Sea Surface Salinity, Seasonal Variability, River Discharge

Numerical study of Internal tide energetics in the Andaman Sea

*Sachiko Mohanty¹, A D Rao¹

1. Indian Institute of Technology Delhi

The interactions of barotropic tides with irregular bottom topography generate internal waves with high amplitude known as large-amplitude internal waves (LAIW). The Andaman Sea is one of the potential region where such waves occur. These waves are an important phenomena in the ocean due to their influence on the density structure and energy transfer into the region. These waves are also important in submarine acoustics, underwater navigation, offshore structures, ocean mixing, biogeochemical processes, etc. over the shelf-slope region.

A three-dimensional MIT general circulation ocean model (MITgcm) is configured over the Andaman sea to investigate the generation and propagation of M_2 internal tides. Initially, the model simulations are validated by using *in-situ* observations of temperature, conductivity and currents from a buoy located at 10.5N, 94E. The spectral energy estimate of density shows that the peak estimate is associated with the semi-diurnal frequency at all the depths in both observations and model simulations. The vertical structure of baroclinic velocity is compared with observations and the analysis suggests that a multi-mode features of baroclinic tides are present at the buoy location. To understand the generation and propagation of internal tides over this region, energy flux and barotropic-to-baroclinic M_2 tidal energy conversion rates are also estimated. The model simulation suggests that the internal tides are mainly generated at the north of Sumatra coast and the regions around the Nicobar islands. The internal tides propagate away from the respective generation sites. The steepness of topography suggests that the internal tide mainly generate at the supercritical slope region and the energy flux reflected back to the deep water from the supercritical slope. The M_2 dissipation rate is also found to be maximum at the generation sites.

Keywords: Internal tide, Andaman Sea, Energetics

Simulation of storm surges by improving the cyclonic wind formulation

*Smita Pandey¹, A D Rao¹

1. Indian Institute of Technology Delhi

Coastal regions of India are highly prone to sea level rise during tropical cyclones. Storm surges due to cyclones can produce high sea levels, especially when their occurrence coincide with the high tides. Storm surges, tides and wind waves are the main factors responsible for the variation in the total water elevation (TWE) near the coast. The accuracy in simulation of storm surges depends not only on the cyclonic track and its intensity, but also on the spatial distribution of winds which include its speed and direction. In the present study, the cyclonic winds are validated using buoy winds for the recent cyclones formed in the Bay of Bengal since 2010 using Jelesnianski wind scheme. It is found that the cyclonic winds computed from the scheme show an underestimate in the magnitude and also a mismatch in its direction. Hence, the wind scheme is suitably modified based on the buoy observations available at different locations using a power law which reduces the exponential decay of winds by about 30%. Moreover, the cyclonic wind direction is also corrected by suitably modifying its inflow angle. The significance of modified exponential factor and inflow angle in the computation cyclonic winds is highlighted using statistical analysis.

In the present study, simulation of storm surges for the recent cyclonic cases along the east coast of India are carried out using both stand-alone hydrodynamical depth integrated ADCIRC model and coupled ADCIRC+SWAN model. The coupled ADCIRC+SWAN model is used to incorporate the contribution of wind waves in the simulation of TWE near the coast. The cyclonic wind distribution is computed using both modified and unmodified Jelesnianski wind scheme. The experiments are performed to validate the TWE generated from the cyclones through computation of surge residuals with the available tide gauge data. On comparison of observed surge residuals with the simulations using modified winds from the uncoupled and coupled models, it is found that the simulated surge residuals are better compared, especially with the inclusion of wave effect through the coupled model.

Keywords: total water elevations, storm surge, cyclonic winds, ADCIRC MODEL

九州西方海域で発生する気象津波の増幅機構に関する数値的研究

Numerical study of the amplification mechanism of the meteo-tsunami originating off the western coast of Kyushu

*福澤 克俊¹、日比谷 紀之¹

*Katsutoshi Fukuzawa¹, Toshiyuki Hibiya¹

1. 東京大学大学院理学系研究科地球惑星科学専攻

1. Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo

A meteo-tsunami is generated by a travelling atmospheric pressure disturbance and is characterized by a rapid sea level rise with a period of several tens of minutes in the coastal areas. In Japan, meteo-tsunamis are frequently observed along the western coast of Kyushu during winter-spring and are called “Abiki” .

In shallow seas such as continental shelf areas, a propagating shallow water wave can be amplified through a resonant coupling to the atmospheric pressure disturbance traveling with nearly the same speed (“the first resonance”). The shallow water wave then enters local coastal areas while being further amplified by “geometrical resonance” effects (“the second resonance”). To predict the meteo-tsunami, therefore, it is essential to clarify the amplification mechanism of the shallow water wave, especially through “the second resonance” effect, in addition to draw the information about the atmospheric pressure disturbance causing “the first resonance” from the synoptic scale atmospheric data.

In this study, we investigate the amplification processes of the meteo-tsunami in Makurazaki Bay that occurred in 2004. The sea level variations in Makurazaki Bay are reproduced well using a barotropic numerical model in which an atmospheric pressure disturbance with a width of about 300 km is assumed to propagate east-southeastward with a constant speed of 31 m/s over the East China Sea. On the continental shelf of the East China Sea, the shallow water wave is amplified through “the first resonance” effect. Then, the shallow water wave propagates further eastward and enters the continental shelf running west to east off Makurazaki Bay. It is shown that the existence of Mt. Kaimon, located at the eastern end of this continental shelf plays an essential role in “the second resonance” ; the reflection of shallow water waves from Mt. Kaimon causes the excitation of the eigen-oscillation on this continental shelf, which resonantly intensifies the eigen-oscillation in Makurazaki Bay.

キーワード：気象津波、九州西方沿岸域、増幅機構

Keywords: Meteo-tsunami, Western coast of Kyushu, Amplification mechanism

Numerical simulations of internal wave dynamics in the vicinity of Izu-Oshima Island, off Sagami Bay, Japan

*増永 英治¹、鈴江 洋太²、内山 雄介²、山崎 秀勝³

*Eiji Masunaga¹, Yota Suzue², Yusuke Uchiyama², Hidekatsu Yamazaki³

1. 茨城大学、2. 神戸大学、3. 東京海洋大学

1. Ibaraki University, 2. Kobe university, 3. Tokyo University of Marine Science and Technology

日本南岸海域における海洋環境は潮汐と黒潮(西岸境界流)の両方が強く影響していることが知られている。近年の研究により、潮汐により励起される非線形性の内部波が沿岸海域において強い流れや混合を引き起こすことがわかってきた。また、黒潮により発生した流れや渦が、伊豆海嶺や島を通過する時に強い物質輸送や混合を引き起こすことが知られている。本研究では、伊豆小笠原海嶺北端に位置する伊豆大島周辺海域を対象海域として内部波の挙動の研究を行った。研究海域の物理構造を再現するためにRegional Ocean Modeling System (ROMS)を用い、3段階のダウンスケーリングを行うことで対象海域の水平解像度を300mに設定した。また内部波の挙動を解析するために、外力として潮汐を与えたモデルと潮汐を考慮しないモデルの2条件の計算を行った。潮汐を外力として与えたモデルでは、潮汐によって励起された非線形性の内部波が再現され強い流れを発生させていた。浅海域(水深250m以下)における運動エネルギーは、潮汐を外力として加えた計算では潮汐を考慮しないモデルに比べ2倍程度高かった。また、内部波のフラックスは潮汐を外力として与えた計算では、約10倍高い結果であった。潮汐は流動場だけではなく水温場にも強い影響を与え、潮汐と内部潮汐によって発生した混合により海嶺と島周辺では水温が2度程度下がることがわかった。本研究による結果から、伊豆大島周辺では潮汐と内部潮汐が流動場にとって支配的であり物理構造に非常に強く関わっていることが明らかとなった。

キーワード：内部波、潮汐、数値計算、黒潮

Keywords: internal waves, tide, numerical simulations, Kuroshio

ツバル国フナフチ環礁ラグーン内の汚濁物質輸送シミュレーション Numerical simulations of mass transports processes in the lagoon of Funafuti Atoll, Tuvalu

*小池 海¹、増永 英治²、横木 裕宗³

*Kai Koike¹, Eiji Masunaga², Hiromune Yokoki³

1. 茨城大学大学院理工学研究科、2. 茨城大学広域水圏科学教育研究センター、3. 茨城大学工学部都市システム工学科
1. Graduate School of Science and Engineering, Ibaraki University, 2. Ibaraki University, Center for water
Environmental Studies, 3. Department of Urban and Civil Engineering College of Engineering, Ibaraki University

サンゴ礁などにより形成される環礁周辺の海洋構造の理解は、サンゴ礁やラグーン内環境の保全にとって必要不可欠である。ツバル国フナフチ環礁では、下水処理施設の整備不足により住宅などから発生した汚濁物質が、ラグーン海岸に流出していることが明らかとなっている(Fujitaら、2013)。しかも、ラグーンと外洋との間での海水交換が生じる場所は、州島の切れ目などごく限られている。したがって、ラグーン内は外洋から隔離されており閉鎖性の高い海域である。州島居住域から流出した汚濁物質は、環礁の形成を担っているサンゴ礁の劣化を引き起こし、ひいては国土の喪失につながる可能性がある。

そこで本研究では、フナフチ環礁ラグーンを対象に海洋数値モデルSUNTANSを用いることでラグーン内の流動構造と物質輸送過程の再現を試みた。典型的な流動構造を再現するために、潮汐に加え6パターンの風速・風向を外力として与えた。また、州島からの汚濁物質を想定したパッシブトレーサーを用い、モデル内での汚濁物質の輸送過程の再現を行った。

計算の結果、ラグーン内の流動場は潮汐に比べ風による影響を強く受けることがわかった。潮流はラグーン中央部では0.008 m/s程度の流れしか発生させなかった。一方、風速5 m/sの風が吹いた場合ラグーン内の流速は0.1 m/sに達した。3パターンの風向条件による計算により、北東の風が吹いた場合に州島から流出した汚濁物質は最も多くラグーン外に輸送された。風速5 m/s・北東風の場合、83%の州島起源の汚濁物質がラグーン外へ20日間で流出した。また、北西の風が吹いた場合にはラグーン内に長時間汚濁物質が滞留し、風速5 m/sの場合では35%の汚濁物質がラグーン外へ20日間で流出する結果となった。本研究による結果から、ラグーン内では風の条件が汚濁物質の輸送に強く影響していることが明らかとなった。