

The Impact of Ocean Surface Currents on Global Eddy Kinetic Energy via the Wind Stress Formulation

*Zhitao Yu¹, Edward Joe Metzger¹

1. US NRL

A more complete wind stress formulation takes into account the ocean surface currents, while the conventional wind stress popularly used in ocean circulation models is only a function of 10-m winds. A pair of 12.5-year (July 2002 –December 2014) HYbrid Coordinate Ocean Model (HYCOM) simulations that only differ in the wind stress formulation are used to study the impact of ocean surface currents on global Eddy Kinetic Energy (EKE). The model results (2004-2014) show that including ocean surface currents in surface wind stress formulation reduces global EKE by more than 40%. To understand the mechanisms behind the large difference, we calculate the global EKE budget using the standard Reynolds averaging procedure. The direct impact of surface wind stress on EKE is through surface wind work. The indirect impact is through changes in the mean circulation that affect the shear production, buoyancy work, and bottom friction. Model Results indicate that the reduction of global EKE is primarily due to the buoyancy work and shear production. Even though the surface eddy wind work is one order larger than buoyancy work, and two orders larger than shear production, it is mainly balanced by bottom friction and thus not the main contribution for the EKE differences.

Keywords: HYCOM, eddy kinetic energy, wind work, baroclinic instability, wind stress

Long-term change and variation of salinity in the western North Pacific subtropical gyre revealed by 50-year long observations along 137°E

*Eitarou Oka¹, Shota Katsura¹, Hiroyuki Inoue², Atsushi Kojima^{2,3}, Moeko Kitamoto^{2,3}, Toshiya Nakano^{2,3}, Toshio Suga^{4,5}

1. Atmosphere and Ocean Research Institute, The University of Tokyo, 2. Global Environment and Marine Department, Japan Meteorological Agency, 3. Oceanography and Geochemistry Research Department, Meteorological Research Institute, 4. Department of Geophysics, Graduate School of Science, Tohoku University, 5. Research and Development Center for Global Change, Japan Agency for Marine-Earth Science and Technology

The 137°E repeat hydrographic section for 50 winters during 1967–2016 has been analyzed to examine interannual to interdecadal variations and long-term changes of salinity and temperature in the surface and intermediate layers of the western North Pacific, with a particular focus on freshening in the subtropical gyre. Rapid freshening on both isobars and isopycnals began in mid-1990' s and persisted for the last 20 years in the upper main thermocline/halocline in the western subtropical gyre. In addition, significant decadal variability of salinity existed in the Subtropical Mode Water (STMW), as previously reported for the shallower layers. An analysis of the 144°E repeat hydrographic section during 1984–2013 supplemented by Argo profiling float data in 2014 and 2015 revealed that the freshening trend and decadal variability observed at 137°E originated in the winter mixed layer in the Kuroshio Extension (KE) region and was transmitted southwestward to 137°E 1–2 years later in association with the subduction and advection of STMW. The mechanism of these changes and variations in the source region was further investigated. In addition to the surface freshwater flux in the KE region pointed out by previous studies, the decadal KE variability in association with the Pacific Decadal Oscillation likely contributes to the decadal salinity variability through water exchange between the subtropics and the subarctic across the KE. Interdecadal change in both the surface freshwater flux and the KE state, however, failed to explain the rapid freshening for the last 20 years.

Keywords: Western North Pacific subtropical gyre, Main thermocline/halocline, Long-term change, Decadal variability, Repeat hydrographic section

Interannual and inter-decadal variability of the North Equatorial Countercurrent in the Western Pacific


*Xiao Chen¹, Bo Qiu

1. Hohai University

Abstract

Interannual and longer timescale variations of the North Equatorial Countercurrent (NECC) in the western Pacific are investigated using the multi-decade (1960-2014) hindcast by the Ocean general circulation model for the Earth Simulator (OFES). The OFES-simulated sea level and upper ocean circulation changes show favorable comparisons with available tide gauge data and repeat hydrographic surveys along the 137°E meridian. An empirical orthogonal function (EOF) analysis reveals that the low-frequency NECC variability is dominated by two distinct modes. The first mode fluctuates interannually and shows strengthening and southward migration of the NECC concurrent with the development of El Niño events. Unlike the extra-tropical western Pacific Ocean circulation variability controlled by wind forcing west of the dateline, the interannual NECC variations are forced by equatorial wind forcing cumulative across the entire Pacific basin. The second mode of the NECC variability has a inter-decadal timescale and is characterized by NECC's progressive weakening in strength, migrating poleward, and broadening in width over the past 50 years. These long-term changes in NECC are caused by the corresponding changes in the trade wind system that weakened and expanded poleward in the past half a century across the Pacific basin.

Keywords: Interannual and Inter-decadal Variability, Western Pacific, Northern Equatorial Countercurrent



Seasonal transport variation over the entire Kuroshio region

*ZHENLONG ZHANG¹, Hirohiko Nakamura¹

1. Kagoshima University

As the observations of the Kuroshio were conducted regionally by different groups, studies for the seasonal variation and its mechanism were also proceeded separately. By using the data collected by satellites and assimilation model's outputs, we focus on the seasonal transport variation over the entire Kuroshio region. Our main purpose is to show the temporal and spatial feature of the seasonal transport variation and give an insight for its driving mechanism.

The results from Aviso data reveal that the surface transport variation reaches maximum in July for most of the Kuroshio region, but its minimum appears in different months. Near Luzon Island and in the East China Sea, the minimum values appear in October or November. For the area of Taiwan, most of the minimum values are found in January or February. Time when the minimum values occur also shows propagation signals: it shifts from September to November in Luzon Strait, while moves gradually from November to June south of Japan. Another feature is that the amplitudes of the surface transport variation are large near Luzon Island, Taiwan, and south of Japan, where there is a land boundary to the west of the Kuroshio. The results from EOF analysis, which is conducted by using an assimilation dataset estimated from the Multivariate Ocean Variation Estimation system, also indicate that the transport variation tends to dominate in the areas where the western land boundary exists.

Czeschel et al. (2012) suggested that fast barotropic waves propagate the signals due to wind-induced coastal upwelling (downwelling) southward along the simple slope off the North American coast and contribute largely to the annual cycle of the Florida current transport. While for the North Pacific western boundary, the barotropic waves are considered to propagate along the complicated coastline with large topographic modulation. This might be a possible cause that leads to the regional features of seasonal transport variation revealed in this study.

Keywords: Kuroshio, seasonal transport variation

Mooring Measurement of the Abyssal Circulations in the Western Pacific Ocean

*Jianing Wang¹, Fan Wang¹

1. Institute of Oceanology, Chinese Academy of Sciences

A scientific observing network in the western tropical Pacific has initially been established by the Institute of Oceanology, Chinese Academy of Sciences (IOCAS). Using fifteen moorings that gives unprecedented measurements in the intermediate and abyssal layers, we present multi-timescale variations of the deep ocean circulations prior to and during 2015 El Niño event. The deep ocean velocities increase equatorward with high standard deviation and nearly zero mean. The deep ocean currents mainly flow in meridional direction in the central Philippine Basin, and are dominated by a series of alternating westward and eastward zonal jets in the Caroline Basin. The currents in the deep channel connecting the East and West Mariana Basins mainly flow southeastward. Seasonal variation is present in the deep jets in the Caroline Basin, associating with vertical propagating annual Rossby wave. The high-frequency flow bands are dominated by diurnal, and semi-diurnal tidal currents, and near-inertial currents. The rough topography has a strong influence on the abyssal circulations, including the intensifications in velocity and internal tidal energy, and the formation of upwelling flow.

Keywords: Abyssal Circulation, Western Pacific, Mooring Measurement

Ocean variability at intermediate depth in the western equatorial Pacific

*Yuji Kashino¹, Takuya Hasegawa¹, Fadli Syamusdin²

1. JAMSTEC Japan Agency for Marine-Earth Science and Technology, 2. BPPT

Intermediate ocean variability at the depth of 300 - 750m in the western equatorial Pacific west of 156E is investigated using data from underwater sensors of the Triangle Trans-Ocean Buoy Network (TRITON) buoys and CTD data by Japan Meteorological Agency, and others. Variability is dominant in the western boundary region. Seasonal variability is generally dominant, but, intraseasonal variability is not negligible around 2N, 138E, where the northwestward flowing New Guinea Coastal Undercurrent retroflects to the east as the Equatorial Undercurrent. There is the anti-cyclonic New Guinea Eddy, which interacts with neighbor eddies around this location. Interannual variability with ENSO and quasi-decadal time scales was observed. Because the pattern of correlations between temperature variability at the intermediate layer and indices of ENSO and quasi-decadal time scales is not simple, variability with these time scale is thought to be associated with thermocline structure variability due to the equatorial waves rather than zonal shift of the warm water pool. We also found freshening trend in the intermediate waters in the western equatorial Pacific. This is likely associated with the freshening of the North Pacific Intermediate Water and Antarctic Intermediate Water.

Keywords: Intermediate ocean variability, Western equatorial Pacific, TRITON buoy, Intermediate waters

Seasonal variability of salinity in the Indian Ocean in response to riverine freshwater forcing using ROMS

*K. K. Sandeep¹, Vimlesh Pant¹

1. Indian Institute of Technology Delhi

Indian Ocean exhibits large spatial and temporal variability in the sea surface salinity distribution. The unique geographical features causes the salinity distribution over the Arabian sea and Bay of Bengal quite different. The Arabian sea being connected to highly saline waters from Red Sea and Persian Gulf remains salty when compared to the Bay of Bengal. Major rivers such as Ganges, Brahmaputra, Mahanadi, Godavari, Krishna, and Irrawaddy debouch an excess amount of freshwater into the Bay. In addition to this, the annual precipitation over Bay of Bengal always remains higher than evaporation when compared to Arabian Sea. Due to these factors the Bay remains fresher than Arabian sea throughout the year and also exhibits large seasonal variability in the circulation patterns. The seasonal variability of sea surface salinity in the Indian ocean is studied by using a ocean general circulation model 'Regional Ocean Modeling System' (ROMS). The model domain extends from 30° S-30°N, 30°E-120°E with quarter degree resolution in the horizontal and 40 sigma levels. The model is initialized with annual mean climatology of tracer fields from World Ocean Atlas 2013 (WOA13) and forced with daily climatological winds from Quikscat and ASCAT and other atmospheric forcing fields from TropFlux. It is found that there is significant contribution to freshening from riverine freshwater discharge in coastal regions near river mouths. Over the Indian Ocean, northern boundaries of Bay of Bengal found to have largest impact of river discharges on SSS simulations during monsoon (June-September) season when the peak discharge from major rivers Ganges, Brahmaputra, and Irrawaddy accumulate freshwater plume along coastal regions in Bay of Bengal. The impact of continental freshwater discharge on the seasonal salinity pattern are investigated. Satellite and in-situ observations are also used to understand the response of riverine freshwater forcing in controlling the spatial and seasonal variations of salinity in various sectors of the Indian Ocean.

Keywords: Indian Ocean, Sea Surface Salinity, Seasonal Variability, River Discharge

Numerical study of Internal tide energetics in the Andaman Sea

*Sachiko Mohanty¹, A D Rao¹

1. Indian Institute of Technology Delhi

The interactions of barotropic tides with irregular bottom topography generate internal waves with high amplitude known as large-amplitude internal waves (LAIW). The Andaman Sea is one of the potential region where such waves occur. These waves are an important phenomena in the ocean due to their influence on the density structure and energy transfer into the region. These waves are also important in submarine acoustics, underwater navigation, offshore structures, ocean mixing, biogeochemical processes, etc. over the shelf-slope region.

A three-dimensional MIT general circulation ocean model (MITgcm) is configured over the Andaman sea to investigate the generation and propagation of M_2 internal tides. Initially, the model simulations are validated by using *in-situ* observations of temperature, conductivity and currents from a buoy located at 10.5N, 94E. The spectral energy estimate of density shows that the peak estimate is associated with the semi-diurnal frequency at all the depths in both observations and model simulations. The vertical structure of baroclinic velocity is compared with observations and the analysis suggests that a multi-mode features of baroclinic tides are present at the buoy location. To understand the generation and propagation of internal tides over this region, energy flux and barotropic-to-baroclinic M_2 tidal energy conversion rates are also estimated. The model simulation suggests that the internal tides are mainly generated at the north of Sumatra coast and the regions around the Nicobar islands. The internal tides propagate away from the respective generation sites. The steepness of topography suggests that the internal tide mainly generate at the supercritical slope region and the energy flux reflected back to the deep water from the supercritical slope. The M_2 dissipation rate is also found to be maximum at the generation sites.

Keywords: Internal tide, Andaman Sea, Energetics

Simulation of storm surges by improving the cyclonic wind formulation

*Smita Pandey¹, A D Rao¹

1. Indian Institute of Technology Delhi

Coastal regions of India are highly prone to sea level rise during tropical cyclones. Storm surges due to cyclones can produce high sea levels, especially when their occurrence coincide with the high tides. Storm surges, tides and wind waves are the main factors responsible for the variation in the total water elevation (TWE) near the coast. The accuracy in simulation of storm surges depends not only on the cyclonic track and its intensity, but also on the spatial distribution of winds which include its speed and direction. In the present study, the cyclonic winds are validated using buoy winds for the recent cyclones formed in the Bay of Bengal since 2010 using Jelesnianski wind scheme. It is found that the cyclonic winds computed from the scheme show an underestimate in the magnitude and also a mismatch in its direction. Hence, the wind scheme is suitably modified based on the buoy observations available at different locations using a power law which reduces the exponential decay of winds by about 30%. Moreover, the cyclonic wind direction is also corrected by suitably modifying its inflow angle. The significance of modified exponential factor and inflow angle in the computation cyclonic winds is highlighted using statistical analysis.

In the present study, simulation of storm surges for the recent cyclonic cases along the east coast of India are carried out using both stand-alone hydrodynamical depth integrated ADCIRC model and coupled ADCIRC+SWAN model. The coupled ADCIRC+SWAN model is used to incorporate the contribution of wind waves in the simulation of TWE near the coast. The cyclonic wind distribution is computed using both modified and unmodified Jelesnianski wind scheme. The experiments are performed to validate the TWE generated from the cyclones through computation of surge residuals with the available tide gauge data. On comparison of observed surge residuals with the simulations using modified winds from the uncoupled and coupled models, it is found that the simulated surge residuals are better compared, especially with the inclusion of wave effect through the coupled model.

Keywords: total water elevations, storm surge, cyclonic winds, ADCIRC MODEL

Numerical study of the amplification mechanism of the meteo-tsunami originating off the western coast of Kyushu

*Katsutoshi Fukuzawa¹, Toshiyuki Hibiya¹

1. Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo

A meteo-tsunami is generated by a travelling atmospheric pressure disturbance and is characterized by a rapid sea level rise with a period of several tens of minutes in the coastal areas. In Japan, meteo-tsunamis are frequently observed along the western coast of Kyushu during winter-spring and are called “Abiki” .

In shallow seas such as continental shelf areas, a propagating shallow water wave can be amplified through a resonant coupling to the atmospheric pressure disturbance traveling with nearly the same speed (“the first resonance”). The shallow water wave then enters local coastal areas while being further amplified by “geometrical resonance” effects (“the second resonance”). To predict the meteo-tsunami, therefore, it is essential to clarify the amplification mechanism of the shallow water wave, especially through “the second resonance” effect, in addition to draw the information about the atmospheric pressure disturbance causing “the first resonance” from the synoptic scale atmospheric data.

In this study, we investigate the amplification processes of the meteo-tsunami in Makurazaki Bay that occurred in 2004. The sea level variations in Makurazaki Bay are reproduced well using a barotropic numerical model in which an atmospheric pressure disturbance with a width of about 300 km is assumed to propagate east-southeastward with a constant speed of 31 m/s over the East China Sea. On the continental shelf of the East China Sea, the shallow water wave is amplified through “the first resonance” effect. Then, the shallow water wave propagates further eastward and enters the continental shelf running west to east off Makurazaki Bay. It is shown that the existence of Mt. Kaimon, located at the eastern end of this continental shelf plays an essential role in “the second resonance” ; the reflection of shallow water waves from Mt. Kaimon causes the excitation of the eigen-oscillation on this continental shelf, which resonantly intensifies the eigen-oscillation in Makurazaki Bay.

Keywords: Meteo-tsunami, Western coast of Kyushu, Amplification mechanism

Numerical simulations of internal wave dynamics in the vicinity of Izu-Oshima Island, off Sagami Bay, Japan

*Eiji Masunaga¹, Yota Suzue², Yusuke Uchiyama², Hidekatsu Yamazaki³

1. Ibaraki University, 2. Kobe university, 3. Tokyo University of Marine Science and Technology

Physical processes off the southern part of Japan mainland are strongly influenced by both tides and Kuroshio (eastern boundary current). The tidal forcing generates internal tides, which results in nonlinear internal bores accompanied by strong flows and turbulent mixing. Currents and eddies induced by Kuroshio generate strong mass transports and turbulent mixing when they cross shallow ridges, such as Izu-Ogasawara Ridge. In this study, we focus on internal wave dynamics around Izu-Oshima Island located on Izu-Ogasawara Ridge, off Sagami Bay, Japan. Regional Ocean Modeling System (ROMS) was employed to simulate physical processes in the study area. By using a one-way offline nesting approach, the horizontal grid spacing was down to 300 m around the study area. Two model case were run to investigate internal wave dynamics: (1) numerical run with the tidal forcing, and (2) numerical run without tidal forcing. Numerical results from the case with the tidal forcing showed strong internal tidal bores generated around the Island, which results in strong currents near the coast. The volume-integrated kinetic energy was 2 times higher in shallow area (depth<250 m) for the case with tides than that of the case without tides. The strength of internal waves (internal wave energy flux) was an order of 10 times higher for the case with tidal forcing than that of the case without tides. In addition, the tidal forcing reduced the surface temperature over the shallow ridge because of strong vertical mixing induced by tidal flows and internal tides. This study indicates that tides and internal tides dominates in the flow field and significantly contribute to oceanic physical processes in the vicinity of Izu-Oshima Island.

Keywords: internal waves, tide, numerical simulations, Kuroshio

Numerical simulations of mass transports processes in the lagoon of Funafuti Atoll, Tuvalu

*Kai Koike¹, Eiji Masunaga², Hiromune Yokoki³

1. Graduate School of Science and Engineering, Ibaraki University, 2. Ibaraki University, Center for water Environmental Studies, 3. Department of Urban and Civil Engineering College of Engineering, Ibaraki University

Understanding oceanic structures in atolls, formed by coral reefs, is essential for maintaining their ecosystems and environment. Contaminated water due to poorly sewage treatment outflows into the lagoon in Funafuti Atoll, Tuvalu (Fujita et al., 2013). However, water exchange between the lagoon and open ocean is limited through narrow channels on the coral reef. Therefore, the sea water in the lagoon is isolated from the open ocean and the lagoon is a highly closed ocean area. Contaminated water discharged from the urban area would causes deterioration of coral reefs, which leads to the changes in the form of the atoll and the loss of the land.

This study attempted to reproduce flow structures and mass transport processes in Funafuti Atoll lagoon by using the SUNTANS model (Fringer et al., 2006). In order to reproduce the typical flow structure in the lagoon, the model was forced by tides and six wind conditions (wind speed and direction). In addition, transport processes of contaminated water from the urban area were reproduced by a passive tracer model.

Numerical results showed that the wind stress strongly influences the flow field in the lagoon compared to the tidal forcing. In the middle of the lagoon, the tidal current generated weak currents, approximately 0.008 m/s. On the other hand, when the wind speed is 5m/s, the flow speed in the middle of the lagoon reached 0.1 m/s. According to numerical runs with three wind direction patterns, northeast winds enhance transports of contaminated water from the lagoon to the open ocean. 83% contaminated water from the urban area flowed into the open ocean within 20 days by northeast winds with the speed of 5m/s. Under the northwest and 5m/s wind condition, water from the urban area stays inside of the lagoon for a long time. 35% of contaminated water flowed into the open ocean within 20 days in this condition. This study suggests that the wind condition plays a significant role in transport processes of contaminated water between the inside of the lagoon of Funafuti Atoll and the open ocean.

Spatial and temporal variability of temperature and salinity in the global ocean from Argo float data

*Shigeki Hosoda¹, Kanako Sato¹

1. Japan Marine-Earth Science and Technology

Statistical analysis for spatial and temporal variability of oceanic temperature and salinity, especially focusing on decorrelation length and signal/noise ratio, is carried out using long-term accumulated Argo float data.

It has been passed 16 years since International Argo program started. The number of active Argo float exceeded 3000 in late 2007, which is the first target in the Argo program, and then the number of accumulated profile data got to 1,000,000 in 2011. As the global Argo observation array is producing real-time oceanic data each horizontally 3x3 degrees every 10 days, we are able to have a huge amount of temperature and salinity data from sea surface to 2000 dbar in the global ocean. The long-term huge Argo data enable us get statistical information of spatial and temporal variability accurately.

Previously such kind of statistical information were estimated for mainly temperature field using expendable BathyThermograph (XBT) data (e.g., White, 1995; Meyer et al., 1982). However, sampling area of XBT data is temporally and spatially biased because of limited observational or voluntary ship courses. Recently statistical analyses were carried out using a huge amount of Argo float data to investigate statistical structure of longer term spatial variability on oceanic fields of not only temperature but also salinity. For example, Resnyanskii et al. (2010) showed spatial covariance and signal/noise ratio for separated temperature and salinity fields into 12 areas using Argo data for 2005-2007, and discussed the characteristics of statistical structures depending on areas, temperature/salinity and depth qualitatively. However, they directly analyzed only spatial covariance, indirectly guessed temporal one by global average current velocity.

Using 16-year Argo float data, here we show 4-D statistical structures of accurate decorrelation length and signal/noise ratio, referring the analytical method of Resnyanskii et al. (2010).

Used Argo profiling data are delayed mode quality controlled temperature and salinity in Jan. 2001 - Dec. 2016 obtained from the Argo Global Data Assembly Centers (GDACs) and the Advanced Quality Controlled data (AQC), which is well quality controlled dataset for real-time QC Argo data. Anomalies are calculated from monthly climatology of World Ocean Atlas 2013 (WOA2013: Locarnini et al., 2013; Zweng et al., 2013) to remove seasonal variability. Here we separate the global ocean into 25 areas, which is finer than in the previous study. After sorting all profiling data for each areas, the profiling data are interpolated on 23 levels from 10 to 2000 dbar (Akima, 1970). Then covariance of anomalies are calculated for temperature and salinity fields, sorting into 20km bins of zonal and meridional distances between observations from 0 - 3000km, respectively. Temporal covariance is also calculated from sorted profiling data into 5-day bins from Jan. 2001 to Dec. 2016, referring spatial covariance estimated above. Based on these temporal and spatial covariance maps, we then estimate decorrelation lengths for time, horizontal and vertical directions every areas.

The temporal and spatial covariance exponentially decrease with distances in all area, which are qualitatively similar to the previous study except for temporal covariance that is the first time to analyze in this study. However, some differences of spatial structures on decorrelation length and signal/noise ratio in the previous study are detected in the spatial covariance of temperature and salinity, especially in the difference between temperature and salinity, surface and deeper layers. These results are obtained because of sorting finer area separation into 25 areas and using long-term Argo float data.

The basic statistical information in this study is estimated based only on the purely observational data

from the global Argo array. Therefore, this information is expected to be useful for not only applications for objective analyses and optimal interpolation mapping but also improvement and validation of data assimilation and numerical simulation models.

Keywords: Decorrelation length of oceanic variability, Argo, Global ocean

Long-term change of the halocline in the global ocean

*Oda Masato¹, Ueno Hiromichi², Yasui Katsura¹

1. Graduate School of Environmental Science, Hokkaido University, 2. Faculty of Fisheries Sciences Graduate School of Fisheries Sciences, Hokkaido University

Distribution and long-term change of the halocline in the upper layers of the world ocean were investigated via analysis of World Ocean Database 2013 (WOD13) using a simple definition of the halocline. The halocline was observed in the tropics, equatorward subtropical regions, and subpolar regions, but was absent from central subtropical regions. A strong halocline tended to occur in areas where the sea surface salinity (SSS) was low. Long-term change of the halocline was also observed in the world ocean: halocline was strengthened by more than 20% in the last thirty years in the global average. The change was relatively strong in the tropics, equatorward subtropical regions, and subpolar regions and was closely related with long-term change of SSS: the lower the SSS was, the stronger the halocline became. The correlation coefficient between the horizontal distributions of the halocline strength change and SSS change was -0.55 in the area of 60°S-60°N.

Keywords: halocline, long-term change, salinity, global ocean, World Ocean Database

Steric sea level variability from the ocean reanalyses intercomparison project in the East Asian marginal seas

*You-Soon Chang¹, Min-Ji Kang¹

1. Kongju National University

Ocean Reanalyses Intercomparison Project (ORA-IP) is successfully coordinated, but most analyses using this product have been limited to the global scale assessments. In this study, regional assessment on steric height variability has been performed focusing on the East Asian marginal seas. Results show that reanalysis ensemble derived from 15-different assimilation systems depicts higher correlation with satellite observations (altimetry-gravimetry) than objective analysis ensemble. Additional results about the long-term variability for the thermosteric and halosteric components of each product will be presented.

Keywords: Ocean Reanalyses Intercomparison Project (ORA-IP), steric height, East Asian marginal seas

Diagnosing formation of the severe drought in SWC during 2009-2010

*Lin Feng¹

1. The First Institute of Oceanography, State Oceanic Administration, China

An extraordinary severe drought hit Southwest China (SWC) from late 2009 to early 2010 and caused enormous losses. The local long-lasting rainfall deficiency which started several months before and the warmer surface air temperature during this period were considered to be the direct meteorological reasons for this disaster. Station observations covering SWC show that there was an almost unbroken rainfall shortage period from May 2009 to March 2010, including two remarkable spells in 2009, May-Jun-July (MJJ) and the boreal autumn (SON). During the first spell, local anomalous descent in the mid-to-low troposphere plays a dominate role to suppress the convection even the water vapor is enough. Diagnosing the vertical motion equation indicates that the local descent is primarily maintained by the anomalous cold temperature advection processes. Both the advection of mean temperature by anomalous wind and the advection of anomalous temperature by mean wind contribute are important. In SON 2009, a quite different situation is proposed that the less than normal northeastward low-level water vapor transport anomaly instead of the descent motions anomaly lead to the rainfall shortage in SWC. Advanced diagnoses illustrate that the low-level water vapor convergence over SWC is significantly less as the transport reduces and the mean water vapor converged by anomalous wind is mainly responsible for this anomaly. Ultimately, the regional circulation anomaly over SWC is controlled by the concurrent El Niño event, which is distinctive and generates an anomalous cyclone centered over the northwest tropical Pacific Ocean.

Keywords: Rainfall , Southwest China, The tropical Pacific

Identification of the Kuroshio and its extension paths from satellite altimetric measurements.

*Yuya Hirano¹, Kazuyuki Uehara¹

1. Tokai University

The Kuroshio is the western boundary current of the subtropical gyre of the North Pacific Ocean, it flows eastward along the south coast of Japan. The Kuroshio has typical two paths. One is called large-meander path and the other is called non-large-meander path (Kawabe, 1995).

After the Kuroshio is separating from east of the Boso peninsula, we call the eastward jet as the Kuroshio Extension (KE). It is considered that the KE has basically quasi-stationary meanders with two ridges located at 144 and 150E. Recently, satellite altimetric measurements of the sea surface height field allow us to have been clarified various meander patterns of the KE due to mesoscale eddies and its decadal changes (Qiu and Chen, 2005).

Using sea surface height of satellite altimetric measurements, to determine the path of flows in the ocean is generally very difficult. Determination of the KE path, in particular, is difficult because of existence of many mesoscale eddies.

In this study, using daily gridded absolute dynamic sea surface height (ADH) distributed by AVISO, we try to determine the paths of the Kuroshio/Kuroshio Extension.

Although previous studies have been individually discussed the Kuroshio paths and the KE paths, we here regard the Kuroshio and the KE as one system.

The determination processes of the path are follows. (1) the daily ADH fields and the surface geostrophic flows derived from those are smoothed by 3-month. (2) the maximum absolute geostrophic flows are detected by scanning north-south at the intervals of 1/4 degree longitude from 130E to 140E. The ADHs corresponding the maximum absolute geostrophic flow detected at each longitude are averaged. This mean ADH from 130E to 140E is determined as the ADH which show the Kuroshio path. (3) the start point of the Kuroshio path is determined by the mean ADH on the 126E longitude line. After scanning the south-north along longitude lines and scanning the east-west along latitude lines in order, the next point is selected the shortest distance from among the position of the same values of ADH. (4) this method is repeated to 170E. One path of the Kuroshio/Kuroshio Extension at a instantaneous section is obtained in this way.

We discuss on the statistics of the Kuroshio/Kuroshio Extension paths obtained by the above method.

Keywords: The Kuroshio, The Kuroshio Extension

Internal tides along the Sanriku Coast, Japan

*Kei Nishina¹, Kiyoshi Tanaka¹, Daigo Yanagimoto¹, Hajime Nishigaki²

1. Tokyo University, 2. Oita University

Hydrographic observations made in several bays (e.g. Otsuchi bay, Toni bay) on the Sanriku Coast, Japan, detected that baroclinic circulations extending over the bays intermittently occur (e.g. Okazaki, 1990, 1994; Tanaka et al. 2016). Previous studies suggest that the main source of this baroclinic circulation is the internal tides (Okazaki, 1990, 1994; Otake et al., 1996, 2009; Ito et al., 1998), however, detailed mechanisms are not clear. In the present study, we perform numerical experiments with a two-dimensional (x-z) nonhydrostatic model (Akitomo et al. 1995) in order to investigate the internal tides along the Sanriku Coast focused on the amplification process of the baroclinic circulation over the bays. Experimental results (Figure 1) indicate that the internal waves enhance the baroclinic circulations over the bays. This internal waves are generated at the shelf break and propagate to the bay mouth.

Keywords: internal tide, Sanriku, shelf-break, nonhydrostatic model




Figure 1. Experimental results of baroclinic cross-shore velocity (color tone, cm/s) and potential density (contour, contour interval 0.5 kg m^{-3})