

## Impact of terrigenous materials on satellite bio-optical variables in the Bay of Bengal

\*Eko Siswanto<sup>1</sup>, Md Latifur Rahman Sarker<sup>2,3</sup>

1. JAMSTEC, 2. Univ. Tek. Malaysia, 3. Rajshahi Univ.

Transported by atmospheric deposition and river discharge, the Bay of Bengal (BoB) largely receives terrigenous iron-contained alluvial dust and dissolved inorganic nitrogen, the micro-macro nutrients which fuel phytoplankton growth and ocean primary productivity (e.g., Jutla et al. 2011; Srinivas et al. 2012; Narvekar and Kumar 2014; Sarangi 2016). Besides nutrients, river discharge also supplies the BoB suspended sediment and dissolved organic carbon, the sizable presence of which may be detrimental for phytoplankton growth and primary productivity. To date however, there is no information how the long-term river discharge variability may interannually affect the above-said in-water bio-optical variables in the BoB. In this analysis, by utilizing satellite-retrieved remote sensing reflectance (Rrs), the optical variables of chlorophyll-a concentration (Chl-a, a proxy for ocean primary productivity), total suspended matter concentration (TSM), and colored dissolved organic carbon absorption coefficient (CDOM) were derived within the period from September 1997 to December 2015. In-situ Ganges-Brahmaputra River (GBR) discharge (a proxy for nutrient input), and satellite-derived rain rate (RR) within the same time-span were also analyzed, and their interannual impacts on the ocean color variables were identified. The flood season in the GBR basin is in summer with the peak (in August) approximately lagged RR (in July) by one month. Chl-a, TSM, and CDOM in the BoB had larger non-seasonal component variances than seasonal ones, indicating that they might be sensitive to climate changes. Observing meridional distributions, the coastal water (with high bio-optical variables) can be roughly separated from the open ocean (with low bio-optical variables) approximately at 21°N. In summer of the years 1998, 2007, and 2011 however, it was obvious that bio-optical variables dispersed more southward to occupy the open ocean. In the same periods, RR and GBR discharge were remarkably higher than those during the normal summer season. This might indicate that high GBR discharge caused by high RR during summers 1998, 2007, and 2011 supplied terrigenous nutrients, suspended solid, and dissolved organic carbon into the BoB. In addition, the summers in 1998, 2007, and 2011 were the moderate La Niña periods, and it is well known that summer rainfall in the upstream GBR basin was anomalously high during La Niña years (e.g., Chowdhury 2003; Cai et al. 2015). Even though the aforesaid results showed increments of bio-optical variables through La Niña teleconnections with RR and GBR streamflow, further analysis is required to clarify whether any factors other than GBR discharge may also modulate the bio-optical variables. This is because, strengthened physical factors such as, wind speed, surface current, and vertical mixing, may entrain nutrients, suspended sediment, and dissolved organic carbon from deeper layer. Terrigenous alluvial dust deposition may also fertilize the BoB, thereby may also enhance Chl-a at interannual time-scale. La Niña teleconnections with atmospheric alluvial dust, wind speed, current, and vertical mixing in the BoB also need to be investigated. Therefore, we will analyze other geophysical variables such as, satellite-derived aerosol absorbing index (a proxy for alluvial dust), wind speed, surface ocean current, and reanalyzed mixed layer depth data, and identify whether or not those factors also interannually modulate the BoB bio-optical variables.

Keywords: Remote sensing, Ocean color, River discharge, Aerosols, La Nina

## Intraseasonal coastal upwelling signal along the southern coast of Java observed by Indonesian tidal station data

\*堀井 孝憲<sup>1</sup>、植木 巖<sup>1</sup>、Syamsudin Fadli<sup>2</sup>、Sofian Ibnu<sup>3</sup>、安藤 健太郎<sup>1</sup>

\*Takanori Horii<sup>1</sup>, Iwao Ueki<sup>1</sup>, Fadli Syamsudin<sup>2</sup>, Ibnu Sofian<sup>3</sup>, Kentaro Ando<sup>1</sup>

1. 海洋研究開発機構 地球環境観測研究開発センター、2. インドネシア技術評価応用庁、3. インドネシア地理空間情報局

1. Research and Development Center for Global Change (RCGC), Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2. Indonesian Agency for the Assessment and Application of Technology (BPPT), 3. Geospatial Information Agency (BIG)

Understanding coastal upwelling along the southwestern coasts of Sumatra and Java and their impacts on ocean surface heat and biogeochemical balance in the eastern Indian Ocean are the main target of international research collaboration, the Eastern Indian Ocean Upwelling Research Initiative (EIOURI). In this study, analyzing sea level data along the coasts of Sumatra and Java, we investigated the coastal upwelling signal that is linked to local sea surface temperature (SST) variability. We used Indonesian tidal station data together with satellite SST data and atmospheric reanalysis data. The sea level variations along the southern coast of Java have a significant coherence with remote wind, local wind, and local SST variations, with an intraseasonal time scale of 20–50 days. Assuming that a coastal upwelling signal would appear as a sea level drop (SLD), we focused on intraseasonal-scale SLD events in the data. Significant upwelling signals are frequently observed during both the boreal summer and winter. To evaluate the impact of the coastal upwelling on local SST, we examined statistical relationships between sea level and SST variations. The results demonstrated that events that occurred during April-August were associated with local SST cooling. The horizontal distribution of the SST cooling was analogous with annual-mean SST, suggesting the importance of intraseasonal-scale coastal upwelling in forming the climatic conditions of the southeastern tropical Indian Ocean.

キーワード：東部インド洋、沿岸湧昇、スマトラ・ジャワ島、潮位データ、東部インド洋湧昇イニシアティブ  
Keywords: Eastern Indian Ocean, Coastal Upwelling, Sumatra and Java, Tidal Station Data, EIOURI

## Improved prediction of the Indian Ocean Dipole Mode by use of subsurface ocean observations

\*土井 威志<sup>1</sup>、Andrea Storto<sup>2</sup>、Swadhin Behera<sup>1</sup>、山形 俊男<sup>1</sup>

\*Takeshi Doi<sup>1</sup>, Andrea Storto<sup>2</sup>, Swadhin Behera<sup>1</sup>, Toshio Yamagata<sup>1</sup>

1. JAMSTEC, 2. CMCC

1. JAMSTEC, 2. CMCC

The numerical seasonal prediction system using the SINTEX-F ocean-atmosphere coupled model has so far demonstrated good performance of predicting the Indian Ocean Dipole Mode (IOD) despite that the system adopts a relatively simple initialization scheme with nudging only the sea surface temperature (SST). However, it is easily expected that the system is not sufficient to capture the subsurface oceanic precondition. Therefore, we have introduced a new three-dimensional variational ocean data assimilation (3DVAR) method that takes 3D profiles of observed ocean temperature and salinity into account. Since the new system has successfully improved the IOD prediction, the present study has clearly shown that the ocean observational efforts in the tropical Indian Ocean are quite useful for improvement of the IOD prediction and may contribute to rich socio-economic activities, particularly in the Indian Ocean rim countries.

キーワード：インド洋ダイポールモード現象、季節予測、海洋垂表層の観測

Keywords: Indian Ocean Dipole Mode, Seasonal prediction, Subsurface ocean observation

## 赤道インド洋中深層での季節内変動により励起された平均鉛直流 Mean subsurface upwelling induced by intraseasonal variability over the equatorial Indian Ocean

\*尾形 友道<sup>1</sup>、升本 順夫<sup>2</sup>、名倉 元樹<sup>3</sup>

\*Tomomichi Ogata<sup>1</sup>, Yukio Masumoto<sup>2</sup>, Motoki Nagura<sup>3</sup>

1. 筑波大学生命環境科学研究科、2. 東京大学大学院理学系研究科、3. (独) 海洋研究開発機構

1. Faculty of Life and Environmental Sciences, University of Tsukuba, 2. Graduate School of Science, The University of Tokyo, 3. Japan Agency for Marine-Earth Science and Technology

A possible formation mechanism of mean subsurface upwelling along the equator in the Indian Ocean is investigated using a series of hierarchical setting of ocean general circulation model (OGCM) integrations and analytical considerations. In an eddy-resolving OGCM with realistic forcing, mean vertical velocity in the tropical Indian Ocean shows rather strong upwelling, with its maximum on the equator in subsurface layer below the thermocline. Heat budget analysis exhibits that horizontal and vertical heat advection due to currents and temperature deviations from the mean balances with vertical advection due to mean equatorial upwelling. Horizontal heat advection is mostly associated with intraseasonal variability with a period from 3- to 91-day, while contribution from longer period (> 91 days) are small. Sensitivity experiments with a coarse-resolution OGCM further demonstrate that such mean equatorial upwelling cannot be reproduced by seasonal forcing only. Adding the intraseasonal wind forcing, especially meridional wind variability with a period of 15 days, generates significant mean subsurface upwelling on the equator. Further experiments with idealized settings confirm the importance of intraseasonal mixed Rossby-gravity (MRG) waves to generate mean upwelling, which appears along the energy “beam” of the MRG wave. An analytical solution of the MRG indicates that wave-induced temperature advection caused by the MRG waves with upward (downward) phase propagation results in warming (cooling) on the equator. This wave-induced warming (cooling) is shown to balance with the mean equatorial upwelling (downwelling), which is consistent with simulated characteristics in the OGCM experiments.

キーワード：インド洋、季節内変動

Keywords: Indian Ocean, intraseasonal variability

## On the trigger and interannual variability of the Natal Pulse

\*山上 遥航<sup>1</sup>、東塚 知己<sup>1</sup>、Qiu Bo<sup>2</sup>

\*Yoko Yamagami<sup>1</sup>, Tomoki Tozuka<sup>1</sup>, Bo Qiu<sup>2</sup>

1. 東京大学大学院理学系研究科地球惑星科学専攻、2. ハワイ大学マノア校

1. Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo, 2. University of Hawaii at Manoa

Variability of the Agulhas Current in the southwestern Indian Ocean is characterized by its solitary meander event, known as the Natal Pulse, which may trigger the pinch off of the Agulhas Rings in the Agulhas retroflection region. These turbulent motions may influence inter-ocean water mass exchange between Indian and Atlantic Oceans and thus the global climate system.

In this study, the trigger of the Natal Pulse, and the interannual variability in the occurrence of Natal Pulse events are investigated using satellite altimeter data and outputs from the high-resolution OGCM (OFES) simulation.

It is found that most of the Natal Pulse events are triggered by anticyclonic eddies propagating from the upstream region, most of which originates from the Southeast Madagascar Current (SEMC), although eddies from Mozambique Channel also contribute to the Natal Pulse event.

Furthermore, interannual variations in the Natal Pulse occurrence correlate with eddy kinetic energy off the southern coast of Madagascar, which suggests that the Natal Pulse is controlled by anticyclonic eddies from the SEMC region. An automatic eddy tracking method supports this idea, and shows that possible eddy corridor is mainly from the SEMC region. The connection between the Natal Pulse and western boundary current along the Madagascar coast implies the potential contribution of the basin-scale wind field.

キーワード : アガラス海流、Natal Pulse、渦-平均流相互作用、蛇行、中規模渦

Keywords: Agulhas Current, Natal Pulse, Eddy-Mean Flow Interaction, Meander, Mesoscale Eddies

## Decadal climate prediction in the southern Indian Ocean using SINTEX-F2 coupled GCM

\*森岡 優志<sup>1</sup>、土井 威志<sup>1</sup>、Behera Swadhin<sup>1</sup>

\*Yushi Morioka<sup>1</sup>, Takeshi Doi<sup>1</sup>, Swadhin Behera<sup>1</sup>

1. 海洋研究開発機構アプリケーションラボ

1. Application Laboratory, JAMSTEC

Decadal climate variation in the southern Indian Ocean plays a crucial role in rainfall variability over southern Africa through changes in moisture transport. The rainfall increase over southern Africa has a strong relation with warm sea surface temperature (SST) and high sea level pressure (SLP) anomalies in the Southwest Indian Ocean. Despite many efforts devoted to understanding the physical mechanisms, few studies have examined the decadal climate predictability in the southern Indian Ocean. This study aims at investigating decadal climate predictability in the southern Indian Ocean using a state-of-the-art coupled general circulation model (GCM), called SINTEX-F2. Here we performed 12 ensemble decadal hindcast experiments using different initial conditions with SST-nudging scheme, starting from Mar 1st in each year of 1982-2005.

On decadal timescale, the observed SST anomalies averaged in the Southwest Indian Ocean show positive peak in late 1990s and negative peak in late 2000s. Both positive and negative peaks are well reproduced in the decadal hindcast runs initiated in 1994 and 1999, respectively. Particularly, the decadal hindcast experiments initiated from 1999 successfully capture a distinct phase change from the positive to negative peak. In addition, the spatial patterns of warm SST and high SLP anomalies observed in late 1990s show a clear eastward propagation from the South Atlantic to the southern Indian Ocean along the Antarctic Circumpolar Current. The decadal hindcast experiments initiated in 1994 successfully capture this observed eastward propagation of SST and SLP anomalies from the South Atlantic, indicating an important role of SST anomalies in the South Atlantic. These results suggest that the SINTEX-F2 decadal hindcast experiments, with simple SST-nudging initialization, have reasonable skills in predicting the decadal climate variability of the southern Indian Ocean, important for the southern African climate.

キーワード：十年規模変動、南インド洋、海面水温

Keywords: Decadal variability, Southern Indian Ocean, Sea surface temperature

# ベンガル湾の生物地球化学におけるエアロゾルの役割

## Impact of aerosol on biogeochemistry in the Bay of Bengal

\*本多 牧生<sup>1</sup>、松本 和彦<sup>1</sup>、Siswanto Eko<sup>1</sup>、金谷 有剛<sup>1</sup>、竹谷 文一<sup>1</sup>、宮川 拓真<sup>1</sup>  
\*Makio Honda<sup>1</sup>, Kazuhiko Matsumoto<sup>1</sup>, Eko Siswanto<sup>1</sup>, Yugo Kanaya<sup>1</sup>, Fumikazu Taketani<sup>1</sup>,  
Takuma Miyakawa<sup>1</sup>

1. 海洋研究開発機構

1. Japan Agency for Marine-Earth Science and Technology

陸から海洋へ供給される大気塵（エアロゾル）が海洋の様々な生物地球化学に影響を与えていることが報告されてきた。現代のみならず氷期においても、陸域起源のエアロゾルが海洋の栄養塩濃度を増加させ、基礎生産力、沈降粒子量の増加を誘引する可能性が指摘されている。一方、陸域エアロゾルの海洋供給は有害金属を海洋へ供給したり、海洋の酸性化を促進させたり等負の効果も考えられる。インド洋北東部に位置するベンガル湾は北側および東西を陸に囲まれた半閉鎖的な熱帯海域である。強い日射に加え、ガンジス川、ブラマプトラ川という世界最大級河川による淡水流入、さらに北半球夏季モンスーン（南西風）による多量の降雨により成層構造が発達するため亜表層からの栄養塩供給が極めて少ない貧栄養海域である。このベンガル湾において海洋表層の基礎生産力を高めるための栄養塩供給メカニズムとしては、海洋内部の中規模渦、サイクロンのような気象擾乱に加え、陸域エアロゾル供給が挙げられる。インド洋特有のモンスーンシステムにより、冬季は北東風が卓越し陸から海に向かって風が吹くため、陸域から天然起源あるいはバイオマスバーニングや化石燃料使用による人為起源のエアロゾルが大量にベンガル湾に供給される。ベンガル湾の特徴として、湾周辺国には世界人口の約1/4が集中している。特にインドやバングラディッシュは大気汚染度（PM2.5濃度）が高いことで知られており、大量の人為起源のエアロゾルがベンガル湾周辺で発生し大気中へ放出されている。アラビア海に比べると、春季はより多くの天然起源・あるいは人為起源のエアロゾルが供給されやすい海域であるとの報告がある。人為起源エアロゾルに含まれるmicronutrientである鉄は、天然起源エアロゾルに比較すると、濃度が高く、かつ溶出しやすい（換言すれば、生物に利用されやすい）との報告がある。したがって将来的にはより多くの鉄が海洋に供給された結果、鉄不足の海域では基礎生産力が増加する可能性がある。またベンガル湾南東部に位置するインドネシアからも山火事、火山噴火、バイオマスバーニングによる大量のエアロゾルが発生し同海域の海洋の生物地球化学に影響を与えている可能性も指摘されている。大量のエアロゾルがベンガル湾海上大気に輸送された場合は日射を遮断し、海洋の一次生産力を低下させる効果もあるはずである。従って生物地球化学におけるエアロゾルの役割を研究する上でベンガル湾は“ホットスポット”と言える。本発表ではベンガル湾の生物地球化学におけるエアロゾルの役割に関する研究例をまとめ、2018年に計画されている観測研究航海について紹介する。

キーワード：ベンガル湾、天然起源・人為起源エアロゾル、生物地球化学

Keywords: Bay of Bengal, natural / anthropogenic aerosol, biogeochemistry

## Physical-biogeochemical variability in the Sumatra-Java upwelling region

\*植木 巖<sup>1</sup>、笹井 義一<sup>1</sup>、笹岡 晃征<sup>1</sup>

\*Iwao Ueki<sup>1</sup>, Yoshikazu Sasai<sup>1</sup>, Kousei Sasaoka<sup>1</sup>

1. 国立研究開発法人海洋研究開発機構

1. JAMSTEC Japan Agency for Marine-Earth Science and Technology

The south of Sumatra-Java is one of the greatest upwelling regions in the southeastern tropical Indian Ocean (SETIO). The upwelling appear under the monsoonal wind, thus it dominates from September to November in usual. However, action-center of the Indian Ocean Dipole (IOD) events, which has interannual variability, also appears in SETIO. Then the seasonality of the upwelling should be modulated by IOD events. In general, the upwelling bring rich nutrient from deeper ocean to the surface, thus biological primary production becomes large at that time. This kind of biogeochemical response against physical condition should also be modulated by monsoon and IOD activities. Although the physical interpretations of IOD mechanism and ocean responses of the Monsoonal wind have been improved by development of observing system and numerical modeling technique in the past decade, understanding of the biogeochemical responses of that has not been so improved because of lack of in situ observations. At the present, the second international Indian Ocean expedition (IIOE-2) is conducted under the international observation efforts. It is good chance to progress our understanding of the physical and biogeochemical features at the Sumatra-Java upwelling region. In this presentation we will introduce our observational and numerical analysis results for the Sumatra-Java upwelling region, which mostly focused on the sea surface temperature and phytoplankton variability.

キーワード：インド洋、湧昇、国際インド洋調査2、東部インド洋湧昇研究イニシャティブ

Keywords: Indian Ocean, Upwelling, IIOE2, EIOURI



## 国際インド洋調査2（IIOE-2）と東部インド洋湧昇研究イニシャティブ（EIOURI）

### International Indian Ocean Expedition-2 (IIOE-2) and Eastern Indian Ocean Upwelling Research Initiative (EIOURI)

\*升本 順夫<sup>1</sup>

\*Yukio Masumoto<sup>1</sup>

1. 東京大学大学院理学系研究科

1. Graduate School of Science, The University of Tokyo

1960年代前半に行われた国際インド洋調査から半世紀が経ち、海洋観測技術も大きく発展した現在でも、インド洋の物理、化学、生物研究には多くの未解明なテーマが残されている。また、人類活動が地球環境に影響を与えていることも明らかとなり、インド洋を含む海洋環境への影響も懸念されている。このような状況のもと、インド洋の総合的な観測を主体とした研究計画として国際インド洋調査-2が企画され、2015年12月から開始されている。その中で、特に海洋湧昇域の重要性が注目され、東部および西部インド洋の湧昇域研究イニシャティブ（EIOURIおよびWIOURI）がIIOE-2の重要な研究プロジェクトとして始まった。これらのプロジェクトでは、物理、化学、生物学的視点から湧昇システムの解明と、これらの間の多様な相互関連性を明らかにすることを目的としている。本発表では、IIOE-2とEIOURIの詳細を紹介する。

キーワード：国際インド洋調査、湧昇システム、分野間連携

Keywords: International Indian Ocean Expedition, Upwelling system, Interdisciplinary collaboration