

Impact of terrigenous materials on satellite bio-optical variables in the Bay of Bengal

*Eko Siswanto¹, Md Latifur Rahman Sarker^{2,3}

1. JAMSTEC, 2. Univ. Tek. Malaysia, 3. Rajshahi Univ.

Transported by atmospheric deposition and river discharge, the Bay of Bengal (BoB) largely receives terrigenous iron-contained alluvial dust and dissolved inorganic nitrogen, the micro-macro nutrients which fuel phytoplankton growth and ocean primary productivity (e.g., Jutla et al. 2011; Srinivas et al. 2012; Narvekar and Kumar 2014; Sarangi 2016). Besides nutrients, river discharge also supplies the BoB suspended sediment and dissolved organic carbon, the sizable presence of which may be detrimental for phytoplankton growth and primary productivity. To date however, there is no information how the long-term river discharge variability may interannually affect the above-said in-water bio-optical variables in the BoB. In this analysis, by utilizing satellite-retrieved remote sensing reflectance (Rrs), the optical variables of chlorophyll-a concentration (Chl-a, a proxy for ocean primary productivity), total suspended matter concentration (TSM), and colored dissolved organic carbon absorption coefficient (CDOM) were derived within the period from September 1997 to December 2015. In-situ Ganges-Brahmaputra River (GBR) discharge (a proxy for nutrient input), and satellite-derived rain rate (RR) within the same time-span were also analyzed, and their interannual impacts on the ocean color variables were identified. The flood season in the GBR basin is in summer with the peak (in August) approximately lagged RR (in July) by one month. Chl-a, TSM, and CDOM in the BoB had larger non-seasonal component variances than seasonal ones, indicating that they might be sensitive to climate changes. Observing meridional distributions, the coastal water (with high bio-optical variables) can be roughly separated from the open ocean (with low bio-optical variables) approximately at 21°N. In summer of the years 1998, 2007, and 2011 however, it was obvious that bio-optical variables dispersed more southward to occupy the open ocean. In the same periods, RR and GBR discharge were remarkably higher than those during the normal summer season. This might indicate that high GBR discharge caused by high RR during summers 1998, 2007, and 2011 supplied terrigenous nutrients, suspended solid, and dissolved organic carbon into the BoB. In addition, the summers in 1998, 2007, and 2011 were the moderate La Niña periods, and it is well known that summer rainfall in the upstream GBR basin was anomalously high during La Niña years (e.g., Chowdhury 2003; Cai et al. 2015). Even though the aforesaid results showed increments of bio-optical variables through La Niña teleconnections with RR and GBR streamflow, further analysis is required to clarify whether any factors other than GBR discharge may also modulate the bio-optical variables. This is because, strengthened physical factors such as, wind speed, surface current, and vertical mixing, may entrain nutrients, suspended sediment, and dissolved organic carbon from deeper layer. Terrigenous alluvial dust deposition may also fertilize the BoB, thereby may also enhance Chl-a at interannual time-scale. La Niña teleconnections with atmospheric alluvial dust, wind speed, current, and vertical mixing in the BoB also need to be investigated. Therefore, we will analyze other geophysical variables such as, satellite-derived aerosol absorbing index (a proxy for alluvial dust), wind speed, surface ocean current, and reanalyzed mixed layer depth data, and identify whether or not those factors also interannually modulate the BoB bio-optical variables.

Keywords: Remote sensing, Ocean color, River discharge, Aerosols, La Nina

Intraseasonal coastal upwelling signal along the southern coast of Java observed by Indonesian tidal station data

*Takanori Horii¹, Iwao Ueki¹, Fadli Syamsudin², Ibnu Sofian³, Kentaro Ando¹

1. Research and Development Center for Global Change (RCGC), Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2. Indonesian Agency for the Assessment and Application of Technology (BPPT), 3. Geospatial Information Agency (BIG)

Understanding coastal upwelling along the southwestern coasts of Sumatra and Java and their impacts on ocean surface heat and biogeochemical balance in the eastern Indian Ocean are the main target of international research collaboration, the Eastern Indian Ocean Upwelling Research Initiative (EIOURI). In this study, analyzing sea level data along the coasts of Sumatra and Java, we investigated the coastal upwelling signal that is linked to local sea surface temperature (SST) variability. We used Indonesian tidal station data together with satellite SST data and atmospheric reanalysis data. The sea level variations along the southern coast of Java have a significant coherence with remote wind, local wind, and local SST variations, with an intraseasonal time scale of 20–50 days. Assuming that a coastal upwelling signal would appear as a sea level drop (SLD), we focused on intraseasonal-scale SLD events in the data. Significant upwelling signals are frequently observed during both the boreal summer and winter. To evaluate the impact of the coastal upwelling on local SST, we examined statistical relationships between sea level and SST variations. The results demonstrated that events that occurred during April–August were associated with local SST cooling. The horizontal distribution of the SST cooling was analogous with annual-mean SST, suggesting the importance of intraseasonal-scale coastal upwelling in forming the climatic conditions of the southeastern tropical Indian Ocean.

Keywords: Eastern Indian Ocean, Coastal Upwelling, Sumatra and Java, Tidal Station Data, EIOURI

Improved prediction of the Indian Ocean Dipole Mode by use of subsurface ocean observations

*Takeshi Doi¹, Andrea Storto², Swadhin Behera¹, Toshio Yamagata¹

1. JAMSTEC, 2. CMCC

The numerical seasonal prediction system using the SINTEX-F ocean-atmosphere coupled model has so far demonstrated good performance of predicting the Indian Ocean Dipole Mode (IOD) despite that the system adopts a relatively simple initialization scheme with nudging only the sea surface temperature (SST). However, it is easily expected that the system is not sufficient to capture the subsurface oceanic precondition. Therefore, we have introduced a new three-dimensional variational ocean data assimilation (3DVAR) method that takes 3D profiles of observed ocean temperature and salinity into account. Since the new system has successfully improved the IOD prediction, the present study has clearly shown that the ocean observational efforts in the tropical Indian Ocean are quite useful for improvement of the IOD prediction and may contribute to rich socio-economic activities, particularly in the Indian Ocean rim countries.

Keywords: Indian Ocean Dipole Mode, Seasonal prediction, Subsurface ocean observation

Mean subsurface upwelling induced by intraseasonal variability over the equatorial Indian Ocean

*Tomomichi Ogata¹, Yukio Masumoto², Motoki Nagura³

1. Faculty of Life and Environmental Sciences, University of Tsukuba, 2. Graduate School of Science, The University of Tokyo, 3. Japan Agency for Marine-Earth Science and Technology

A possible formation mechanism of mean subsurface upwelling along the equator in the Indian Ocean is investigated using a series of hierarchical setting of ocean general circulation model (OGCM) integrations and analytical considerations. In an eddy-resolving OGCM with realistic forcing, mean vertical velocity in the tropical Indian Ocean shows rather strong upwelling, with its maximum on the equator in subsurface layer below the thermocline. Heat budget analysis exhibits that horizontal and vertical heat advection due to currents and temperature deviations from the mean balances with vertical advection due to mean equatorial upwelling. Horizontal heat advection is mostly associated with intraseasonal variability with a period from 3- to 91-day, while contribution from longer period (> 91 days) are small. Sensitivity experiments with a coarse-resolution OGCM further demonstrate that such mean equatorial upwelling cannot be reproduced by seasonal forcing only. Adding the intraseasonal wind forcing, especially meridional wind variability with a period of 15 days, generates significant mean subsurface upwelling on the equator. Further experiments with idealized settings confirm the importance of intraseasonal mixed Rossby-gravity (MRG) waves to generate mean upwelling, which appears along the energy “beam” of the MRG wave. An analytical solution of the MRG indicates that wave-induced temperature advection caused by the MRG waves with upward (downward) phase propagation results in warming (cooling) on the equator. This wave-induced warming (cooling) is shown to balance with the mean equatorial upwelling (downwelling), which is consistent with simulated characteristics in the OGCM experiments.

Keywords: Indian Ocean, intraseasonal variability

On the trigger and interannual variability of the Natal Pulse

*Yoko Yamagami¹, Tomoki Tozuka¹, Bo Qiu²

1. Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo, 2. University of Hawaii at Manoa

Variability of the Agulhas Current in the southwestern Indian Ocean is characterized by its solitary meander event, known as the Natal Pulse, which may trigger the pinch off of the Agulhas Rings in the Agulhas retroflection region. These turbulent motions may influence inter-ocean water mass exchange between Indian and Atlantic Oceans and thus the global climate system.

In this study, the trigger of the Natal Pulse, and the interannual variability in the occurrence of Natal Pulse events are investigated using satellite altimeter data and outputs from the high-resolution OGCM (OFES) simulation.

It is found that most of the Natal Pulse events are triggered by anticyclonic eddies propagating from the upstream region, most of which originates from the Southeast Madagascar Current (SEMC), although eddies from Mozambique Channel also contribute to the Natal Pulse event.

Futhermore, interannual variations in the Natal Pulse occurrence correlate with eddy kinetic energy off the southern coast of Madagascar, which suggests that the Natal Pulse is controlled by anticyclonic eddies from the SEMC region. An automatic eddy tracking method supports this idea, and shows that possible eddy corridor is mainly from the SEMC region. The connection between the Natal Pulse and western boundary current along the Madagascar coast implies the potential contribution of the basin-scale wind field.

Keywords: Agulhas Current, Natal Pulse, Eddy-Mean Flow Interaction, Meander, Mesoscale Eddies

Decadal climate prediction in the southern Indian Ocean using SINTEX-F2 coupled GCM

*Yushi Morioka¹, Takeshi Doi¹, Swadhin Behera¹

1. Application Laboratory, JAMSTEC

Decadal climate variation in the southern Indian Ocean plays a crucial role in rainfall variability over southern Africa through changes in moisture transport. The rainfall increase over southern Africa has a strong relation with warm sea surface temperature (SST) and high sea level pressure (SLP) anomalies in the Southwest Indian Ocean. Despite many efforts devoted to understanding the physical mechanisms, few studies have examined the decadal climate predictability in the southern Indian Ocean. This study aims at investigating decadal climate predictability in the southern Indian Ocean using a state-of-the-art coupled general circulation model (GCM), called SINTEX-F2. Here we performed 12 ensemble decadal hindcast experiments using different initial conditions with SST-nudging scheme, starting from Mar 1st in each year of 1982-2005.

On decadal timescale, the observed SST anomalies averaged in the Southwest Indian Ocean show positive peak in late 1990s and negative peak in late 2000s. Both positive and negative peaks are well reproduced in the decadal hindcast runs initiated in 1994 and 1999, respectively. Particularly, the decadal hindcast experiments initiated from 1999 successfully capture a distinct phase change from the positive to negative peak. In addition, the spatial patterns of warm SST and high SLP anomalies observed in late 1990s show a clear eastward propagation from the South Atlantic to the southern Indian Ocean along the Antarctic Circumpolar Current. The decadal hindcast experiments initiated in 1994 successfully capture this observed eastward propagation of SST and SLP anomalies from the South Atlantic, indicating an important role of SST anomalies in the South Atlantic. These results suggest that the SINTEX-F2 decadal hindcast experiments, with simple SST-nudging initialization, have reasonable skills in predicting the decadal climate variability of the southern Indian Ocean, important for the southern African climate.

Keywords: Decadal variability, Southern Indian Ocean, Sea surface temperature

Impact of aerosol on biogeochemistry in the Bay of Bengal

*Makio Honda¹, Kazuhiko Matsumoto¹, Eko Siswanto¹, Yugo Kanaya¹, Fumikazu Taketani¹, Takuma Miyakawa¹

1. Japan Agency for Marine-Earth Science and Technology

It has been reported that aerosol supplied from the land to the ocean affects ocean biogeochemistry. Not only in the modern age, but also during the glacier age, macro- and micro-nutrient in aerosol possibly increases ocean nutrients resulting in increases of primary productivity and settling particles. On the other hand, aerosol input possibly has negative impact: the harmful trace metals in aerosol might damage ocean primary producer and some chemical substances might accelerate the ocean acidification. The Bay of Bengal (BoB) is located in the northeast of the Indian Ocean. The BoB is semi-closed tropical maritime surrounded by the land on its north, east, and west. In addition to strong insolation, large amount of fresh water input from Ganges and Brahmaputra rivers that are one of largest river in the world and large precipitation during the boreal summer monsoon season strongly stratify the ocean. As a result, the supply of nutrient from the subsurface is strictly inhibited and the BoB is oligotrophic. As mechanisms of nutrients supply to the sun-lit layer in the BoB, in addition to passes of meso-scale cyclonic eddy and of meteorology turbulence such as cyclone, aerosol input is very important. Because of the unique monsoon system in the Indian Ocean, the northeast wind is pre-dominant during winter season. Thus, especially in winter, natural and anthropogenic aerosol that originates from the consumption of fossil fuel and the biomass burning are largely transported from the land to the BoB. Another characteristic of the BoB is that about one-quarter of the world population live around the BoB. Especially, Indo and Bangladesh are well known as a country that magnitude of air pollution (concentration of PM_{2.5}) is quite high and gigantic anthropogenic aerosol are emitted. It is reported that, in spring, an amount of anthropogenic aerosol input to the BoB is much higher than that to the Arabian Sea. It has been reported that concentration of iron (Fe) in the anthropogenic aerosol that serves as micronutrient is higher and dissolved more easily (namely, more bioavailable) than that in the natural aerosol. Thus in future, increase of anthropogenic aerosol input might enhance ocean primary productivity in the pelagic ocean with Fe deficiency. Moreover, large amount of aerosol that is attributed to wildfire, volcanic explosion and biomass burning are emitted from Indonesia located in the southeast of the BoB. These aerosols possibly affect biogeochemistry in the BoB too. There is another negative feedback that irradiance to the BoB would decrease resulting decrease of primary productivity if large amount of aerosol were transported over the BoB. Therefore the BoB is a “hot spot” for the study on impact of aerosol on ocean biogeochemistry. This presentation reviews previous study on this issue and introduces a scientific cruise planned in 2018.

Keywords: Bay of Bengal, natural / anthropogenic aerosol, biogeochemistry

Physical-biogeochemical variability in the Sumatra-Java upwelling region

*Iwao Ueki¹, Yoshikazu Sasai¹, Kousei Sasaoka¹

1. JAMSTEC Japan Agency for Marine-Earth Science and Technology

The south of Sumatra-Java is one of the greatest upwelling regions in the southeastern tropical Indian Ocean (SETIO). The upwelling appear under the monsoonal wind, thus it dominates from September to November in usual. However, action-center of the Indian Ocean Dipole (IOD) events, which has interannual variability, also appears in SETIO. Then the seasonality of the upwelling should be modulated by IOD events. In general, the upwelling bring rich nutrient from deeper ocean to the surface, thus biological primary production becomes large at that time. This kind of biogeochemical response against physical condition should also be modulated by monsoon and IOD activities. Although the physical interpretations of IOD mechanism and ocean responses of the Monsoonal wind have been improved by development of observing system and numerical modeling technique in the past decade, understanding of the biogeochemical responses of that has not been so improved because of lack of in situ observations. At the present, the second international Indian Ocean expedition (IIOE-2) is conducted under the international observation efforts. It is good chance to progress our understanding of the physical and biogeochemical features at the Sumatra-Java upwelling region. In this presentation we will introduce our observational and numerical analysis results for the Sumatra-Java upwelling region, which mostly focused on the sea surface temperature and phytoplankton variability.

Keywords: Indian Ocean, Upwelling, IIOE2, EIOURI

International Indian Ocean Expedition-2 (IIOE-2) and Eastern Indian Ocean Upwelling Research Initiative (EIOURI)

*Yukio Masumoto¹

1. Graduate School of Science, The University of Tokyo

It has been half a century since the International Indian Ocean Expedition (IIOE) took place in 1960' s. Although in situ and remotely sensed observing technology have been improved significantly, our understanding of physical, chemical, and biological oceanography in the Indian Ocean is still quite limited. It has also been discussed that recent human activities affect ocean environment, including conditions in the Indian Ocean. Considering such situations, the second IIOE (IIOE-2) was planned and launched in Dec. 2015 to investigate Indian Ocean conditions and their variations from physical, chemical, biological, and geological viewpoints and to understand their mutual interactions. Upwelling system is among important processes to be highlighted under IIOE-2, and Eastern and Western Indian Ocean Upwelling Research Initiatives (EIOURI and WIOURI) have been established as key research initiatives. Detailed descriptions of IIOE-2 and EIOURI will be presented.

Keywords: International Indian Ocean Expedition, Upwelling system, Interdisciplinary collaboration