

酸素発生型光合成の痕跡は3.2億年前のフィグツリー層群に記録されているか？

Early oxygenic photosynthesis in 3.2 Ga Fig Tree Group, Barberton Greenstone Belt?

*掛川 武¹

*Takeshi Kakegawa¹

1. 東北大学大学院理学研究科地学専攻

1. Graduate School of Science, Tohoku University

Reconstructing Archean ecosystem is important to constrain the evolution of early life. Less metamorphosed sedimentary rocks in ca. 3.2 Ga Fig Tree Group gives us an opportunity to examine where and how the early life habituated. Here I report results of geological and geochemical studies on Fig Tree Group, in particular for Jesefsdal (southern section) and Waggon Road Mine (northern section) areas. For both areas, clastic sedimentary rocks were dominant at the base. Northern section is more dominated by fine-grained black shales rich in various sulfides. Southern section is more dominated by conglomerate. Such horizontal change and mineral characteristic suggest that northern section deposited in deep basin where submarine hydrothermal activities were taking place, and southern section deposited shallow and high energy sedimentation environments. Most of these sedimentary rocks are rich in Cr (up to 900 ppm in black shale). Chromite is often found under microscope. Such finding suggests significantly high flux of clastic materials from ultramafic rocks in ca. 3.4 Ga Onverwacht Group to sedimentary basin of early Fig Tree Group. Some chromite crystals in examined samples have distinct chemistry (very low Mg#) compared to detrital chromite (high Mg #). This suggests that a part of chromite were chemically precipitated from 3.2 Ga ocean water. This further implies the presence of oxidized Cr species (e.g., Cr⁶⁺) in 3.2 Ga shallow ocean water. Those clastic sedimentary rocks are overlain by banded iron formations. In particular, silica-rich band in banded iron formation in northern section contain appreciable amounts of organic carbon (up to 0.2 wt %). Surprisingly carbon isotope compositions are constant (at around -25 per mil) from early clastic sedimentary rocks to banded iron formations, although their sedimentary environments and ages were different. Such homogeneous carbon isotope compositions require single and high productive primary producer, which maybe lived in photic zone, through sedimentation of Fig Tree Group. Presence of oxygenic photosynthesis in 3.2 Ga oceans well explains oxidation of Cr (and also Fe²⁺) and high primary production.

キーワード：シアノバクテリア、南アフリカ、フィグツリー

Keywords: cyanobacteria, South Africa, Fig Tree

西グリーンランドの太古代かんらん岩体から何がわかるか？ What can we learn from ultramafic rocks in Mesoarchaeon orthogneisses in the Fiskefjord region of southern West Greenland?

*森下 知晃¹、Szilas Kristoffer^{2,3}、Pearson Graham⁴

*Tomoaki Morishita¹, Kristoffer Szilas^{2,3}, Graham Pearson⁴

1. 金沢大学理工研究域自然システム学系、2. スタンフォード大学、3. 南デンマーク大学、4. アルバータ大学

1. School of Natural System, College of Science and Technology, Kanazawa University, 2. Stanford University, 3. University of Southern Denmark, 4. University of Alberta

Ultramafic rocks occur sporadically as variably sized, lens-shaped bodies (e.g., Seqi, Miaggoq and Ulamertoq) within Mesoarchaeon (ca. 3000 Ma) amphibolite- to granulite-facies orthogneisses in the Akia terrane of southern West Greenland (Garde, 1997 *Geol. Greenland Surv. Bull.*; Szilas et al., 2015a *GeoResJ*). Two contrasting models can be proposed for the origin of peridotite bodies in the studied area and other Archaean belts in Greenland: (1) residual mantle peridotite after partial melting, and (2) ultramafic cumulates (Friend & Nutman, 2011 *Geology*; Szilas et al., 2015a, 2015b *Gondwana Res.*). In either case, these ultramafic bodies would provide new insights into the links between magmatic-metasomatic processes and crust-continent formation during the Archaean. The Ulamertoq body is a relatively large ultramafic body, 1 x 1.5 km in size. Here we document its field occurrence and petrological characteristics as an example of the ultramafic bodies in the studied area in the context of magmatic and metasomatic processes during the Archaean age. The Ulamertoq peridotite body and also other ultramafic bodies are crosscut by granitic rocks at ca. 3000 Ma (U-Pb zircon; Szilas et al., 2016 AGU abstract). The Ulamertoq peridotite body suffered extensive metasomatism of hydrous silica-rich metasomatic fluids/melts and provides evidence for subduction-like metasomatism during the Archaean. Consequently, although it is difficult to find primary igneous features in peridotite bodies from Archaean belts, our field observations and geochemical data from these peridotite bodies support their cumulate origin (Szilas et al., 2015a, b).

キーワード：30億年かんらん岩体の産状、30億年前の島弧交代作用、30億年前のかんらん岩集積岩

Keywords: Occurrence of Mesoarchaeon peridotite body, Mesoarchaeon Subduction-related metasomatism, Archaean ultramafic cumulate

The Hadean environment inferred from mineral inclusions within the oldest terrestrial zircons

The Hadean environment inferred from mineral inclusions within the oldest terrestrial zircons

*山本 伸次¹、田中 大貴²、坂田 周平³、大林 秀行⁴、平田 岳史⁴、磯崎 行雄⁵、小宮 剛⁵

*Shinji Yamamoto¹, Daiki Tanaka², Shuhei Sakata³, Hideyuki Obayashi⁴, Takafumi Hirata⁴, Yukio Isozaki⁵, Tsuyoshi Komiya⁵

1. 横浜国立大学大学院環境情報研究院、2. 横浜国立大学大学院工学部、3. 学習院大学理学部、4. 東京大学理学部、5. 東京大学総合文化研究科

1. Graduate School of Environment and Information Sciences, Yokohama National University, 2. College of Engineering Science, Yokohama National University, 3. Department of Chemistry, Gakushuin University, 4. Graduate School of Science, The University of Tokyo, 5. Graduate School of Arts and Sciences, The University of Tokyo

Terrestrial rock records during Hadean era (> 4.0 Ga) are scarcely preserved due to surface erosion, subduction, and intense meteorite bombardment. Clues about conditions during this time can be deduced from detrital zircon and its mineral inclusions as old as 4.4 Ga recovered from metasedimentary rocks at Jack Hills in the Narryer Gneiss Complex, Western Australia [1-2]. Hadean zircons from Jack Hills contain various mineral inclusions such as muscovite, quartz, plagioclase, biotite, hornblende, apatite, magnetite, and biogenic carbon (graphite). Investigations of these mineral inclusions have suggested the emergence of a hydrosphere, granitic continental crust, sedimentary cycling, geo-magnetic field, and biosphere in Hadean earth [3-6]. Especially, chemistry of apatite inclusions in zircon reflects the compositions of whole rocks, thus can characterize the host magma [7]. However, the percentage of Hadean zircons to detrital zircons in Jack Hills was as small as 5% [8]. Moreover, skepticism about the preservation of such inclusions have recently raised because of secondary metamorphic overprint [9]. Owing to these reasons, large number of age analysis for zircon grains and precise identification of primary mineral inclusions must be operated to derive reliable information for Hadean earth environment. To overcome this, we have developed rapid and precise dating technique using laser ablation ICP-mass spectrometer (LA-ICP-MS). Mineral inclusions in the zircons were investigated using laser Raman microspectroscopy, equipped with high-speed Raman mapping system. Our preliminary results show that primary apatite and melt inclusions are successfully recovered from zircons of over 4.0 Ga out of 3000 checked grains. The chemistry of apatite and melt inclusions are commonly utilized to characterize host rock, missing Hadean crust. Combination of rapid U-Pb dating with LA-ICP-MS and mineral identification with laser Raman leads us to trace a missing Hadean environment from the oldest zircon of the Earth.

References

- [1] Compston, W. and Pidgeon, R.T., 1986, *Nature*, 321, 766-769. [2] Wild, S.A., Valley, J.W., Peck, W.H., and Graham, C.M., 2001, *Nature*, 176-178. [3] Hopkins, M., Harrison, T.M., and Manning, C.E., 2009, *Nature*, 456, 493-49. [4] Harrison, T.M., 2009, *Annu. Rev. Earth Planet. Sci.*, 37, 479-505. [5] Bell, E.A., Boehnke, P., Harrison, T.M., and Mao W.L., 2015, *PNAS*, 112, 144158-14521. [6] Tarduno, J.A., Cottrell, R.D., Davis, W.J., Nimmo, F., and Bono, R.K., 2015, *Science*, 349, 521-524. [7] Jennings E.S., Marshcall, H.R., Hawkesworth, C.J., and Storey, C.D., 2011, *Geology*, 39, 863-866. [8] Holden, P., Lanc, P., Ireland, T.R., Harrison, T.M., Foster, J.J., and Bruce, Z., 2009, *Int. Jour. Mass Spectrometry*, 286, 53-63. [9] Rasmussen B., Fletcher, I.R., Muhling, J.R., Gregory, C.J., and Wild, S.A., 2011, *Geology*, 39, 1143-1146.

キーワード : Hadean, zircon, mineral inclusion
Keywords: Hadean, zircon, mineral inclusion

Geology of the Eoarchean Nuvvuagittuq supracrustal belt: Constraints on the geochemical signatures of ^{142}Nd anomaly and a geochemical variation of volcanic rocks.

*小宮 剛¹

*Tsuyoshi Komiya¹

1. 東京大学大学院総合文化研究科広域科学専攻

1. Department of Earth Science & Astronomy Graduate School of Arts and Sciences The University of Tokyo

The Nuvvuagittuq supracrustal belt is one of the oldest supracrustal belts so that the geology, geochemistry and geobiology provide important insights into the early evolution of the earth. The minimum age (ca. 3.8 Ga) of the supracrustal rocks are provided by U-Pb dating of zircons from surrounding granitic gneisses and thin felsic intrusions into the supracrustal belt. Besides, the mafic and ultramafic rocks were suggested to be formed in the Hadean as old as 4.2 Ga based on a pseudo-isochron of $^{147}\text{Sm}/^{144}\text{Nd}$ - $^{142}\text{Nd}/^{144}\text{Nd}$. However, the interpretation of the pseudo-isochron of $^{147}\text{Sm}/^{144}\text{Nd}$ - $^{142}\text{Nd}/^{144}\text{Nd}$ is highly controversial; the apparent good correlation between them is also regarded as mixing between chondritic $^{142}\text{Nd}/^{144}\text{Nd}$ and low $^{142}\text{Nd}/^{144}\text{Nd}$ sources. The geological interpretation of the felsic intrusions is also on debate, and felsic volcanoclastic sediments intercalated with the supracrustal rocks are also proposed for the protoliths. In this case, the age of zircons from the rocks is interpreted as a depositional age of the sediments, namely the age of the supracrustal belt.

The Nuvvuagittuq supracrustal belt (NSB) is located on the eastern coast of Hudson Bay, in the Inukjuak Block of the Hudson Bay Terrane of the Northeastern Superior Province of Canada, and contains ultramafic rocks, mafic rocks, banded iron formation (BIF), chert, conglomerate, gabbroic and felsic intrusions. The belt forms an arcuate structure with moderately dipping and NW-plunging hinge line. In addition, the belt was also tightly to isoclinally folded before the folding to form a syncline structure. Previous work suggested three groups of mafic-ultramafic suites with different chemical compositions such as TiO_2 contents and REE patterns: High-Ti, depleted Low-Ti and enriched Low-Ti Ujaraaluk units in ascending order, respectively, and considered the ultramafic rocks as sills.

We made a detailed geological map of the southwestern part of the NSB at a 1:5000 scale, and found many geologic differences from the previous works. The belt comprises ultramafic volcanic rocks, mafic volcanic rocks, mafic intrusions, BIF, chert, "carbonate rocks", "putative" conglomerates and fine-grained felsic intrusions, and pegmatitic intrusions.

The previous work suggested that the southwestern part forms the synclinal structure, but our geologic reappraisal shows that the volcanosedimentary sequences of ultramafic and mafic volcanic units overlain by the BIF and chert mostly strike NS and dip to the east. The stratigraphy that ultramafic rocks are overlain by the BIF layers indicates that they are not sills but flows under the water. Although there are some coarse-grained amphibolites, possibly originating from gabbroic intrusions, many NS-trending structures are not due to the gabbroic intrusions but due to the felsic intrusions. Many of the gabbroic intrusions, interpreted by previous works, are not intrusions, but they originate from mafic volcanic rocks and apparently form linear structures due to felsic intrusions on both sides. It is because there are no chilled margins along the rims and the grain sizes are varied within the linear structures. Although a banding structure occurs along the boundary with mafic rocks in some places, a large chert layer (silica formation) should be referred as a pegmatitic intrusion, which is one of many large pegmatite intrusions in the belt. Two types of felsic intrusions are ubiquitously distributed over the belt: pegmatite and fine-grained felsic intrusions. The felsic intrusions are quite enigmatic; the felsic intrusions fade away into

conglomerate-like structures in some places and form many linear structures within the mafic rock units. We, preliminarily, consider that the felsic intrusions account for the conglomerates and three geochemical groups of basaltic volcanics due to assimilation of the felsic rocks into the mafic rocks as well as a pseudo-isochron of $^{147}\text{Sm}/^{144}\text{Nd}$ - $^{142}\text{Nd}/^{144}\text{Nd}$.

キーワード : 142Nd同位体異常、初期分化、初期太古代

Keywords: 142Nd isotope anomaly, Early differentiation, Eoarchean

ジルコンから読み解く大陸地殻進化

Zircon perspectives on the evolution of the continental crust

*飯塚 毅¹、山口 能央¹、板野 敬太¹、日比谷 由紀¹、鈴木 和恵¹

*Tsuyoshi Iizuka¹, Takao Yamaguchi¹, Keita Itano¹, Yuki Hibiya¹, Kazue Suzuki¹

1. 東京大学

1. University of Tokyo

One of the Earth's unique features among the known terrestrial planets is the presence of a chemically evolved massive crust—the continental crust. It is generally accepted that the modern continental crust has an andesitic bulk composition and is vertically stratified in terms of lithological and chemical compositions—from lower portions dominated by mafic rocks to upper portions dominated by granitoids. Yet there is considerable debate as to when and how the continental crust has evolved to its present form. In this presentation, we explore perspectives on the evolution of the continental crust emerged from the zircon U–Pb age and Hf isotope data with supplementary zircon O isotope data.

Zircon, a common accessory phase in granitoids, can be precisely dated by the U–Pb system. Zircon Hf isotopic composition is a function of crustal residence time of the magmatic protolith, whereas the O isotopic composition is a sensitive record of reworking of mature sediments such as pelite. An integration of U–Pb, Lu–Hf and O isotopic data for detrital zircons from modern large rivers indicates that: (1) granitoid magmatism has played a significant role in the crustal differentiation and crustal reworking over geologic history, (2) the supercontinent cycle has controlled the evolution of the continental crust by regulating the rates of crustal generation and intra-crustal reworking processes and the preservation potential of granitoid crust, and (3) ~25% and ~80% of the preserved continental crust would be formed by 2.5 Ga and 1.0 Ga, respectively. Given that the continental growth is a combined result of the addition of new continental crust and recycling of continental crust into the mantle, the actual net growth of continental crust could be remarkably earlier than this. Taking into account the crustal residence times of continental crust recycled back into the mantle, we propose a continental growth model that stable continental crust was firstly established in the Paleo- and Mesoproterozoic and significantly grew in the Paleoproterozoic.

キーワード：大陸地殻、ジルコン、同位体

Keywords: continental crust, zircon, isotope

U-Pb zircon geochronology of the North Pole Dome adamellite in the eastern Pilbara Craton

*浅沼 尚¹、大林 秀行²、澤木 佑介¹、坂田 周平³、鈴木 和恵⁴、北島 宏輝⁵、平田 岳史⁴、丸山 茂徳¹
 *Hisashi Asanuma¹, Hideyuki Obayashi², Yusuke Sawaki¹, Shuhei Sakata³, Kazue Suzuki⁴, Kouki Kitajima⁵, Takafumi Hirata⁴, Shigenori Maruyama¹

1. 東京工業大学、2. 京都大学、3. 学習院大学、4. 東京大学、5. ウィスコンシン大学

1. Tokyo Institute of Technology, 2. Kyoto University, 3. Gakushuin University, 4. The University of Tokyo, 5. University of Wisconsin

Supracrustal rocks around the North Pole Dome area, Western Australia, provide valuable information regarding early records of crustal growth, surface environments and biosphere. Owing to the occurrence of the oldest known microfossils, the North Pole Dome area has attracted interests from many researchers. The Paleoproterozoic successions (Warrawoona Group) in this area mainly comprise mafic-ultramafic greenstones with intercalated cherts and felsic lavas. Previous age constraints on the succession have been mainly based on zircon U-Pb geochronology of felsic rocks (e.g., adamellite, rhyolite and tuff). However, most zircon grains have suffered from metamictization and contain anomalously high contents of common Pb, which makes interpretation of the U-Pb data complicated. Actually, previous zircon U-Pb data were highly scattered even plotted on Concordia curve within their analytical errors. In order to provide more convincing chronological constraints, we attempted to acquire U-Pb Concordia ages, which had been widely accepted as the best estimate of U-Pb age with the smallest uncertainties.

The North Pole Dome adamellite intrudes into the bottom of the Warrawoona Group, and most zircons separated from the adamellites (95NP207 and 96NP208) suffered from severe metamictization. We selected less metamictized domains using a pre-ablation technique in conjunction with elemental mapping and then conducted *in situ* U-Pb isotopic analyses with a laser ablation inductively coupled plasma mass spectrometry. Most analyzed domains contain certain amounts of common Pb ($^{204}\text{Pb}/^{206}\text{Pb} > 0.0001$), whereas we have obtained three and five concordant data points with less common Pb ($^{204}\text{Pb}/^{206}\text{Pb} < 0.0001$). These concordant data-sets yield Concordia ages of 3445 ± 23 and 3454 ± 17 Ma, respectively. Although these ages are almost similar to those from previous reports, the new ages are based on more reliable data with the smaller errors owing to avoidance of metamictized domains. These ages constrain the intrusive age of the North Pole Dome adamellite and the minimum depositional age of the Warrawoona Group and indicates that the microfossils discovered in this area had appeared by ca. 3454 Ma. In addition, a single xenocrystic zircon grain with less common Pb ($^{204}\text{Pb}/^{206}\text{Pb} < 0.0001$) shows a $^{207}\text{Pb}/^{206}\text{Pb}$ age of ca. 3545 Ma, supporting the idea that the sialic basement of the Pilbara Craton existed prior to 3500 Ma.

In general, old zircon grains often suffer from metamictization and contain detectable level of common Pb, which are unsuitable to constrain precise igneous ages of the host rocks. In contrast, the *in situ* U-Pb zircon dating combined with the pre-ablation technique can obviate to measure such metamictized domains, and has the potential for yielding precise and accurate geochronological data even from metamict zircons.

キーワード：ノースポール地域、ジルコンU-Pb年代学、メタミクト化、初期鉛、U-Pbコンコーディア年代

Keywords: North Pole Dome, U-Pb zircon geochronology, Metamict, Common Pb, U-Pb Concordia age

太古代大気組成解明に向けた高波長分解能4種硫黄同位体SO₂吸収断面積測定

High-resolution ^{32/33/34/36}S SO₂ absorption cross-section measurements for revealing Archean atmospheric composition

*遠藤 美朗¹、小川 萌子²、新開 真純²、セバスチアン ダニエラチェ^{2,3}、上野 雄一郎^{1,3,4}

*Yoshiaki Endo¹, Moeko Ogawa², Masumi Shinkai², Sebastian Danielache^{2,3}, Yuichiro Ueno^{1,3,4}

1. 東京工業大学大学院、2. 上智大学、3. 東京工業大学地球生命研究所、4. 海洋研究開発機構

1. Tokyo Institute of Technology, 2. Sophia Univ., 3. ELSI, 4. JAMSTEC

Many geological and geochemical records suggest a reducing Archean atmosphere. Revealing Archean atmospheric composition is important proxy for the understanding of the origin and evolution of life. The discovery and explanation of sulfur mass-independent fractionation (S-MIF) signatures in Archean sedimentary rocks possess as a key to unravel Archean atmospheric composition. Our SO₂ photochemical experiments generated large S-MIF ($\Delta^{33}\text{S} > +5\text{‰}$) and reproduced basic character of the Archean S-MIF signature ($\Delta^{36}\text{S}/\Delta^{33}\text{S} = -1$) under a specific condition for the first time (Endo et al. 2016). Self-shielding of SO₂ photodissociation and intersystem crossing (ISC) from singlet SO₂ to triplet SO₂ are shown as key mechanisms. Next, we simulated large S-MIF signature ($\Delta^{33}\text{S} > +5\text{‰}$) in our box numerical model, and we showed that Archean S-MIF trend ($\Delta^{36}\text{S}/\Delta^{33}\text{S} = -1$) can be explained when there are several ppm level of SO₂ (like the plume of volcanic gas) and 2% (2 kPa) CO or 3% (3 kPa) CH₄ in the atmosphere. But box model's calculation is not completely correct because photochemical reaction rate and fractionation factor (such as self-shielding) strongly depends on the altitude. Then we need to develop 1-D atmospheric model.

Our group's new 1-D model which focuses on UV spectra because reactions and fractionation factor (such as self-shielding) change delicately as a function of irradiative photon flux. Fractionation factor of photodissociation can be calculated by isotopologue cross-section and irradiative photon flux. Then in order to develop 1-D atmospheric model including sulfur isotopes, SO₂ isotopologue cross-sections (^{32/33/34/36}SO₂) are necessary. Although SO₂ isotopologue cross-section have been measured, they are too low-resolution to estimate self-shielding (Danielache et al. 2008, Endo et al. 2015). Here, we report preliminary results of high resolution ($\sim 1\text{cm}^{-1}$) ^{32/33/34/36}SO₂ absorption cross-sections and estimation of fractionation factor including self-shielding effect.

Referces: Endo et al. (2016), EPSL, Danielache et al. (2008), JGR Atmospheres, Endo et al. (2015), JGR Atmospheres

キーワード：太古代大気、硫黄同位体、光化学

Keywords: Archean atmosphere, Sulfur isotope, Photochemistry

西オーストラリア・ピルバラ地域デキソンアイランド層における32億年前の重晶石および黄鉄鉱硫黄同位体不均質

3.2 Ga sulfur isotopic heterogeneity of barite and pyrite microcrystals in Dixon Island Formation, Pilbara, Western Australia.

*三木 翼¹、清川 昌一¹、高畑 直人²、石田 章純³、伊藤 孝⁴、池原 実⁵、佐野 有司²

*Tsubasa Miki¹, Shoichi Kiyokawa¹, Naoto Takahata², Akizumi Ishida³, Takashi Ito⁴, Minoru Ikehara⁵, Yuji Sano²

1. 九州大学大学院理学府地球惑星科学専攻、2. 東京大学大気海洋研究所、3. 東北大学大学院理学研究科、4. 茨城大学教育学部、5. 高知大学海洋コア総合研究センター

1. Department of Earth and Planetary Sciences, Graduate School of Science, Kyushu University, 2. Atmosphere and Ocean Research Institute, the University of Tokyo, 3. Department of Geoscience, Graduate school of Science, Tohoku University, 4. Faculty of Education, Ibaraki University, 5. Center for Advanced Marine Core Research, Kochi University

中期太古代(32~28億年前)は還元的大気海洋環境の遷移的酸化を示す大規模縞状鉄鉱層の形成(Windley, 1995)など多くの地球史上重要なイベントがあった時代として知られている。当時の硫黄サイクルや酸化還元環境を知る上で硫黄の安定同位体比($\delta^{34}\text{S}(\text{‰}) = ((^{34}\text{S}/^{32}\text{S})_{\text{sample}} / (^{34}\text{S}/^{32}\text{S})_{\text{standard}} - 1) \times 1000$)は重要なプロキシである。各硫黄化学種の $\delta^{34}\text{S}$ はマントルや海洋などのリザーバーごとに大局的な平衡状態にある。しかし局所的にそれらが混合することや酸化還元反応により化学種が変わることで値は大きく変化し、とくに還元性物質である硫化鉱物および酸化性物質である硫酸塩鉱物の $\delta^{34}\text{S}$ 変動は酸素濃度上昇や硫酸還元菌の活動活発化と密接に関わっていると考えられる。近年の微小領域高精度の分析法の向上により詳細な変化を識別できるようになったことから、本研究では32億年前の堆積物に注目し、西オーストラリア州ピルバラ地域のデキソンアイランド層中にわずかに残存する重晶石および黄鉄鉱について高空間分解能二次イオン質量分析計(NanoSIMS)による $\delta^{34}\text{S}$ 微小領域分析を行った。

デキソンアイランド層は下位からコマチアイト-流紋岩質凝灰岩部層、黒色チャート部層、多色チャート部層からなる。多色チャート部層中の黄鉄鉱試料について、DXCL掘削により得られた未風化黄鉄鉱試料は数mmから1mm以下の非常に細かなラミナを成し、 $\delta^{34}\text{S}$ 値は $-10.1 \sim +26.8\text{‰}$ ($n=143$; Avg. = $+8.9\text{‰}$)(坂本, 2010MS)、初期にできた微小球殻状黄鉄鉱内でも $10\mu\text{m}$ で $5\sim 10\text{‰}$ の同位体変化が見られる(三木, 2015MS)。

また、重晶石層はデキソンアイランド層黒色チャート部層の下部に5~8層残っているが全体に珪化しており、珪化を免れた重晶石が $200\mu\text{m}$ 以下の結晶としてごくわずかに保存されている。また重晶石が結晶形を保って珪化した偽晶は非常に細粒の黄鉄鉱を含む。この地層から3層準を選んで岩石試料を採取し、粒界粉碎により重晶石29粒および付随する黄鉄鉱19粒を取り出してNanoSIMS分析を行った。その結果、重晶石からは $-7.1 \pm 1.0 \sim +18.7 \pm 0.9\text{‰}$ (Avg. = $+0.4 \pm 1.3\text{‰}$) という値が得られた。一方、共出した黄鉄鉱は $+2.1 \pm 2.0 \sim +22.3 \pm 5.9$ (Avg. = $+11.4 \pm 2.8\text{‰}$)であった。

堆積物中で硫酸還元菌が関与した場合、最終生成物である黄鉄鉱の $\delta^{34}\text{S}$ は負の値を取ることで知られているが、本研究ではほとんど差がなく、むしろ黄鉄鉱が重い値を持った。最近の研究では32億年前の硫黄同位体の値は高い値をとっており、硫酸還元菌が関与しない反応経路もしくは火山活動などの外からの流入が考えられる。またレイリー分別により正にシフトすることはラグーンや全球凍結等硫酸供給が閉じた環境を意味するため、 $\delta^{34}\text{S}$ 値が高くなる原因を識別することが重要になる。

キーワード：太古代、硫黄同位体、黄鉄鉱、重晶石

Keywords: Archean, sulfur isotope, pyrite, barite

古原生代Transvaal超層群Hotazel層の縞状鉄鉱層およびMn堆積物の地質学・地球化学的研究：希土類元素組成から示唆される堆積環境の酸化還元と熱水の影響の変化

The petrographical and geochemical characteristics of the BIFs in the Hotazel Formation from Transvaal Supergroup: Implications for redox and hydrothermal contributions in their depositional environments.

*青木 翔吾¹、坂田 周平²、中田 亮一³、柏原 輝彦³、大野 剛²、高橋 嘉夫⁴、Tsikos Harilaos⁵、小宮 剛¹

*Shogo Aoki¹, Shuhei Sakata², Ryoichi Nakada³, Teruhiko Kashiwabara³, Takeshi Ohno², Yoshio Takahashi⁴, Harilaos Tsikos⁵, Tsuyoshi Komiya¹

1. 東京大学大学院総合文化研究科、2. 学習院大学理学部化学科、3. 海洋研究開発機構、4. 東京大学大学院理学研究科、5. Department of Geology, Rhodes University

1. Graduate School of arts and science, the University of Tokyo, 2. Department of Chemistry, Gakushuin University, 3. Japan Agency for Marine-Earth Science and Technology, 4. Graduate School of Science, the University of Tokyo, 5. Department of Geology, Rhodes University

The surface environments on the earth have evolved from anoxic to oxic. The oxidation has occurred discontinuously mainly at two times throughout the earth history: the Neoproterozoic to the Paleoproterozoic and the Neoproterozoic. The first oxidation event is known as “the Great Oxidation Event (GOE)”. In the Paleoproterozoic Hotazel Formation of the Transvaal Supergroup, South Africa, there are the manganese ores hosted by the banded iron formations at the three stratigraphic horizons, which is one of the most conspicuous evidences for GOE.

Previous studies in the Hotazel Formation have focused only on their metallogenic studies of the Mn ores such as post-depositional alterations (e.g. Gutzmer and Beukes, 1997). On the other hand, their sedimentary environment changes responsible for depositions of the manganese rocks and the BIFs have not been constrained fully. In this study, we tried to estimate redox and hydrothermal contributions in the sedimentary environments from the lowermost BIFs to the lowest Mn-rock layer based on stratigraphic variations of some geochemical proxies (some major element contents such as Mn, Ca and Fe, and REE + Y patterns).

In the studied strata, the Mn/Fe ratios and Ca/Fe ratios from the lowermost BIFs to the lowest Mn rocks show increasing trends, suggesting that precipitations of Mn-oxide minerals and Ca-carbonate minerals were becoming prevailed relative to that of Fe-oxide minerals in the sedimentary environments. These trends suggest that the sedimentary environments were becoming more oxic and shallower.

Whole-rock REE + Y contents in the BIFs show weakly positive correlations with Fe contents and strongly positive correlations with phosphorus (P) contents. Moreover, micro-scale elemental imaging in the lowermost BIFs shows that apatite occur as spots in the Fe-oxide bands, and REE + Y is concentrated in those spots. Those REE + Y distributions in the BIFs suggest that REE + Y might have been primarily derived from adsorbents on Fe oxyhydroxide and secondarily moved into phosphorous minerals at the diagenesis. On the other hand, REE + Y in the lowest Mn-rock layer is positively correlated with Mn contents, suggesting that those elements might have been derived from adsorbents on Mn oxyhydroxide. However, secondarily movement of REE + Y associated with the diagenesis forming apatite might be less influential on REE + Y patterns because anomalous behaviors of Eu and Ce are seen regardless of P contents. PAAS-normalized REE + Y patterns show that the lowermost BIFs overlying the Ongeluk

Formation show positive Eu anomalies characteristics of high-temperature hydrothermal fluids (e.g. Bau and Dulski, 1999). On the other hand, the Mn rocks show negative Ce anomalies similar to modern oxic seawater (e.g. Alibo and Nozaki, 1999).

Above stratigraphic variations of Mn/Fe, Ca/Fe ratios and REE + Y patterns in the analyzed strata suggest that the Paleoproterozoic ocean was composed of double-layered structure. The deep ocean was anoxic and subject to contributions of hydrothermal fluids, resulting in precipitations of Fe-oxide minerals (the BIFs). On the other hand, the shallow ocean was oxic with active primary productions, resulting in Mn-oxide and Ca-carbonate minerals.

キーワード：縞状鉄鉱層、Mn堆積物、希土類元素組成

Keywords: Banded Iron Formations, Manganese rocks, Rare Earth Elements

カナダ/古原生代トランスハドソン変動帯に残された堆積盆の層序復元: Reconstruction Paleoproterozoic sedimentary basin stratigraphy in Trans Hadoson Orogeny, Canada

*清川 昌一¹、元村 健人¹、Bleeker Wouter²、Price Dave³

*Shoichi Kiyokawa¹, Kento Motomura¹, Wouter Bleeker², Dave Price³

1. 九州大学大学院理学研究院地球惑星部門、2. カナダ地質調査所、3. Hadbay Mine

1. Department of Earth and Planetary Sciences Faculty of sciences, Kyuushu University, 2. Geological Survey of Canada, 3. Hadbay Mine

古原生代は、大気中の酸素濃度上昇事件 (Grate Oxidation Event) や、真核生物の台頭が始まり、大陸集合・分裂イベントを伴って地球表層が大きく変化したと言われる (e.g. Holland, 1994, Condie 1997, Kopp et al, 2005). 表層の急激な酸化作用から、海底は極度の還元環境に陥り、硫化物に富むユーキシニック海洋が広がるとされている (Lyons et al., 2014). しかし、大陸が安定化するこの時代の深海底の地質学的証拠は原生代の深海底の地層は付加や衝突帯に巻き込まれており、その復元は詳細な地質調査による層序復元が重要になる。我々は古原生代における比較的深い海底の環境変遷を明らかにするために、カナダ・スペリオルクラトンの北縁にあたるトランスハドソン造山帯中に残されるフリントロン帯およびケープスミス帯に注目した。特に黒色頁岩を含む地層に注目し、地質調査および鉱山が取得する掘削コア記載・サンプリングにより、比較的深い海洋環境における海底状態の復元を試みている。

フリントロン帯 (Trout Lake (Embury lake) Formation トロウトレイク層) : 島弧火山活動によって形成されている火山岩帯に沿って分布する砂岩・泥岩からなるタービダイト層からなる。露頭では造山運動の影響により千枚岩化しており、タイト褶曲が面構造に沿って見られる。Hadbay社の掘削コアより、400mの連続試料について現地にて記載し、黒色頁岩部分を中心に200個におよび試料を取得している。褶曲解析により約150mの連続層序が復元できた。薄片観察にて、これらの地層は石英粒子に乏しい火山岩起源の砂岩と泥岩の互層であり、島弧起源から供給された比較的深い堆積盆でできた地層と思われる。現在、砂岩および石英斑岩の年代測定、炭素同位体測定を行っている。

ケープスミス帯 (Nuvilik Formation, Povungnituk Group : ヌービリック層, ブブンマタク層群)

ケープスミス帯は19億年前のシート岩脈をもつオフィオライトからなるWatts層群、構造的下位の玄武岩およびコマチアイト溶岩を主体とするChukotat層群、Watts層群の下位に整合的關係で分布する大陸棚起源堆積岩 (黒色頁岩) のPovungituk層群からなる。Chukotat層群の最下部に位置するコマチアイト溶岩はサーマルエロージョンによる侵食w起こしており、溶岩最下部はNi-Cu硫化物鉱物鉱床 (Relgan鉱山) を形成する (Leshner 2017)。我々はコマチアイト溶岩直下に分布する黒色頁岩 (千枚岩) からなるNuvilik層

(Povungituk層群) について、Relgan鉱山の許可を得て露頭調査を行なった。また、2箇所のコアKikialik siteにおいて鉱山内から掘削した300mコア (468069 core) と Katinniq siteのzone 5-8 地域での表層から800mパイロットコア(718-3485 core)について、詳細な層序記載および試料採取を行った。薄片観察では、黒色頁岩には石英粒子をおよび有機物に富む特徴を持つ。スペリオルクラトンの北縁部にあたる2地域での比較的深い海底堆積物について、層序復元を陸上調査および掘削コア記載を行った。

キーワード : 古原生代、トランスハドソン造山運動、有機物深海堆積物

Keywords: Paleoproterozoic, Trans Hudson Orogeny, rganic rich deep sea sediment

Lithology and Depositional Age of Paleoproterozoic Volcaniclastic Sequence from Ashanti Belt of Birimian Supergroup, Southwest Ghana

*吉丸 慧¹、清川 昌一¹、坂井 志緒乃¹、伊藤 孝²、池原 実³、堀江 憲路⁴、竹原 真美⁴、佐野 貴司⁵、Nyame Frank⁶、Tetteh Georg⁷

*Satoshi Yoshimaru¹, Shoichi Kiyokawa¹, Shiono Sakai¹, Takashi Ito², Minoru Ikehara³, Kenji Horie⁴, Mami Takehara⁴, Takashi Sano⁵, Frank K Nyame⁶, George M Tetteh⁷

1. 九州大学理学府地球惑星科学専攻、2. 茨城大学、3. 高知大学、4. 国立極地研究所、5. 国立科学博物館、6. ガーナ大学、7. タルクワ鉱山技術大学

1. Department of Earth and Planetary Science, Kyushu University, 2. Ibaraki University, 3. Kochi University, 4. National Institute of Polar Research, 5. National Museum of Nature and Science, 6. University of Ghana, 7. University of Mines and Technology

Paleoproterozoic Birimian volcaniclastic successions occur along the coast near Cape Three Points, in Ashanti belt, southwest Ghana. In this study, the depositional environments and bioactivities recorded in the c.a. 2.3 Ga rocks were investigated. We report the structure and stratigraphy in the area, improved depositional age of the sediments by zircon U-Pb dating and C isotope features of the bioactivities in the paleo-ocean.

The Ashanti belt, generally showing NE-SW strike, composed mainly of andesitic basalts, volcaniclastic rocks and belt type granitoids (Perrouty et al., 2012). The greenstone is unconformably overlain by Tarkwaian conglomerates and metasedimentary rocks. The maximum depositional age of the overlying metasedimentary rocks and the oldest age of the intruded rock into Birimian volcanics in the Sekondi region is 2154 ± 2 Ma and 2174 ± 2 Ma, respectively (U-Pb zircon: Oberthür et al., 1998).

Detailed field investigations were performed to reconstruct about 1000m out of over 2000m thick stratigraphy of the volcaniclastic sequence in the Cape Three Points area. The rocks, affected by greenschist-facies metamorphism, generally strike N-S, mainly dip $60-80^\circ$ to the east and show fining upward stratigraphy. Using a west vergent thrust fault which occurs in the central part of the area, we sub-divided the area into two, a western zone and an eastern zone. The eastern zone consists of approximately 800 m thick fining upwards sequence which appears to have been repeated two times within the sequence. Thick basaltic lava in the lowermost part of the sequence is replaced by or changes into sediments produced by gravity flows. The uppermost part of the sequence reveals thick basaltic andesite and/or pyroclastic layers which then changes to fine altered volcaniclastics containing organic material. The western zone has not yet been reconstructed the stratigraphy.

$\text{TiO}_2/\text{Al}_2\text{O}_3$ ratios from EPMA analyses of chromite in basaltic rocks suggest that these rocks originated in a volcanic arc system. Whole-rock trace element compositions tend to show low concentrations of Nb and high LREEs, which also supports derivation from volcanic arc.

A foliated porphyritic dyke which occurs in the Cape Three Points area was dated by SHRIMP at the National Institute of Polar Research. Zircon grains in the dyke yielded a weighted mean ^{204}Pb -corrected $^{207}\text{Pb}/^{206}\text{Pb}$ age of 2265.6 ± 4.6 Ma (95% confidence), which suggests that the volcaniclastic sequence was deposited before 2265.6 ± 4.6 Ma and was deformed afterwards. This age is the oldest in the Ghanaian greenstone terrane (Loh and Hirdes, 1999). It would seem, therefore, that rocks in the Cape Three Points area record the history of early volcanic activity in the Birimian greenstone terrain.

We measured organic carbon isotope ratio of some black shale samples. The TOC varies from 0.02 %- 0.3 % and the $\delta^{13}\text{C}$ values are -35‰ - -15‰ (N=5). These low concentrations of the organic matter suggest dilution by the continual influx of volcanogenic clastics.

The thick fining-upward volcaniclastic sequence and chemical compositions may suggest the Cape

Three Points Formation was deposited on mid-deep sea floor beside an oceanic volcanic arc. The deposition may have occurred before 2265.6 ± 4.6 Ma suggesting these sediments record oceanic environmental changes from Makganyene glaciation toward Lomagundi Event (2.3 Ga -2.06 Ga, Bekker and Holland, 2012 and Kopp et al., 2005).

キーワード：ガーナ・ビリミアン超層群、古原生代

Keywords: Birimian Supergroup, Ghana, Paleoproterozoic

東エジプト El Dabbah 地域の層序と堆積場の復元：新原生代の縞状鉄鉱層 Reconstruction of stratigraphy and tectonic setting in El Dabbah, Eastern Egypt: Neoproterozoic banded iron formations

*鈴木 大志¹、清川 昌一¹、池原 実²、佐野 貴司³、ダウッド マヘル⁴、アブエルハッサン モハメッド⁴
*TAISHI SUZUKI¹, Shoichi Kiyokawa¹, Minoru Ikehara², Takashi Sano³, Maher Dawood⁴,
Mohamed Abouelhassan⁴

1. 九州大学、2. 高知大学、3. 国立科学博物館、4. メノフィア大学

1. kyushu Univ., 2. Kochi Univ., 3. National Museum of Nature and Science, 4. Menoufia Univ.

In the Eastern Desert of Egypt, Neoproterozoic Banded Iron Formations (BIFs) are reported within Nubia greenstone belt whose geological structure and stratigraphy are not well understood. In this study, the geological structure was established to reconstruct the stratigraphy including Iron formations (IFs) in El Dabbah in the middle of the Eastern Desert.

Greenstone sequence in El Dabbah area, strike-slip basin on Nubia shield during pan African orogeny, is covered by Hammamat Group. The greenstone sequence preserved the thick volcano-sedimentary rocks with gabbro, coarse-grained volcanoclastics rocks, pillow lavas, black shales, and BIFs. Reconstruction of stratigraphy in this greenstone sequence, which is 4000m in total thickness, composes Basal, Lower, Middle, and Upper members. There are 13 iron sections within this greenstone sequence.

Especially, BIFs sequence preserved within Lower and Upper members. BIFs sequences are well preserved within massive - pillow lavas sections and it contain laminated greenish - black shales. BIFs are composed of magnetite and/or hematite. Magnetite are oxidized from rims and become hematite.

We have determined major and trace element compositions of whole rock of volcanics and BIFs by using XRF and ICP-MS analysis. Trace elements compositions indicate that the volcanics are arc origin in term of a Nb-Zr-Y discrimination diagram and a MORB normalized spider diagram. Major elements in BIFs in this area are characterized by their higher content of Al_2O_3 than the other Archean BIFs. In addition, REE in BIFs show pattern of light REE enrichment and negative Eu anomalies.

This Volcanoclastic sedimentary sequence would be related at island arc sedimentary sequence with volcanic activity.

キーワード：縞状鉄鉱層、新原生代

Keywords: banded iron formation, Neoproterozoic

Toward a better understanding of trace element availability in Paleo-proterozoic seawater

*澤木 佑介¹、佐藤 友彦²、丸山 茂徳²

*Yusuke Sawaki¹, Tomohiko Sato², Shigenori Maruyama²

1. 東京大学、2. 東京工業大学大学院 地球生命研究所

1. The University of Tokyo, 2. Tokyo Institute of Technology

One of the large gaps between prokaryotic and eukaryotic lives lies in the utilization of trace elements in seawater. Eukaryotes have higher Cu and Zn requirements than prokaryotes; for example all eukaryotes have Cu-Zn-SOD (Superoxide Dismutase), whereas prokaryotes have Fe-Mn-SOD. Phylogenomic reconstruction suggested that above-mentioned protein structures evolved either concurrent with or after the emergence of the Eukaryotic domain. Therefore Cu and Zn would be two of the most likely trace metals to provide a geochemical barrier specific to eukaryotic evolution, and deficits of Cu and Zn in seawater have been hypothesized as one of the explanations for the delayed diversification of eukaryotes.

Previous works have tried to decipher a secular variation of marine Zn inventory based mainly on Zn concentrations in ancient sediments. They suggested that the amount of bio-available Zn remained relatively unchanged through time (Robbins et al., 2013; Scott et al., 2013), but stoichiometric assessment concerning removal process of elements during sedimentation still remains difficult. In order to quantitatively track secular variations of Cu and Zn availabilities in seawater, we adopted Cu and Zn isotopes of carbonate rocks and black shales, because these isotopes are expected to reflect marine inventories of each element.

Francevillian Group in Gabonese Republic was recently established as a typical sedimentary sequence for the Paleoproterozoic, and it contains macroscopic structures interpreted as colonial eukaryotic organisms. We hypothesize that these organisms might have evolved in concert with the oxidation of the atmosphere-ocean system and increases of marine Cu and Zn inventories at that time. Although Chi Fru et al. (2016) recently reported eight Cu isotopic ratios from sediments in the Francevillian Group, sequences round macroscopic structures have never been investigated. We collected some carbonate rocks and black shales from the Francevillian Group, and made thin sections from them. SEM-EDS analysis demonstrated that Cu and Zn are enriched in pyrite grains. Powdery samples were prepared by micro-drilling, and acid digestion was performed with aqua regia at 150 Celsius degrees for 48 hours. For isotopic analysis, Cu and Zn were purified using anion exchange on AG-MP-1M resin column. Bulk Cu and Zn isotopic ratio were obtained with a multi collector inductively coupled plasma mass spectrometry (NEPTUNE) at UC Davis. We will report the preliminary results in this presentation.

キーワード：銅同位体比、亜鉛同位体比、ガボン、前期原生代

Keywords: Cu isotope ratio, Zn isotope ratio, Gabon, The Paleo-proterozoic

ガボン・前期原生代フランスヴィル層群における生物多様性 Biodiversity in the Paleoproterozoic Francevillian Group, Gabon

*佐藤 友彦¹、澤木 佑介¹、丸山 茂徳¹

*Tomohiko Sato¹, Yusuke Sawaki¹, Shigenori Maruyama¹

1. 東京工業大学

1. Tokyo Institute of Technology

The Paleoproterozoic (~2.2 Ga) macrofossils are discovered one after another from the sedimentary sequence of the Francevillian Group in Gabon; pyritized macrofossils from Socoba section in Franceville Basin (Albani et al., 2010, *Nature*; 2014, *PLOS ONE*) and nodular fossils from Akou section in Okondja Basin (Moussavou et al., 2015, *J Geol Geosci*; Edou-Minko et al., 2017, *J Geol Geosci*). These fossils are interpreted as microbial colony or colonial eukaryotic organisms on the basis of their complex structures and macro sizes. They are reported from the black shales of the FB Unit; however, the fossil horizons are not correlated precisely because of the poor stratigraphic correlation between the basins. In order to clarify the interbasinal appearance of the fossils and the corresponding environmental changes, we conducted a geological fieldwork and investigated the litho-, bio- and chemo-stratigraphy in the Franceville, Lastoursville, and Okondja Basins. These intracratonic rift basins on the Archean basement rocks were filled up with the Paleoproterozoic Francevillian Group; i.e., FA (fluvial sandstones with uranium ore at the top, including the well-known Oklo nuclear reactors), FB (black shales and carbonates, containing manganese-rich carbonates and the fossil horizon at the upper part), FC (cherts), FD (black shales), and FE (sandstones). We will report the preliminary stratigraphic correlations between the Franceville, Lastoursville, and Okondja Basins, especially on the fossiliferous FB Unit.

キーワード：前期原生代、大型化石、微化石、ガボン

Keywords: Paleoproterozoic, macrofossil, microfossil, Gabon

エディアカラ紀海洋の硫酸濃度上昇イベント

Oceanic sulfate increase events in the Ediacaran ocean

*松浦 史宏¹、斎藤 誠史²、澤木 佑介³、上野 雄一郎^{1,2,4}*Fumihiko Matsuura¹, Masafumi Saitoh², Yusuke Sawaki³, Yuichiro Ueno^{1,2,4}

1. 東京工業大学 地球生命研究所、2. 海洋研究開発機構、3. 東京大学 総合文化研究科、4. 東京工業大学理工学研究科地球惑星科学専攻

1. EARTH-LIFE SCIENCE INSTITUTE, Tokyo Institute of Technology, 2. Japan Agency for Marine-Earth Science & Technology (JAMSTEC), 3. Graduate School of Arts and Sciences, College of Arts and Sciences, The University of Tokyo, 4. Department of Earth & Planetary Sciences, Tokyo Institute of Technology

It is thought that the development of life was closely linked to seawater chemistry, especially redox conditions. Since metazoans diversified in the Ediacaran, revealing redox condition in Ediacaran ocean is crucial for evolution of life. Recent sulfur isotopic values ($\delta^{34}\text{S}$) of carbonate associated sulfate (CAS) and chromium reducible sulfur (CRS) of Ediacaran sediments indicate that the ocean-atmosphere system was progressively oxidized during the Neoproterozoic. Those data further suggest that sulfur cycle was closely related to Shuram excursion, the largest carbon isotope anomaly in the Ediacaran. Ediacaran sediments in South China are ideal to unravel possible linkages between oceanic redox conditions and biological activity, because various proxies have been provided by a number of works. A previous work reported both $\delta^{34}\text{S}_{\text{CAS}}$ and $\delta^{34}\text{S}_{\text{CRS}}$ from Ediacaran Doushantuo Formation at Three Gorges, South China. Their $\delta^{34}\text{S}_{\text{CAS}}$ data were highly scattered, and this is possibly attributed to contamination from CRS during separation processes. Therefore stratigraphic profile of the $\delta^{34}\text{S}_{\text{CAS}}$ has not been fully determined, which leaves ambiguity in reconstruction of the oceanic redox nature.

We carried out sulfur isotope analyses of CAS and CRS of Ediacaran drill core samples collected from the Three Gorges. To reduce contamination from CRS fraction into CAS fraction, we employed an improved method to extract the pure CAS. The newly obtained $\delta^{34}\text{S}_{\text{CAS}}$ values display a smooth curve above the lower part of Member 2 of the Doushantuo Formation and range from +18.7‰ to +46.4‰, except for a datum of -3.7‰. The $\delta^{34}\text{S}_{\text{CRS}}$ values also display a smooth curve, and range from -18.6 to +42.8‰. The differences between $\delta^{34}\text{S}_{\text{CAS}}$ and $\delta^{34}\text{S}_{\text{CRS}}$ values are negatively correlated with $\delta^{34}\text{S}_{\text{CRS}}$ values, suggesting that the $\delta^{34}\text{S}_{\text{CRS}}$ values likely reflect degree of isotopic fractionation during sulfate reduction. The $\delta^{34}\text{S}_{\text{CAS}}$ data in sediments deposited before Shuram excursion are unsynchronized among South China, USA, Mexico, and Oman. The heterogeneous $\delta^{34}\text{S}_{\text{CAS}}$ data possibly arose from low oceanic sulfate concentration, because residence time of oceanic sulfate fell below a mixing time of the various ocean basins when sulfate concentration was low.

The stratigraphic profile of $\delta^{34}\text{S}_{\text{CRS}}$ exhibits two negative excursions, and we firstly found that the both excursions coincide with the positive excursions of the $^{87}\text{Sr}/^{86}\text{Sr}$ ratio and the negative excursions of the $\delta^{13}\text{C}$ value of the carbonate ($\delta^{13}\text{C}_{\text{carb}}$). These correlations indicate that enhanced oxidative weathering increased oceanic sulfate concentration and accelerated release of isotopically light carbon through sulfate reduction of organic matter. Furthermore, the $\delta^{34}\text{S}_{\text{CAS}}$ decreases from +23.5‰ to +18.7‰ parallel with the Shuram excursion. This decreasing trend during the Shuram excursion is comparable to those in Mexico, USA, and Oman. Therefore decrease of $\delta^{34}\text{S}_{\text{CAS}}$ concomitant with the Shuram excursion was a global trend, and this fact further suggests increase of oceanic sulfate concentration during this period. During the Shuram excursion, $\delta^{13}\text{C}_{\text{carb}}$ values started to co-vary with $\delta^{13}\text{C}$ values of the organic carbon, which indicates that a large oceanic organic carbon pool was exhausted owing to elevated oceanic sulfate level and extensive sulfate reduction. Our new results indicate that enhanced oxidative weathering supplied sulfate into ocean and induced high oceanic sulfate level and subsequent Shuram excursion

during the late Ediacaran.

小惑星-彗星衝突による寒冷化-大量絶滅事件の頻度

Frequency of significant cooling and subsequent mass extinctions by asteroid-comet impacts

*海保 邦夫¹

*Kunio Kaiho¹

1. 東北大学大学院理学研究科

1. Graduate School of Science, Tohoku University

For decades, rule of asteroid-comet impacts for climate changes and extinctions of biota has been a subject of debate. A consistent impact rate since the end of the late heavy bombardment and vulnerability of comet suggests approximately 70 and 10 significant Cooling Events enough to cause Mass Extinctions (CEME) after the late heavy bombardment 3.8 Ga and in the Phanerozoic after diversification of multicellular animals, respectively. Many scientists have sought evidence of impact for all mass extinctions. However, only one mass extinction likely occurred by asteroid impact in the Phanerozoic. Here I show that approximately 20 and a few CEME caused by asteroid impacts are possible after 3.8 Ga and in the Phanerozoic, respectively. High concentration of hydrocarbon in target rocks is essential to cause CEME, as an impact burned hydrocarbon in the target rocks forming stratospheric soot, which caused extreme global cooling and draught. The high content areas enough for CEME occupy 13–15% and 40% of the Earth surface for 9–15 km and 20–30 km asteroid impacts, resulting in the decrease of possible number of CEME.

キーワード：大量絶滅、小惑星-彗星衝突、地球寒冷化

Keywords: mass extinctions, asteroid-comet impacts, global cooling

白亜紀セノマニアン／チューロニアン境界のOAE2に対する南半球高緯度の反応

Response to Cretaceous Cenomanian/Turonian OAE2 in southern high latitude

*長谷川 卓¹、牟田 宗一郎²、後藤（桜井） 晶子¹、クランプトン ジェームス³

*Takashi Hasegawa¹, Soichiro Muta², Akiko S. Goto¹, James Crampton³

1. 金沢大学自然システム学系、2. 金沢大学大学院自然科学研究科（現・石油天然ガス金属鉱物資源機構）、3. ニュージーランド地質調査所

1. Division of Global Environmental Science and Engineering, Institute of Natural Science and Technology, Kanazawa University, 2. Graduate School of Natural Science and Technology, Kanazawa University (present affiliation: JOGMEC), 3. GNS Science, New Zealand

白亜紀セノマニアン／チューロニアン(C/T)期境界では、短期間に大量の有機物が広範囲の海底に堆積する「海洋無酸素事件（OAE）2」と呼ばれるイベントが生じたことが知られている。このイベントは顕生代において重要な炭素循環の攪乱イベントと考えられ、テチス域および大西洋域の大陸海を中心に多くの研究がなされてきた。このOAE2期に相当する層位範囲では有機物および炭酸塩の炭素同位体比に特徴的な正のエクスカッション(CIE)がみられることが知られており、炭素同位体比層序による国際対比がなされてきた。しかしながら白亜紀当時の海洋の大半を占めていた太平洋域がOAE2にどのように反応したのかについての知識は乏しい。

本研究では南太平洋高緯度域の海域・陸域にその炭素循環の攪乱の影響がいかに現れるかを理解するため、C/T階境界上下の堆積岩から得られた抽出性有機物に関する総括的な研究を行った。試料を採集した露頭はニュージーランド南島のマルボロ地区である。

バイオマーカー分析から明らかになったホモホパンインデックス(HHI)は世界の他地域のOAE2相当層準付近には知られていない周期的な増減変動を示し、この堆積場の酸化還元環境が周期的に富酸素－貧酸素環境を繰り返していたことを示唆する。OAE2の最初期段階では強い酸素欠乏が生じ、その後急速に酸化的環境に移行した。ステラン／ホパン(S/H)比はこれとほぼ同期して大きく低下したが、HHIの低下（海底の富酸素化）より約10万年程度遅れて生じた。この真核生物由来バイオマーカーの減少は海洋有機物の海底への輸送量の低下、すなわち生物生産の減少を示唆している。海底に到達する有機物が減少することが海底の酸素消費量を減らすことに連動して貧酸素環境から富酸素環境へのシフトを誘導したという考察は、酸化的環境に移行するタイミングが生物生産減少に先行していることを説明できない。酸素に富む水塊の流入が海洋の富酸素化を誘導したという解釈が、上述のOAE2層準のバイオマーカー変動を最も良く説明できるようだ。

OAE2層準における陸域由来の多環芳香族分布は、高緯度南太平洋域において、OAE2直前まで徐々に自然火災を引き起こしやすい気候が卓越していくことを示していた。その後CIEの開始と共にその自然火災頻度は減少し、OAE2期全般に渡ってその頻度は低かったと解釈される。球果植物由来のジテルペノイドと多様な植物に由来するカダレンを用いた高等植物指標はOAE2前後の層準を含めて全体として低下していく傾向を示したが、特にOAE2層準では低い値を示した。このことはOAE2層準では植物群集の中で球果植物が減退していたことを示唆する。

海洋と陸域の環境を示唆するバイオマーカー指標等から、南半球高緯度太平洋域において海洋、陸域の双方で深刻な生息環境の変化が示唆された。この地のOAE2期における環境変化はテチス域およびその他の地域のそれとはどこも似ておらず、OAE2のメカニズムが従来考えられていたよりも複雑であることを示している。

キーワード：白亜紀、セノマニアン、チューロニアン、OAE

Keywords: Cretaceous, Cenomanian, Turonian, OAE

古生代末の二段階の絶滅：四川省・朝天セクションにおける層序学的研究 The double extinction at the end-Paleozoic: Stratigraphy at Chaotian, Sichuan, China

*齋藤 誠史¹、磯崎 行雄²、姚 建新³、紀 戦勝³

*Masafumi Saitoh¹, Yukio Isozaki², Jianxin Yao³, Zhansheng Ji³

1. 独立行政法人海洋研究開発機構、2. 東京大学、3. 中国地質科学院

1. Japan Agency for Marine-Earth Science and Technology, 2. University of Tokyo, 3. Chinese Academy of Geological Science

約2億5000万年前の古生代末では、顕生代最大の大量絶滅事件がおきた。この事件は、ペルム紀中-後期（G-L）境界およびペルム-トリアス紀（P-T）境界における、二段階の絶滅からなる。発表者らはこれまで、南中国・四川省の朝天セクションにおいて、G-L境界層およびP-T境界層の層序学的研究を行ってきた。本発表では、これまでに得られた成果について報告する。

キーワード：大量絶滅、朝天、環境変動

Keywords: mass extinction, Chaotian, environmental changes

The latest research on REY-rich mud in the Pacific Ocean

*加藤 泰浩^{1,2,3}、中村 謙太郎¹、藤永 公一郎^{2,1}、安川 和孝^{1,2}、高谷 雄太郎^{4,1}、大田 隼一郎^{3,1}、田中 えりか¹、見邨 和英¹、飯島 耕一^{3,1}、町田 嗣樹^{3,1}、野崎 達生^{3,1,5}、木村 純一³、岩森 光^{3,6}

*Yasuhiro Kato^{1,2,3}, Kentaro Nakamura¹, Koichiro Fujinaga^{2,1}, Kazutaka Yasukawa^{1,2}, Yutaro Takaya^{4,1}, Junichiro Ohta^{3,1}, Erika Tanaka¹, Kazuhide Mimura¹, Koichi Iijima^{3,1}, Shiki Machida^{3,1}, Tatsuo Nozaki^{3,1,5}, Jun-Ichi Kimura³, Hikaru Iwamori^{3,6}

1. 東京大学・工学系研究科、2. 千葉工業大学・次世代海洋資源研究センター、3. 海洋研究開発機構、4. 早稲田大学・創造理工学部、5. 神戸大学・理学研究科、6. 東京工業大学・理学院

1. School of Engineering, The University of Tokyo, 2. ORCeNG, Chiba Institute of Technology, 3. JAMSTEC, 4. School of Creative Science and Engineering, Waseda University, 5. Graduate School of Science, Kobe University, 6. School of Science, Tokyo Institute of Technology

Deep-sea mud enriched in rare-earth elements and yttrium (REY), termed as REY-rich mud, has been expected as a novel mineral resource for the critical elements. A recent discovery of the extremely REY-rich mud with the maximum total REY content of ~8,000 ppm in the Japanese Exclusive Economic Zone (EEZ) around Minamitorishima Island makes it realistic to economically develop the highly promising deep-sea mineral resource.

In the presentation, we report the progress of our researches on the REY-rich mud within the Minamitorishima EEZ towards the world's first development of deep-sea mineral resources. Moreover, we also introduce our latest findings including the basin-wide spatial distribution of the muds with a high resource potential and various genetic components identified in the western North and central to eastern South Pacific REY-rich muds. Our accumulation and integration of the scientific knowledge about deep-sea sediments including REY-rich mud strongly suggest an intrinsic linkage between the formation of deep-sea mineral resources and Earth system dynamics such as climate change, geochemical cycles, and plate tectonics, which can offer new insights into resource geology, oceanography, paleoclimatology, and solid earth science.

キーワード：海底鉱物資源、レアアース泥、南鳥島、北西太平洋、南太平洋

Keywords: deep-sea mineral resource, REY-rich mud, Minamitorishima Island, western North Pacific, South Pacific

南鳥島南方海域における超高濃度レアアース泥層の側方分布 Lateral distribution of extremely REY-rich mud layer in the southern part of the Minamitorishima EEZ

*藤永 公一郎^{1,2}、安川 和孝^{2,1}、野崎 達生^{3,2,4}、町田 嗣樹^{3,2}、飯島 耕一^{3,2}、舘野 ひとみ²、川崎 健寛²、高橋 亜夕³、天川 裕史³、鳥本 淳司³、深海 雄介³、鈴木 勝彦³、中村 謙太郎²、加藤 泰浩^{2,1,3}、MR15-02 乗船者一同

*Koichiro Fujinaga^{1,2}, Kazutaka Yasukawa^{2,1}, Tatsuo Nozaki^{3,2,4}, Shiki Machida^{3,2}, Koichi Iijima^{3,2}, Hitomi Tateno², Takehiro Kawasaki², Ayu Takahashi³, Hiroshi Amakawa³, Junji Torimoto³, Yusuke Fukami³, Katsuhiko Suzuki³, Kentaro Nakamura², Yasuhiro Kato^{2,1,3}, MR15-02 cruise members

1. 千葉工業大学・次世代海洋資源研究センター、2. 東京大学・工学系研究科、3. 海洋研究開発機構、4. 神戸大学・理学研究科

1. Ocean Resources Research Center for Next Generation, Chiba Institute of Technology, 2. School of Engineering, The University of Tokyo, 3. JAMSTEC, 4. Graduate School of Science, Kobe University

Recently, we discovered deep-sea sediment containing ~7,000 ppm of total rare-earth elements and yttrium (REY), which was termed ‘extremely REY-rich mud’ (Iijima et al., 2016), within the Japanese exclusive economic zone (EEZ) around Minamitorishima island. To clarify the mineralogical/chemical characteristics and distribution of REY-rich mud in the Minamitorishima EEZ, we conducted eight research cruises over the four years. In the MR15-02 cruise by R/V Mirai from June 22 to July 17, 2015, we focused on investigating the lateral continuity of the extremely REY-rich mud layer in the southern part of the Minamitorishima EEZ. During the cruise, we collected 16 sediment cores by piston coring from the southwestern to southeastern areas of the Minamitorishima EEZ. Here we report visual core descriptions and bulk chemical compositions of the deep-sea sediment core samples, and discuss the extent of a highly promising area for future development of REY-rich mud in the Minamitorishima EEZ.

キーワード：レアアース、レアアース泥、南鳥島、海底鉱物資源

Keywords: rare earth elements and yttrium (REY), REY-rich mud, Minamitorishima Island, deep-sea mineral resource

Origin of deep-sea sediments within the Minamitorishima EEZ based on downhole variation of bulk chemical composition and neodymium isotopic ratios

*田中 えりか¹、安川 和孝^{1,2}、中村 謙太郎¹、宮崎 隆³、Vaglarov Bogdan³、藤永 公一郎^{2,1}、岩森 光^{3,4}、加藤 泰浩^{5,1,2,3}

*Erika Tanaka¹, Kazutaka Yasukawa^{1,2}, Kentaro Nakamura¹, Takashi Miyazaki³, Bogdan S Vaglarov³, Koichiro Fujinaga^{2,1}, Hikaru Iwamori^{3,4}, Yasuhiro Kato^{5,1,2,3}

1. 東京大学工学系研究科システム創成学専攻、2. 千葉工業大学、3. 海洋研究開発機構、4. 東京工業大学理学院地球惑星科学系、5. 東京大学工学系研究科エネルギー・資源フロンティアセンター

1. Sys. Innovation, Univ. of Tokyo, 2. Chiba Institute of Technology, 3. JAMSTEC, 4. Dept. of Earth and Planetary Sciences, TITECH, 5. FRCER, Univ. of Tokyo

In 2011, the deep-sea sediments containing a high concentration of rare-earth elements and yttrium (REY) were discovered in the Pacific Ocean [1]. Moreover, the presence of "highly/extremely REY-rich mud" was confirmed within the Japanese exclusive economic zone (EEZ) surrounding Minamitorishima Island in 2013 [2].

On the basis of geochemical characteristics in major- and trace-elemental composition, it was reported that the deep-sea sediment layers within the Minamitorishima EEZ can be classified into several distinct groups including the highly/extremely REY-rich mud, although they are apparently very similar pelagic brown clay [3]. However, the origin and formation mechanism of the highly/extremely REY-rich mud has not been completely unraveled yet.

To elucidate the origin of deep-sea sediment including REY-rich mud within the Minamitorishima EEZ, isotopic composition of neodymium (one of the rare-earth elements), together with major and trace element compositions, can provide an important constraint. We aim to decipher geochemical end-members characterized by distinctive chemical compositions and isotopic ratios that enable us to specify their sources, fluxes and processes of supply [4, 5]. Here, we investigated (1) bulk chemical compositions by XRF and ICP-MS analyses and (2) bulk Nd isotopic ratios using Thermal Ionization Mass Spectrometry (TIMS) throughout a piston core, KR13-02 PC05 of 11.45 m in length, which contains the Extremely REY-rich mud and the other several characteristic layers. We report the downhole variations of bulk chemical composition and neodymium isotopic ratios, and discuss the origin of the highly/extremely REY-rich mud on the basis of their geochemical features.

References

- [1] Kato et al. (2011) *Nature Geoscience* 4, 535-539.
- [2] Iijima et al. (2016) *Geochemical Journal* 50, 557-573.
- [3] Nakamura et al. (2016) *JpGU2016*
- [4] Goldstein, O' Nion and Hamilton (1984) *Earth and Planetary Science Letters* 70,221-236.
- [5] Grousset and Biscaye (2005) *Chemical Geology* 222, 149-167.

キーワード : Nd同位体、化学層序、深海堆積物、南鳥島EEZ、レアアース泥

Keywords: Nd isotopes, Chemostratigraphy, Deep-sea Sediments, Minamitorishima EEZ, REY-rich mud

全岩化学組成に基づく北太平洋の遠洋性深海堆積物層序

A chemostratigraphic correlation of deep-sea pelagic clays in the North Pacific Ocean: Insights from the ODP Sites 1149 and 1179

*見邨 和英¹、山本 克志¹、中村 謙太郎¹、安川 和孝^{1,2}、大田 隼一郎^{3,1}、藤永 公一郎^{2,1}、町田 嗣樹^{3,1}、加藤 泰浩^{1,2,3}

*Kazuhide Mimura¹, Katsushi Yamamoto¹, Kentaro Nakamura¹, Kazutaka Yasukawa^{1,2}, Junichiro Ohta^{3,1}, Koichiro Fujinaga^{2,1}, Shiki Machida^{3,1}, Yasuhiro Kato^{1,2,3}

1. 東京大学大学院工学系研究科、2. 千葉工業大学次世代海洋資源研究センター、3. 海洋研究開発機構

1. School of Engineering, the University of Tokyo, 2. ORCeNG, Chiba Institute of Technology, 3. JAMSTEC

Pelagic clay, referred to as red clay, is one of the common types of seafloor sediment, especially in the Pacific Ocean floor [1]. It is known that the pelagic clay deposits at remote areas in the ocean with considerably slow sedimentation rates of only 0.1-0.5 cm/kyr. Recently, this type of sediment is also recognized as a new deep-sea mineral resource. In 2011, Kato et al. [2] reported that some parts of the deep-sea pelagic clay in the Pacific Ocean contain high concentrations of rare-earth elements and yttrium (REY). They defined the pelagic clay containing more than 400 ppm of total REY as “REY-rich mud” and concluded that this sediment could be a prospective resource for the critical elements.

Pelagic clay is, however, not so well studied as other types of sediments, such as carbonate and neritic sediments, probably due to the lack of visible features and little availability of microfossils to determine its depositional age. Therefore, detailed stratigraphy of the pelagic clay layers including REY-rich mud is not well understood, although it deposited throughout the Cenozoic era [3]. To clarify the stratigraphy of the pelagic clay layers including REY-rich mud, analyses of long and fully recovered deep-sea sediment cores are needed. Here we focused on the ODP Sites 1149 and 1179 in the North Pacific Ocean, both of which were recovered continuously from seafloor to basement rock.

In this study, we provide the results of bulk chemical analyses of sediment samples from the ODP cores. By comparing the multi-elemental compositions of these cores and those of GPC3-LL44 [4], we constructed a general chemostratigraphy of pelagic clay layers in the North Pacific Ocean. We also quantified the relative contributions of each geochemical end-member causing the chemostratigraphic variations of the pelagic clay layers. Based on the results, we discuss the deposition processes of pelagic clays including REY-rich mud in the North Pacific Ocean.

References

- [1] Dutkiewicz et al. (2015) *Geology* 43, 795-798.
- [2] Kato et al. (2011) *Nature Geoscience* 4, 535-539.
- [3] Yasukawa et al. (2016) *Scientific Reports* 6, 29603.
- [4] Kyte et al. (1993) *Geochemica et Cosmochimica Acta* 57, 1719-1740.

キーワード：遠洋性粘土、化学層序、レアアース泥、北太平洋、ODP Site 1149、ODP Site 1179

Keywords: pelagic clay, chemostratigraphy, REY-rich mud, North Pacific Ocean, ODP Site 1149, ODP Site 1179

南太平洋におけるレアアース泥の分布と成因の解明

Genesis and distribution of rare-earth elements and yttrium-rich mud in the South Pacific Ocean

*中村 謙太郎¹、李 斯¹、安川 和孝^{1,2}、見邨 和英¹、藤永 公一郎^{2,1}、大田 隼一郎^{3,1}、町田 嗣樹^{3,1}、加藤 泰浩^{1,2,3}

*Kentaro Nakamura¹, Siyang Li¹, Kazutaka Yasukawa^{1,2}, Kazuhide Mimura¹, Koichiro Fujinaga^{2,1}, Junichiro Ohta^{3,1}, Shiki Machida^{3,1}, Yasuhiro Kato^{1,2,3}

1. 東京大学・工学系研究科、2. 千葉工業大学・次世代海洋資源研究センター、3. 海洋研究開発機構

1. School of Engineering, University of Tokyo, 2. ORCeNG, Chiba Institute of Technology, 3. JAMSTEC

In 2011, Kato et al. [1] reported the potential of deep-sea sediment containing high concentrations of rare-earth elements and yttrium (REY-rich mud) in the Pacific Ocean as a new source for REY. It has been demonstrated that the REY-rich mud are distributed mainly in two regions: the eastern South Pacific and central North Pacific. In the North Pacific Ocean, REY-rich mud has also been discovered within the Japanese exclusive economic zone (EEZ) around Minamitorishima island [2]. Subsequently, detailed geochemical study on the REY-rich mud in the North Pacific Ocean proceeded rapidly, clarifying its distribution, stratigraphic position, and structural components including the host phases of REY [3, 4]. In striking contrast to the substantial progress in the North Pacific Ocean, REY-rich muds in the South Pacific Ocean has still not been well understood, even though their high Σ REY concentrations (> 2000 ppm) [1]. This is, at least in part, due to the lack of sediment cores recovered from the South Pacific Ocean (especially southern part of the South Pacific Ocean). IODP cores recovered by Exp. 329 [5] can provide a unique opportunity to elucidate genesis and distribution of REY-rich mud in the South Pacific Ocean.

In this study, we present the results of geochemical analysis of sediment samples obtained from the IODP Exp. 329 cores and discuss the distribution, stratigraphic position, and constituting components of the South Pacific Ocean REY-rich mud. We also compare our results with those in the North Pacific Ocean.

References : [1] Kato et al. (2011) *Nature Geoscience* 4, 535-539. [2] Iijima et al. (2016) *Geochemical Journal* 50, 557-573. [3] Nakamura et al. (2016) *JpGU Abstract*. [4] Mimura et al. (2017) *JpGU Abstract*. [5] D' Hondt et al. (2011) *Proc. IODP*, vol. 329.

キーワード : レアアース泥、海底鉱物資源、IODP Exp. 329、全岩化学組成、南太平洋

Keywords: REY-rich mud, deep-sea mineral resources, IODP Exp. 329, bulk chemical composition, South Pacific Ocean

前期始新世「超温暖期」のインド洋深海堆積物から統計的に抽出された地球システムのフィードバック機構

Earth system feedbacks statistically extracted from the Indian Ocean deep-sea sediments during the early Eocene hyperthermals

*安川 和孝^{1,2}、中村 謙太郎¹、藤永 公一郎²、加藤 泰浩^{1,2,3}、池原 実⁴

*Kazutaka Yasukawa^{1,2}, Kentaro Nakamura¹, Koichiro Fujinaga², Yasuhiro Kato^{1,2,3}, Minoru Ikehara⁴

1. 東京大学大学院工学系研究科、2. 千葉工業大学次世代海洋資源研究センター、3. 海洋研究開発機構、4. 高知大学海洋コア総合研究センター

1. School of Engineering, The University of Tokyo, 2. Ocean Resources Research Center for Next Generation, Chiba Institute of Technology, 3. JAMSTEC, 4. Center for Advanced Marine Core Research, Kochi University

The most prominent global warming event in the Cenozoic era was the Paleocene-Eocene Thermal Maximum (PETM) at ~56 Ma, which is characterized by a rapid global warming by 5 to 8°C, severe ocean acidification, and a distinct negative carbon isotope ($\delta^{13}\text{C}$) excursion both in the marine and terrestrial realm. These features suggest a massive injection of ^{13}C -depleted greenhouse gas to the ocean-atmosphere system. Moreover, multiple PETM-like global warming episodes termed 'hyperthermals' during the early Eocene period (56~52 Ma), accompanying rapid and pronounced negative excursions in $\delta^{13}\text{C}$, have also been recognized over the past dozen years. Geologic records of the hyperthermals have so far been reported from around the globe (e.g., the Pacific, Atlantic, and Arctic Oceans, Europe and North America). However, albeit the third largest ocean on the planet, the Indian Ocean is almost a blank area where only a few published data of the hyperthermals are available. Here we have constructed a comprehensive geochemical data set including major- and trace-element contents, $\delta^{13}\text{C}$, and CaCO_3 contents of 250 bulk sediment samples taken from ODP Sites 752 and 738, located in the southeastern Indian Ocean and the Indian sector of the Southern Ocean, respectively. The analytical results show that the sediments of these cores record multiple carbon isotope excursions and reductions of carbonate contents, probably corresponding to the PETM and the early Eocene hyperthermals including the Eocene Thermal Maximum 2 (ETM2), H2 and I1/I2 events, and ETM3. We applied Independent Component Analysis to the high-dimensional compositional data matrix, and extracted four geochemical independent components that collectively account for 85.6% of the total sample variance. One of the components involving Ba content and $\delta^{13}\text{C}$ indicates a signature of a negative feedback in Earth system that efficiently sequestered the excess carbon in recovery phases of the hyperthermals.

キーワード：深海堆積物、インド洋、気候変動、前期始新世超温暖期、独立成分分析、多変量解析

Keywords: deep-sea sediment, Indian Ocean, climate change, Eocene hyperthermals, Independent Component Analysis, multivariate analysis

Geochemistry and Re-Os and ^{146}Sm - ^{142}Nd isotope systematics of mafic rocks in the Acasta Gneiss Complex: Discovery of the oldest terrestrial rock and implications for the Earth-forming materials

*小宮 剛¹、越田 溪子¹、石川 晃¹、鈴木 勝彦²

*Tsuyoshi Komiya¹, Keiko Koshida¹, Akira Ishikawa¹, Katsuhiko Suzuki²

1. 東京大学大学院総合文化研究科広域科学専攻、2. 国立研究開発法人海洋研究開発機構・海底資源研究開発センター
1. Department of Earth Science & Astronomy Graduate School of Arts and Sciences The University of Tokyo, 2. Research and Development Center for Submarine Resources, Japan Agency for Marine-Earth Science and Technology

The first billion years' history of the earth is still poorly understood because terrestrial rocks with the ages are scarcely preserved. The Acasta Gneiss Complex (AGC) is one of the rare Eoarchean geologic bodies, located in the western margin of the Slave Craton, Canada, and is composed of felsic to intermediate gneisses with subordinate amounts of mafic rocks. This study aims to obtain physico-chemical constraints on the Hadean mantle and propose a new picture of early mantle evolution based on geological, petrological and geochemical investigations of the oldest mafic rocks in AGC. The mafic rocks mainly consist of fine to coarse-grained hornblende and plagioclase with small amount of quartz, chlorite, garnet and ilmenite. They commonly suffered from amphibolite to upper amphibolite facies metamorphism. The mafic rocks are subdivided into three groups based on the rare earth element (REE) patterns: highly variable light REE-enriched pattern, flat REE pattern and slightly light REE-enriched pattern with positive Eu anomaly, respectively. The samples, which have flat REE patterns and whose high field strength element (HFSE) and REE contents are well correlated with immobile Zr contents, were selected as the least altered samples to estimate their source mantle because the infiltration of metamorphic fluids/melts increases the light REE contents relative to the middle and heavy REE contents and more severely disturbs other trace element contents than Zr contents. The least altered mafic rocks have chondritic trace element relative abundances with negative Nb and Ta anomalies. This implies that the Nb and Ta were partitioned into the metallic core to form a Nb and Ta-deficit primitive mantle. The least altered mafic rocks show a positive correlation on a $^{187}\text{Re}/^{188}\text{Os}$ vs $^{187}\text{Os}/^{188}\text{Os}$ diagram, yielding a formation age of 4272 ± 300 Ma. The age is consistent with the field occurrence of mafic rocks because they were intruded by orthogneisses with 4.03 to 3.6 Ga ages. The highly radiogenic initial $^{187}\text{Os}/^{188}\text{Os}$ ratio suggests that their source material was a pre-late veneer mantle with a high Re/Os ratio. However, geochemical signatures, which the initial $^{187}\text{Os}/^{188}\text{Os}$ ratio overlaps with a chondritic value within the error and their highly siderophile element (HSE) abundances are similar to those of modern basalts, indicate that their source mantle had modern mantle-like high HSE contents, implying that the late veneer event and later homogenization took place before 4.27 Ga.

Their $^{142}\text{Nd}/^{144}\text{Nd}$ ratios are identical to those of a modern mantle with a suprachondritic value. Their chondritic initial $^{143}\text{Nd}/^{144}\text{Nd}$ ratio and REE patterns, and suprachondritic $^{142}\text{Nd}/^{144}\text{Nd}$ ratios indicate that their source material had a chondritic Sm/Nd ratio and a suprachondritic (modern mantle value) $^{142}\text{Nd}/^{144}\text{Nd}$ ratio at 4.27 Ga. This is the first evidence that > 3.7 Ga source mantle had a modern mantle-like (normal) $^{142}\text{Nd}/^{144}\text{Nd}$ ratio. Two possible scenarios account for the Hadean mantle with the primitive mantle-like trace element contents and normal $^{142}\text{Nd}/^{144}\text{Nd}$ ratio at 4.27 Ga. One is that the early mantle convection was rapid enough to homogenize large-scale mantle heterogeneity due to a magma ocean until 4.27 Ga. The Acasta mafic rocks was derived from the homogenized primitive mantle, whereas the 3.8-3.7 Ga Isua mafic rocks with excess μ ^{142}Nd values were formed from a shallow depleted mantle possibly due to progressive extraction of primitive crusts. This scenario is supported by the HSE contents

and ^{187}Re - ^{187}Os isotopes. Another model suggests that the extent of a magma ocean was limited and non-melting primitive parts remained in the deep mantle, and that the Acasta mafic rocks were formed from the deep primitive mantle, whereas the Isua mafic rocks were derived from the early differentiated upper mantle.

キーワード：冥王代、初期分化

Keywords: Hadean, Early differentiation

Hydrothermal silicification of mafic and ultramafic rocks in the Barberton Greenstone Belt, South Africa: Relationship between bioessential element influxes into the seawater and secular variation of atmospheric CO₂ contents

*小宮 剛¹、梅田 早織¹、石川 晃¹

*Tsuyoshi Komiya¹, Saori Umeda¹, Akira Ishikawa¹

1. 東京大学大学院総合文化研究科広域科学専攻

1. Department of Earth Science & Astronomy Graduate School of Arts and Sciences The University of Tokyo

The presence and composition of liquid water on the earth's surface through geologic time are essential to sustain biological system. Therefore, it is important to understand the chemical evolution of elemental cycles of an oceanic environment. Especially, it is necessary to understand ocean floor alteration of oceanic crusts because they supply bioessential elements into the ocean, but the quantitative estimates still lack for the Archean hydrothermal systems.

We analyzed major and trace element contents of the silicified, carbonated and metamorphosed (non-silicified/non-carbonated) volcanic rocks, including sixty-five basalts and seventy-three peridotitic and basaltic komatiites and sixteen overlying cherts in the Komati and Hooggenoeg formations in the Onverwacht Group of the Mesoarchean Barberton Greenstone Belt, South Africa. Some of the basalt and komatiites underwent severe hydrothermal silicification and carbonation as well as hydrothermal metamorphism up to the amphibolite facies condition. The silicified peridotitic komatiites are more enriched in SiO₂, Rb, Ba and U contents and depleted in FeO^{tot}, MgO, CaO, transition elements such as Ni and Co, and Sr contents than the non-silicified/carbonated basalts and peridotitic komatiites. The carbonated basalts and peridotitic komatiites are enriched in CaO, Rb, Ba, U and Sr, and depleted in FeO^{tot}, MgO, and the transition element contents.

The chemical contrast indicates that Co and Ni are significantly released from the peridotitic komatiites due to the silicification and carbonation. We performed petrographic and petrological study of the peridotitic komatiites to understand the petrological cause. The non-silicified/non-carbonated peridotitic komatiites commonly contain olivine and serpentine, but the silicified and carbonate rocks contain quartz, potassium-rich white micas and carbonate minerals, replacing the magmatic minerals. The silicate minerals such as the serpentine and olivine can host the transitional metals such as Co and Ni, but the alteration minerals such as quartz, potassium-rich white micas and carbonate minerals cannot contain the transition elements compared with the serpentine and olivine. The disappearance of the serpentine and olivine due to the silicification and carbonation caused decrease of Co and Ni contents of altered peridotitic komatiites, namely supply of the transition elements into the seawater.

Previous study suggested that the basalts under the sedimentary cherts underwent silicification but avoided carbonation, whereas the basalts under the silicification zone suffered from only the carbonation. However, our geological mapping showed that the upper part of the basalt (komatiite) unit suffered from both silicification and carbonation. In addition, disseminated carbonate minerals are replaced by quartz in some partially silicified rocks. The evidence indicates that the carbonation was followed by the silicification because of change of pH condition of the hydrothermal fluid from alkaline to acidic conditions possibly due to mixing with acidic seawater. The change of pH condition caused that the carbonate minerals were dissolved and replaced by silica minerals.

The carbonation of ocean floor basalts was caused by high atmospheric CO₂ contents and ceased due to its decrease around 2.7 Ga. On the other hand, Ni and Co contents of banded iron formation indicate that

Ni and Co contents of seawater also decreased around 2.7 Ga. Previous work suggested that the decrease of Ni contents was due to decrease of komatiitic volcanic activity in the Late Archean. But, it is well known that the komatiite occurred at least until 1.9 Ga. This study proposes that the decrease of the marine Ni and Co contents was not due to disappearance of peridotitic komatiites but decrease of atmospheric CO₂ contents in the Late Archean because the silicification and/or carbonation are necessary to supply Ni and Co from oceanic crust to seawater.

キーワード：珪化作用、海洋遷移元素量、生命必須元素、コマチアイト、海洋組成の経年変化

Keywords: Archean silicification, marine transition element content, bioessential element, komatiite, secular change of seawater composition

Geochemical mapping of living animal eggs and embryos, and the Ediacaran Weng' an embryo fossils for taxonomic identification of the oldest animal embryo fossils

*小宮 剛¹、大河原 綾乃¹、大久保 奈弥²、坂田 周平³、大野 剛³

*Tsuyoshi Komiya¹, Ayano Ohkawara¹, Nami Okubo², Shuhei Sakata³, Takeshi Ohno³

1. 東京大学大学院総合文化研究科広域科学専攻、2. 東京経済大学経済学部、3. 学習院大学

1. Department of Earth Science & Astronomy Graduate School of Arts and Sciences The University of Tokyo, 2. Graduate School of Economics, Tokyo Keitai University, 3. Gukushuin University

The Weng' an biota is an Ediacaran biotic group, and has some unique features such as phosphatization of soft tissues of the original organisms and predominance of microfossils. This study focuses on the spherical phosphatized microfossils, which have multi-layered chorions with spiny or reticulated decorations on those surfaces, and whose internal spheres are divided into one to several hundred vesicles. Previous studies have interpreted them as animal embryos because of the palintomic cell division, but the taxonomic identification, namely animal phylum, is still on debate because the morphology is quite different from any modern animal eggs/embryos. In order to identify a taxonomic group of original organisms, more quantitative assessment than the physical analysis should be required. This study tries to establish chemical methodology to study the Ediacaran fossils, and performed four geochemical analyses: micro-CT analyses for reconstruction of three-dimensional structures of the spheroidal fossils without the destruction, laser Raman analyses to obtain distribution of organic matter within the fossils, compositional mapping of twelve trace element contents (B, Na, Al, Mn, Fe, Co, Ni, Cu, Zn, Sr, Ba, and Pb) on red algae and eggs, embryos and gastrulas of living cnidarian Anthozoa and mollusk Gastropoda, and compositional mapping on the four *Megasphaera* (1-cell) stage fossils and two *megaclonophycus* (many cell) stage fossils in the Weng' an. The chemical analyses were performed with LA-ICP-MS, housed at the Gakushuin University.

The compositional mapping shows that the trace element distribution in the living organisms is greatly different not only between algae and animal eggs but also between animal eggs of different phyla. The cnidarian eggs show a uniform distribution in boron and iron contents and enrichment in Na and Cu contents along the rims. The distribution of Ni, Zn, Sr, Ba and Pb contents within the eggs changes with the development stages. The Ni contents are uniformly distributed at the cleavage and gastrulation stages, whereas are concentrated in the outer parts at the blastocyst stage. Zn, Sr, Ba and Pb contents are concentrated in the outer parts at the cleavage and the blastocyst stages, and are uniformly distributed in the gastrulation stage. Al, Mn and Co could not be detected. On the other hand, internal parts of the mollusk eggs are enriched in Na, Co, Cu and Zn contents whereas the rim parts are enriched in Mn, Fe, Sr, Ba and Pb contents. The centers of the red algae are abundant in B, Na, Ni, Cu, Zn, Sr, Ba and Pb contents. Mn and Fe could not be detected in the red algae.

The trace element contents are not necessarily correlated with their requirement; for example Fe, a bioessential element for photosynthesis, contents are quite low in the algae. On the other hand, the trace element mapping reveals that Sr, Pb and Ba contents are high in the rims at the early development and decrease with development. We considered the reason why they are removing from the eggs with development because they are non-bioessential elements. The difference in distribution of bioessential elements in the eggs between the cnidarian and mollusk can be used to identify the phylum of an egg. We also conducted mappings of fifteen elements including Cr, I and U on the Weng' an fossils. We considered transitional distribution of B, Al, Mn, Fe, Sr and Ba contents between the fossils and

surrounding dolomite, and uniform distribution of Co and Ni within the fossils as taphonomic homogenization. Because Cu is highly distributed in enclosed vesicles of all fossils and the Cu contents sharply change between the fossils and the surrounding, Cu possibly represents a common feature of the original organism of the animal embryos. We compared distributions of Cu, Zn and Pb between the living organisms and Weng' an animal embryos. The distributions are different between them so that we cannot identify the fossils as cnidarian or molluscs.

キーワード：エディアカラ紀、陡山沱、動物胚化石

Keywords: Ediacaran, Doushantuo, Animal embryo fossils

Mid-Archean low geothermal gradient metamorphism: constraints from phase relationships in metamorphosed BIF from the Inyoni shear zone of the Barberton granite-greenstone belt, South Africa

*青木 一勝¹、山本 伸次²、澤木 佑介³、浅沼 尚³、小宮 剛⁴

*Kazumasa Aoki¹, Shinji Yamamoto², Yusuke Sawaki³, Hisashi Asanuma³, Tsuyoshi Komiya⁴

1. 岡山理科大学基礎理学科、2. 横浜国立大学環境情報研究院、3. 東京工業大学地球惑星科学専攻、4. 東京大学総合文化研究科

1. Department of Applied Science Okayama University of Science, 2. Graduate School of Environment and Information Sciences, Yokohama National University, 3. Department of Earth and Planetary Sciences, Tokyo Institute of Technology, 4. Graduate School of Arts and Sciences, The University of Tokyo

The Barberton granite-greenstone belt is one of the oldest and best-preserved examples of Archean geology in the world. High-P amphibolitic rocks associated with ca. 3.23 Ga subduction-collisional event were reported from the Inyoni shear zone situated south of the Barberton belt (Dziggel et al., 2002; Moyen et al., 2006; Nédélec et al., 2012; Cutts et al., 2013). According to those studies, the high-P rocks formed under geothermal gradients of ca. 12–30°C/km, which is similar to those found in the Phanerozoic subduction and collision zones. Therefore, other metamorphic rocks exposed in the Inyoni shear zone must preserve mineralogical and petrological evidences showing such a low geothermal gradient metamorphism. In this study, in order to verify the low geothermal gradients at the mid-Archean subduction-collision zone, quartz-rich layers in metamorphosed BIF from the Inyoni shear zone have been examined for the metamorphic P-T conditions, based on the phase relationships combined with microscopic observation, mineral compositions and thermodynamic calculations.

Studied samples (BF152 and BF153) are composed mainly of quartz, garnet, grunerite, amphibole, clinopyroxene and magnetite. Taking into account the microscopic observation and EPMA analyses (WDS mode) of each mineral, equilibrium mineral assemblage during the metamorphism are shown as follows; assemblage 1 (A1): Grt+Cpx+Gru+Mag+Qtz --> assemblage 2 (A2): Grt+Cpx+Gru+Hbl+Mag+Qtz --> assemblage 3 (A3): Act+Mag+Qtz+Gru or Hbl.

To provide a framework for understanding the change of mineral assemblages and to constrain the metamorphic P-T conditions, P-T pseudosection and isopleth calculations have been performed by forward modeling with a computer program PERPLEX ver. 6.6.6 (Connolly, 2005 and its update) with an internally consistent dataset of Holland & Powell (1998 and its update). Moreover, garnet-hornblende geothermometer (Graham & Powell, 1984 and Perchuk et al., 1985) using the program THERMOBAROMETRY ver. 2.1 (Spear & Kohn, 1999) and the average P-T calculations of THERMOCALC ver. 3.33 with the computer program AX2 (Holland and Powell, 1998 and its update) were carried out as necessary. The bulk rock composition was confirmed by XRF analysis. The chemical states of iron and their relative abundances in the sample were measured by using ⁵⁷Fe Mössbauer spectroscopy. The effective bulk compositions of each assemblage were calculated by volume estimates combined with chemical analysis and abundance ratio of the minerals.

The calculated pseudosection diagrams show that the stability field of each assemblage are about over 10 kbar and 580–690°C (A1), over 4 kbar and 500–600°C (A2) and lower than 4 kbar and 530°C (A3), respectively. These results are consistent with the change of mineral assemblage inferred by the petrography for the samples. These results lead us to that the samples record a series of metamorphism from peak to retrograde conditions. The P-T conditions of A2 given in combination with average P calculation and Grt-Hbl geothermometry are P = ca. 10 kbar and T = ca. 512–538 °C. These results correspond with both the equilibrium phase diagrams and isopleth results. On the other hand, the

isopleth result of A1 is not consistent with the stability field of A1 in the pseudosection diagram. This suggests that the chemical compositions of minerals that up the A1 were modified at a late stage where A2 formed. Thus, we cannot constrain the specific P-T conditions of A1, but we may say at least that A1 underwent higher-grade metamorphism than A2.

The estimated P-T conditions indicate that the A2 formed under geothermal gradients of ca. 15–20°C/km. This gradient corresponds to previous works for the highest-grade rocks in the same area (Dziggel et al., 2002; Moyen et al., 2006). Furthermore, our study suggests that the metamorphism that A1 underwent was higher P-T conditions than that of A2. Hence, it is possible that the geothermal gradients along the subduction-collisional zone when A1 formed were almost as much as or lower than ca. 15–20°C/km. Such gradient gives close agreement with that of subduction zone like Japan and of collision zone such as Himalaya and Kokchetav Massif. These features suggest the possibility that the mid-Archean crust was sufficiently cool and rigid as already mentioned by Moyen et al. (2006), and some of the crustal materials were subducted into much deeper depths than previously considered.

Petrological study of clinopyroxene-bearing garnet amphibolite in the Barberton granite-greenstone belt, South Africa

*加藤 大地¹、青木 一勝¹、小宮 剛²

*Daichi Kato¹, Kazumasa Aoki¹, Tsuyoshi Komiya²

1. 岡山理科大学、2. 東京大学

1. Okayama University of Science, 2. The University of Tokyo

The approximately 3.5-3.2 Ga Barberton granite-greenstone belt (BGGB) is one of the oldest and best-preserved examples of Archean geology in the world. Cpx-bearing Grt amphibolites were reported from the Inyoni Shear zone in south of this belt (Moyen et al., 2006). These rocks record the highest-grade metamorphism compared with other rocks in this belt, and are considered to form under geothermal gradients of ca. 12-20°C/km, which is similar to those found in recent subduction zone. However, the specific metamorphic P-T trajectory from subduction to exhumation has not yet been clarified. In this study, petrological study for the Grt-amphibolite has been examined for mineral assemblages and compositions in detail to discuss the metamorphic history of the BGGB.

Microscopic observation has indicated that the studied samples contain quartz (Qz), garnet (Grt), amphibolite (Amp), clinopyroxene (Cpx), plagioclase (Pl) and epidote (Ep) with minor opaque mineral. Chemical compositions obtained from each mineral by EPMA analyses are as follows; Grt, Cpx and Pl compositions were homogeneous and no chemical zoning was confirmed. Amps were defined as Ca-amphibole. There was a little difference of Al contents between core-mantle (Amp1) and rim (Amp2) part. Amp2 growth was confirmed along fracture part in Grt and the circumference of Cpx. Epidote occurred as matrix grains (Ep1) and component of symplectite (Ep2 + Qz) which is originally Grt. Mn content of Ep1 was slightly higher than that of Ep2. The petrography and mineral compositions indicated that the studied samples record the change of mineral assemblage from Grt + Amp1 + Cpx + Pl + Qz + Ep1 (AS1) to Grt + Amp2 + Cpx + Pl + Qz + Ep2 (AS2).

The metamorphic P-T conditions have been estimated by garnet-clinopyroxene geothermometer (Ai, 1994; Nakamura, 2009) and the average P-T calculations of THERMOCALC ver. 3.3.3 with the computer program AX (Holland and Powell, 1998 and its update). P-T pseudosection and isopleth calculations were performed by a computer program PERPLEX ver. 6.7.4 (Connolly, 2005 and its update). These results showed that the stability P-T field of AS1 and AS2 are at P = ca. 9.8-11.3 kbar and T = ca. 625-675 °C.

The estimated P conditions are lower than that of previous work in the same area (Moyen et al., 2006). Our samples probably record the later stage metamorphism. Integration of our new results with published data suggest that the geothermal gradients at the prograde metamorphism associated with 3.23 Ga subduction-collisional event was not at least exceeding ca. 20°C/km and the initial retrograde P-T trajectory was isothermal decompression or decompression with increasing temperature like a clockwise path.

Distribution and zircon U-Pb dating of the Idiwhaa Tonalitic Gneiss in the Acasta Gneiss Complex

*上原 啓幹¹、越田 溪子¹、山本 伸次²、石川 晃³、青木 翔吾³、坂田 周平⁴、小宮 剛³

*Hiroki Uehara¹, Keiko Koshida¹, Shinji Yamamoto², Akira Ishikawa³, Shogo Aoki³, Shuhei Sakata⁴, Tsuyoshi Komiya³

1. 東京大学大学院理学系研究科地球惑星科学、2. 横浜国立大学環境情報研究院、3. 東京大学大学院総合文化研究科広域科学専攻、4. 学習院大学理学部化学科

1. Department of Earth and Planetary Science, Graduate School of Science, University of Tokyo, 2. Graduate school of Environment and Information Sciences, Yokohama National University, 3. Department of Earth Science and Astronomy, Graduate School of Arts and Sciences, University of Tokyo, 4. Department of Chemistry, Faculty of Science, Gakushuin University

It is assumed that the earth experienced many significant events, such as giant impact, magma ocean, core formation, large-scale mantle differentiation and early crustal formation in the Hadean. However, the investigation for the early earth evolution is basically based on geochemistry of meteorites and lunar rocks so that the quantitative analyses are still poorly understood and need the terrestrial rocks. So far, the oldest terrestrial rock was found in the Acasta Gneiss Complex (AGC) of the western part of the Slave Province, Canada (Bowring and Williams., 1999), and goes back to 4.03 Ga on the basis of U-Pb dating of zircons from the orthogneisses. The AGC comprises the 3.6-4.0 Ga felsic and layered gneiss suites with minor mafic rocks. In addition, a Hadean material of a 4.2 Ga zircon xenocryst was also found from ca. 3.94 Ga tonalitic gneiss. It is expected that the Hadean granitic crust occurred in the area.

Recently, Reimink et al. (2014) discovered the Idiwhaa Tonalitic Gneiss (ITG) with well-preserved 4.02 Ga igneous zircons in the northern part of the AGC. The ITG is garnet-biotite-hornblende tonalitic gneiss, and has high total iron contents, low Mg-numbers, and flat REE patterns with negative Eu anomalies, different typical Archean TTGs. They concluded that the tonalite magma was formed in a plume-related tectonic setting. Moreover, Reimink et al. (2016) reported the ITG unit near a central area of the AGC.

We carried out geological survey in this area in 2015 to obtain more detailed distribution of ITG unit over the AGC. The geology and geochemistry of the orthogneisses indicate that some orthogneisses in the southern part of the AGC are also related with the ITG, and suggests that the ITG unit is extensively distributed over the AGC. In addition, we will report the U-Pb ages of zircons from their rocks with LA-ICP-MS.

キーワード：アカスタ片麻岩体、ジルコン、U-Pb年代

Keywords: Acasta Gneiss Complex, Zircon, U-Pb dating

世界最古(>3.92Ga)の堆積岩の地質学的産状と化学組成から推定する初期 太古代の海洋組成

Occurrence and compositions of the oldest sedimentary rocks in the Nulliak supracrustal rocks, Labrador (>3.92 Ga): Implications for a chemical composition of the Eoarchean seawater

*吉田 聡¹、越田 溪子¹、石川 晃²、小宮 剛²

*Yoshida Satoshi¹, Keiko Koshida¹, Akira Ishikawa², Tsuyoshi Komiya²

1. 東京大学理学系研究科地球惑星科学専攻、2. 東京大学総合文化研究科広域科学専攻

1. Department of Earth and Planetary Science, The University of Tokyo, 2. Department of Earth Science & Astronomy
Graduate School of Arts and Sciences The University of Tokyo

The earth is a unique planet where liquid water and life exist through geologic time. Therefore, estimate of a surface environment on the early earth is very important to understand the origin of life on the earth and gives an insight into the possibility of life on the earth-type planets. However, we have little knowledge of the environment of the early earth, because the first 500 m.y. history of the earth was almost completely lost. The surface environment of the early earth has been ever investigated from geochemistry and geobiology of the Isua supracrustal belt, southern West Greenland because the supracrustal rocks had the oldest age of 3.8 to 3.7 Ga up to now. Recently, older (>3.92 Ga) supracrustal rocks, called Nulliak supracrustal rocks, were found in the Saglek Block, northern Labrador. The supracrustal rocks contain banded iron formation (BIF), carbonate rocks, and clastic rocks of pelitic rocks and conglomerates as well as mafic and ultramafic rocks. Although the Isua supracrustal belt suffered from the extensive carbonation so that the carbonate rocks are skeptically considered as chemical sediments, the Nulliak supracrustal rocks avoided the later carbonation. In this study, we analyzed major and trace element compositions of carbonate rocks to estimate the trace element compositions of the Eoarchean seawater.

The Saglek block is located in the northeastern part of the Labrador Peninsula, northeast Canada, and is underlain by the Archean orthogneisses and supracrustal rocks. They underwent granulite to amphibolite facies metamorphism but escaped extensive carbonation and silicification. The supracrustal rocks contain ultramafic and mafic rocks, BIF, chert, carbonate rocks, pelitic rocks and conglomerates, and were intruded by the Mesoarchean mafic dikes (Saglek Dyke) and Eoarchean orthogneisses. Especially, in the St. John' s Harbour South area, the supracrustal rocks were intruded by the 3.92 Ga orthogneiss, indicating the supracrustal rocks have over 3.92 Ga age so that they are the oldest supracrustal rock in the world.

The carbonate rocks occur in the St. John' s Harbour South (SJHS), St. John' s Harbour East (SJHE), Big Island and Pangertok Inlet areas. We classified the carbonate rocks into three groups based on the field occurrence. The first group is characterized by the occurrence within mafic/ultramafic rock units and is composed of the carbonate rocks in the SJHS. The second group occurs between the BIFs and mafic volcanic rocks, and the type locality is the Pangertok Inlet. The third group comprises the carbonate rocks interlayered with or underlying the pelitic rocks in the SJHE and Big Island.

We analyzed major and trace element (Rb, Sr, Y, Zr, Nb, Cs, Ba, rare earth element (REE), Hf, Ta, Pb, Th, and U) contents of the carbonate rocks. Some carbonate rocks have high Zr, Ti and Al₂O₃ contents, possibly due to involvement of detrital and volcanic materials. We selected the carbonate rocks with low Zr, Ti and Al₂O₃ contents in order to remove the influence of contamination of the detrital and volcanic materials because they are not incorporated into the carbonate minerals. Their PAAS-normalized REE+Y

patterns display flat to slightly LREE-depleted patterns with positive La, Eu and Y anomalies and without Ce anomaly. The positive Y anomalies indicate that the carbonate rocks were chemical sediments precipitated from seawater, and the lack of the Ce anomaly indicates that the Eoarchean seawater was anoxic as many previous works suggested. The presence of the Eu anomalies suggests that the carbonate rocks were deposited in marine environments strongly influenced by hydrothermal fluid.

The Eu anomalies are apparently correlated with lithostratigraphy and accompanied rocks; those in SJHS have larger positive Eu anomalies than those in the SJHE and Big Island, possibly due to the degree of the influence of hydrothermal influxes related to the distance from the volcanic sources. On the other hand, even the carbonate rocks accompanied with pelitic rocks have the distinct Eu anomalies, indicating that the seawater surrounding a continent had the Eu anomaly so that the seawater was dominated by hydrothermal influxes. In addition, there is a positive correlation between the Y and Eu anomalies.

キーワード：初期太古代、初期生命、表成岩

Keywords: Eoarchean, Early life, Supracrustal rocks

縞状鉄鉱層の遷移金属元素濃度の空間分布：海洋生命必須元素濃度の経年変化と生物進化への影響

Transitional-element compositions at whole-rock and micro scale of the Banded Iron Formations : Implications for secular variations of bioessential elements in the ocean and their influences on biological evolutions.

*青木 翔吾¹、坂田 周平²、中田 亮一³、柏原 輝彦³、大野 剛⁴、高橋 嘉夫⁴、Tsikos Harilaos⁵、小宮 剛¹

*Shogo Aoki¹, Shuhei Sakata², Ryoichi Nakada³, Teruhiko Kashiwabara³, Takeshi Ohno⁴, Yoshio Takahashi⁴, Harilaos Tsikos⁵, Tsuyoshi Komiya¹

1. 東京大学大学院総合文化研究科、2. 学習院大学理学部化学科、3. 海洋研究開発機構、4. 東京大学大学院理学研究科、5. Department of Geology, Rhodes University

1. Graduate School of arts and science, University of Tokyo, 2. Department of Chemistry, Gakushuin University, 3. Japan Agency for Marine-Earth Science and Technology, 4. Graduate School of Science, the University of Tokyo, 5. Department of Geology, Rhodes University

It is generally thought that the surface environments have evolved from anoxic to oxic conditions throughout the earth history. The increase in oxygen of the atmosphere has provoked the evolutions of many genera and metabolic types. Especially, the increase in atmospheric oxygen around the Great Oxidation Event might have provoked the evolutions of the eukaryotic organisms in the Paleoproterozoic, which are more dependent on bioavailable oxygen (Albani et al., 2010; Han and Runnegar, 1992). On the other hand, all organisms are dependent on various transitional elements, which are used mainly for some metabolic coenzymes. The relative dependencies on the elements are different depending on their taxa and their metabolic types (Zerkle et al., 2005; Dupont et al., 2006). For example, prokaryotes are more dependent on Co than eukaryotes for synthesizing methionine, one of the essential amino acids (Dupont et al., 2006). Therefore, secular variations of transitional element compositions in seawater might have been influential on the biological evolutions.

Chemical compositions of the banded iron formations (BIFs) can be proxies for the chemical composition of seawater. In this study, we analyzed the transitional element compositions, especially for Co, of the BIFs from the Eoarchean to Paleoproterozoic at micro to whole-rock scales to estimate the secular variations of the bioavailabilities of those elements.

The Eo- and Mesoarchean BIFs show that whole-rock Co compositions are positively correlated with Fe contents, and not positively correlated with their Al, Ti and HFSE contents. Moreover, the elemental distribution mappings of the Mesoarchean BIFs show that the distribution of Co corresponds to Fe-rich bands. On the other hand, the Paleoproterozoic BIFs in the Hotazel Formation, South Africa, which are interlayered with manganeseiferous sedimentary rocks, show positive correlations of bulk Mn contents with Co contents. The micro-scale distribution mappings of elements in the Mn-poor BIFs, which have lower Co contents than the Eo- to Mesoarchean BIFs, show that the distribution of Co corresponds to not Fe-enriched bands but Al-enriched spots.

Positive correlations of Co with Fe contents in the Eo- and Mesoarchean BIFs at whole-rock and micro scale suggest that much Co in them is derived from adsorbed element on Fe oxyhydroxide in the contemporary seawater. On the other hand, the positive correlations between Mn and Co contents in the Paleoproterozoic BIFs and Mn rocks suggest that much of Co in them exists in Mn-oxide minerals.

Moreover, the Co distribution in Mn-poor BIFs corresponding to Al-enriched spots indicates that Co in them is derived not from the seawater, but from the clastic materials. Therefore, bioavailable Co in the Paleoproterozoic ocean might be limited because divalent Co was fixed as insoluble trivalent forms within Mn-oxide minerals.

Cobalt is mainly used for cobalamin (vitamin B12), which is concerned with every metabolic activity in cell. For example, methylated cobalamin (Met H) catalyze the synthesis of methionine, which is an amino acid constituting the terminal domain of all proteins (e.g. Croft et al., 2005). Only prokaryote can synthesize Met H from dissolved Co in aquatic environment, and depend more heavily on Co than eukaryotes (Dupont et al., 2006). On the other hand, a small group of prokaryotes and eukaryotes can use Zn-bearing enzyme (Met E) in place of Co, and they belong to a monophyletic group (Helliwell et al., 2011).

The decrease in bioavailability of Co during the period from the Mesoarchean to the Paleoproterozoic might have encouraged the prosperity of Co-independent prokaryotes, which were the ancestors to the eukaryotes.

キーワード：縞状鉄鉱層、遷移金属元素、生命必須元素

Keywords: Banded Iron Formations, Transitional Elements, Bioessential Elements

Hydrogen Isotopic Composition of Archean Seafloor Basalts & Gabbros

*村居 呂一¹、上野 雄一郎¹、Julien Foriel

*Tomokazu Murai(Ueno)¹, Yuichiro Ueno¹, Julien Foriel

1. 東京工業大学

1. Tokyo Institute of Technology

The oceans are the main water reservoir in Earth's surface. The evolution of the hydrogen isotopes composition of seawater reflects the hydrogen budget in Earth's ocean and atmosphere through time. We have analyzed hydrogen and oxygen isotopic composition of Archean seafloor basalts and gabbros in 3.2 Ga Pilbara Cleaverville Formation, Western Australia. The Cleaverville Formation has undergone regional sub-greenschist to amphibolite facies metamorphism and was interpreted by Shibuya et al., 2007 as an metamorphized ophiolite. So it is possible that hydrated Cleaverville Formation basalts and gabbros preserved δD reflecting the seawater value at 3.2 Ga.

Our results show a clear correlation between metamorphic facies in hydrated ocean-floor minerals and H and O isotopic compositions. The measured δD values are positively correlated with $\delta^{18}O$ values and H_2O content in response to metamorphic grade. By analogy with modern seafloor basalts, the correlation between δD and H_2O content implies seafloor basalt hydrothermal alteration and dehydration with metamorphism. From this hypothesis, we estimate that seawater at 3.2 Ga was depleted in deuterium by about 20% compared to modern seawater.

キーワード：水素同位体、大古代、地球史

Keywords: Hydrogen isotope, Archean, Earth's History

ガーナ・ビリミアン帯ケープスリーポイントにおける陸上掘削について; GHBコアの詳細な岩相と層序

Lithologies and stratigraphy of GHB cores drilled at Cape Three Points area in the Birimian Supergroup, Ghana

*坂井 志緒乃¹、清川 昌一¹、伊藤 孝²、池原 実³、山口 耕生⁴、尾上 哲治⁵、吉丸 慧¹、三木 翼¹
、George.M. Tetteh⁶、Frank.K. Nyame⁷

*Shiono Sakai¹, Shoichi Kiyokawa¹, Takashi Ito², Minoru Ikehara³, Kosei E. Yamaguchi⁴, Tetsuji Onoue⁵, Satoshi Yoshimaru¹, Tsubasa Miki¹, George M Tetteh⁶, Frank K Nyame⁷

1. 九州大学 大学院理学府 地球惑星科学専攻、2. 茨城大学教育学部、3. 高知大学海洋コア総合研究センター、4. 東邦大学, NASA Astrobiology Institute、5. 熊本大学大学院自然科学研究科、6. University of Mining and Technology, Ghana、7. University of Ghana

1. Department of Earth and Planetary Sciences, Graduate School of Sciences, Kyusyu University., 2. Faculty of Education, Ibaraki University, 3. Center for Advanced Marine Core Research, Kochi University, 4. Toho University and NASA Astrobiology Institute, 5. Earth and Environmental Sciences, Graduate School of Science and Technology, Kumamoto University, 6. University of Mining and Technology, Ghana, 7. University of Ghana

初期原生代 (2.5-2.1Ga) は、生命による光合成の結果生じた寒冷化、地球表層の酸素濃度が飛躍的に上昇し酸化環境へシフトした大酸化事変 (GOE)、初期大陸の形成など地球表層環境に大きな変化が生じた時代として知られる (van Kranendonk, 2012)。本研究は初期原生代の酸素濃度上昇時期の深海底環境の復元を目指し、ガーナ・ビリミアン帯のグリーンストーン帯中の堆積岩についての陸上掘削コア (GHBコア: 2015年掘削) について詳細な岩相記載・層序復元を行った。

ガーナ南西部に位置する西アフリカクラトン西部には太古代の地質体が分布しており、南東部には古原生代の地質体であるビリミアン超層群が分布している。ビリミアン超層群 (2.2-2.1Ga) は広く緑色岩体が分布し、層厚が1000mを超える火山砕屑岩を主とする地層が分布する。本地域の海岸線には、連続性のよい深海底堆積物が露出する。GHB掘削によりビリミアン超層群セフィ層群の最上部の地層について試料を取得した。GHBコアはエジル湾岸 (Latitude: 4°45' 23" N, Longitude: 2°02' 15" E) から掘削され、コアの全長は195mであり、深さ30mまでは風化が著しいが、以深では連続コアを取得できた。コアは帯磁率測定とCTによる内部構造の測定を行った後、半割して岩相の記載を行った。これらの試料について薄片を作成し鏡下観察を行った。

GHBコアは変形や褶曲がなく最下部から最上部まで連続した地層からなる。風化の著しい上部30mを除き岩相・層序の特徴により下位から4つのUnitに分けた。Unit 1 (Depth:30-60m) はシルト質砂岩、黒色頁岩、緑色火山砕屑岩の薄層を繰り返す。火山砕屑岩は細粒砂岩サイズの角閃石や斜長石、石英が多く含まれる。地層中には薄く凝灰岩層が挟まれる。Unit 2 (Depth:61-120m) は主に塊状緑色火山砕屑岩や、緑色-緑灰色の火山砕屑岩とシルト質砂岩の薄層の繰り返しから成り、貫入岩が厚さ約20mに渡ってこれを貫く。Unit 1と比較して火山砕屑岩の鉱物組成に大きな変化はないが、層厚の観点では、Unit 1で見られたcm~mmスケールの薄層に加え、十数cm~数十cmにわたり緑色火山砕屑岩が厚く堆積した地層が何度も出現する特徴がある。Unit 3 (Depth:120-174m) はシルト質砂岩、頁岩の薄層の間にしばしば黒色頁岩層を挟む。炭質部は鏡下スケールでも観察でき、緑色-緑灰色火山砕屑岩のシルトサイズの粒子がなすmmスケールのラミネーションの間にも挟在する。Unit 4 (Depth:175-195m) は緑色火山砕屑岩と頁岩、黒色頁岩層から構成され、非常に細かい粒子から成るcm~mmスケール、あるいは1mm以下の厚さの薄層を繰り返す。黒色頁岩は不透明黒色物層を多く含み、その中には反射顕微鏡薄片観察により不透明鉱物 (黄鉄鉱) が観察された。黄鉄鉱と炭質部の分布は不均一である。黄鉄鉱は部分的に炭質な縞に濃集して観察され、それらの縞の上下の緑色火山砕屑岩からは観察されない。

以上、GHBコアは風化の著しい上部30mを除き下部から上部まで約165mの連続的な地層が観察される。コ

アは全般に珪質な緑色火山碎屑岩から構成され、鏡下観察でカーボネートがみられ、mmスケール以下のラミネーションまで非常に良く保存されている。Unit1からUnit2では緑色火山碎屑岩が主構成物であり、塊状緑色火山碎屑岩層が繰り返すことから、近隣の火山からの火山性碎屑物の供給が増加したと考えられる。Unit3からUnit4では頁岩やシルト質砂岩の頻度が増加し、上方へ向けて構成粒子が細粒化する。

キーワード：Ghana、ビルミアン帯、初期原生代、コア、層序

Keywords: Ghana, Birimian, early Proterozoic, Drilled core, Lithologies

Stratigraphy, Lithology and Geochemistry of 1.9Ga ocean floor sediments in the Flin Flon and Cape Smith Belts, Canada

*元村 健人¹、清川 昌一¹、池原 実²

*Motomura Kento¹, Shoichi Kiyokawa¹, Minoru Ikehara²

1. 九州大学、2. 高知大学

1. Kyushu University, 2. Kochi University

The Flin Flon and Cape Smith belts in the Trans-Hudson Orogen, Canada, include volcanics, volcanoclastics and alternation of sandstone and black shale, which accumulated about 1.9 Ga. Black shale preserve organic matter, and is a clue to study about Paleoproterozoic sea environment. Therefore, we did 1)describing detail of lithology, 2)forming geological column, 3) reconstruction of stratigraphy and 3)measurement of chemical composition and organic carbon isotope ratio ($\delta^{13}\text{C}_{\text{org}}$) of black shale about drilling core TS07-01(Embury Lake, the Flin Flon belt) and drilling core 718.3333 (Povungnituk Group, the Cape Smith belt) to reveal marine biology of 1.9 Ga. These cores are mostly composed by alternation of sandstone and black shale.

TS07-01 from the Flin Flon belt is 480m at total length, and has alternation of sandstone and black shale (470m) and intrusion of rhyolite (10m). A 300m scale asymmetrical fold with several minor parasitic folds are identified due to describing the detail of the core. Reconstructed stratigraphy sequence is 280m long. Sandstone contains quartz and plagioclase and black shale is composed by clay mineral, silt size quartz and pyrite. From top to bottom of the core, major elements composition of black shale are constant and similar to that of Post-Archean Australian average shale (PAAS). Organic carbon contents (TOC) is 2.1 wt% on average, and $\delta^{13}\text{C}_{\text{org}}$ is about -37‰ to -26‰.

718.3333 from the Cape Smith belt is 60m in total and composed by continuous stratigraphy of alternation of sandstone and black shale. Sandstone mainly contains fine quartz, and black shale include clay mineral and some silt size quartz and pyrite. Chemical composition is characterized by enrichment of S and organic carbon, and lack of Fe and P. TOC is 3.5wt% on average. $\delta^{13}\text{C}_{\text{org}}$ is from -38‰ to -33‰. Two reconstructed stratigraphic sequences of Trans-Hudson Orogen and its chemical features indicate the depositional area and environment did not change because mineral and chemical composition has no variety. $\delta^{13}\text{C}_{\text{org}}$ show cyanobacteria, chloromatiaceae and chloroblaceae lived 1.9 Ga ocean, and methanogenic bacteria may acted in sediment.

中国三峡地域エディアカラ系における銅元素化学層序

Copper chemostratigraphy of the Ediacaran in the Three Gorges area, South China

*耿野川¹、大野剛²、青木翔吾¹、坂田周平²、石川晃¹、小宮剛¹

*Yechuan Geng¹, Takeshi Ohno², Shogo Aoki¹, Shuhei Sakata², Akira Ishikawa¹, Tsuyoshi Komiya¹

1. 東京大学大学院総合文化研究科広域科学専攻、2. 学習院大学大学院自然科学研究科化学専攻

1. Department of Earth Science & Astronomy Graduate School of Arts and Sciences The University of Tokyo, 2. Department of Chemistry at Gakushuin University

The Ediacaran-Cambrian transition is characterized by several events such as emergence of metazoans and sudden changes of surface environment. Especially, the appearance and evolution of the metazoans in the Ediacaran are one of the most important issues of the evolution of life. Some candidates of causes of the drastic evolution were proposed, for example increase of oxygen contents of atmosphere and ocean, suppression and subsequent liberation of biological activity by the Snowball Earth event, and increase of marine nutrient contents, but they cannot fully account for the dramatic surface environmental change and biological evolution so that the cause(s) is still enigmatic. We focus our attention on the increase of marine nutrient contents because it also caused redox change of the atmosphere and ocean.

Copper is one of bioessential elements, which shows a vertical recycled- and scavenged-type profile in the modern ocean. The copper is incorporated into various proteins but hemocyanin is one of the most famous Cu-bearing proteins. The hemocyanin has a function, which transports oxygen throughout bodies of some invertebrate animals such as arthropods and molluscs. We study chemostratigraphies of copper contents and isotope ratios from the Ediacaran to the Cambrian in order to elucidate a geobiological cycle of copper through the Ediacaran. The copper chemostratigraphy will enable us to obtain new insights into the evolution of the life and environment. But, the stable isotope geochemistry of Cu is poorly known because of the lack of a suitable analytical technique.

We carried out on-land drilling of the sedimentary succession in Three Gorges area, South China to collect more continuous and fresher samples through the Ediacaran. In order to analyze the copper contents and isotope ratios, powdery samples were prepared from fresh-cut surfaces of the drill cores using a micro-drill with a 3 mm-diameter bit. The sampling points were carefully selected to avoid acquisition of late-stage diagenetic carbonate minerals and veins. The powdered samples were dissolved with HF-HNO₃-HClO₄ acid digestion, and a split of the sample solution was diluted with 2% HNO₃. We analyzed the Cu, Zn, Ni and Ti contents with ICP-MS, housed at Gakushuin University.

And the next step, we are going to analyze the isotope data of all the samples from Doushantuo formation. Each sample was subsequently dissolved in 1 ml of 10 N HCl and insoluble particle was centrifuged out. The separation of transition elements on strongly basic anion exchange resins in hydrochloric media follows a connective procedure (Kraus and Moore, 1953). Van der Walt et al. (1985) demonstrated that the macroporous form (AG MP-1) of strongly basic anion exchange resins has higher distribution coefficients for Cu(II), Fe(III) and Zn(II) in concentrated HCl. We will measure the Cu isotope compositions on a MC-ICP-MS.

We have got the concentration data from Doushantuo Formation's Member 1 to Member 4. The concentration of Cu increased in ca.580Ma, when the Gaskiers Glaciation was determined. And decreased in ca.550Ma, when is in the middle of Shuram Excursion. And then at the end of Doushantuo Formation, the concentration of Cu increased sharply. About the variation of Ni and Zn, we found that

there is a same pattern with Cu. Because the samples were also analyzed for the concentration of iron and manganese, and iron isotope values of pyrites (Sawaki et al, 2016), we directly compared between the concentration of Fe, Mn and Cu. We suggested that in oxic marine environment, the concentration of Cu had been increased, and in anoxic marine environment, it had been decreased. Because that molluscs appeared from ca.550Ma, and the arthropods which belong to the Ediacara Fauna appeared from ca.580Ma, we consider that the concentration variation of Cu in seawater, have important correlation with the appearance and evolution of the metazoans in Ediacaran.

キーワード：銅元素、エディアカラ紀、濃度、同位体、海水

Keywords: Copper, Ediacaran, concentration, Isotope, seawater

Estimation of Depositional Environment of Ediacaran Carbonates by Trace Element Abundances

*宮田 桃子¹、高橋 嘉夫¹、渡辺 勇輔¹、古山 精史朗²、狩野 彰宏¹

*Momoko Miyata¹, Yoshio Takahashi¹, Yusuke Watanabe¹, Seishiro Furuyama², Akihiro Kano¹

1. 東京大学大学院理学系研究科地球惑星科学専攻、2. 産業技術総合研究所

1. Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo, 2. Geological Survey of Japan

Introduction

In later Neoproterozoic, at least 3 times glacial event occurred including Marinoan and Gaskiers glacial periods (Hoffman and Li, 2009). Diamicite deposited during the Marinoan glacial event was covered by cap carbonate, which indicates that the sudden change of temperature occurred (Hoffman and Li, 2009). It is thought that by this change living species were selected, which eventually leads to the Cambrian explosion (Hoffman et al., 1998). Hence, it is thought that these cap carbonates give an important hint about the most important environmental change after glacial period for primary animal diversification in the earth history.

Trace elements in deposited sediments can be a proxy of the depositional environment. REE abundances in carbonate rocks reflect those in seawater (Tanaka et al., 2006). Parameters in the REE pattern can be useful to interpret the depositional environment in each sample: (i) Ce anomaly can be proxy of paleoceanographical redox state (Wang et al., 2014); (ii) Y/Ho and tetrad effect include information of water mass in the ocean; (iii) slope of REE pattern (= Pr/Er) reflects water depth in the ocean, if we can assume that variation of REE pattern can be similar in the late Neoproterozoic ocean.

In this study, these REE concentration were compared with carbon isotope ratio that was measured at Kunimitsu et al. (2011). Kunimitsu et al. (2011) divided the section into four units based on the carbon isotope variation. Among them, Unit 1 and 3 correspond to the period after the Marinoan glacial event and cooling period during Gaskiers glacial event, respectively.

Samples and Methods

In this study, Carbonates collected at Yanjiaping in Hunan Province, China were used as analysis samples (Kunimitsu et al., 2011). This section covers the interval from late Cryogenian to earliest Cambrian (655-542Ma; Amthor et al., 2003).

Trace element concentrations in carbonate were measured by Inductively Coupled Plasma - Mass Spectrometry (ICP-MS). Samples were crashed by hummer and grinded by agate mortar. Carbonates were dissolved into acetic acid and diluted to 10 or 1000 times. In and Bi were used as internal standard for the measurement by ICP-MS.

Results and discussion

REE patterns are measured by ICP-MS and normalized by using Post-Archaean Australian Shale (PS). Ce and Eu anomaly was calculated by using PAAS as below, whereas the degree of tetrad effect level was reflected by the Dy/Dy* ratio. Variation of trace elements were also examined.

Carbonate samples used in this study is same as the carbonate samples analyzed in Kunimitsu et al. (2012) and Furuyama et al. (2012) and discussed by segmenting from Unit 1 to 4 much as Kunimitsu et al. (2012).

It was clear that the degree of Ce anomaly is correlated with Y/Ho ratio. Although Ce anomaly is primary redox indicator, but the degree of negative Ce anomaly increases with increase of the water depth in the

ocean, but almost no Ce anomaly can be expected in river water. Y/Ho ratio, which is originally sensitive to ionic-covalent character of reactions, increases with the increase of the ratio. As a result, there is a clear relationship between the two parameters with plots of modern seawater, river water, and coastal water. The close correlation showed the environmental change that occurred within the Yanjiaping section is the frequent shifts between coastal or estuarine environment and deeper marine environment. In particular, water mass from deep ocean must have been supplied when negative Ce anomaly and Y/Ho ratio became large. Such trends were clearly found in Unit 3, which can explain the supply of nutrients including phosphorous to form phosphate sediments within the Unit 3.

Similar discussion is valid when we plot Y/Ho ratio and Dy/Dy* ratios. In this plot, it was again clear that depositional environment changed greatly during the Unit 3.

Increase of the slope of the REE pattern toward heavy REE correlated with the increase of the Y/Ho also shows the supply of deep sea water mass was supplied to the shallow depositional environment in the Unit 3.

Relatively intense Ce anomaly in 0-20 cm from the boundary with Nantuo formation in Unit 1 is notable, when black shale sediments were predominant, which suggests reductive environment. The positive Ce anomaly in carbonite, or seawater is very unique signature, which cannot be observed in normal oxic environment. Concentration of Mn in the carbonate was also high during the period. These results suggested that ferromanganese nodule which generally have positive Ce anomaly and can be formed right after the end of snowball earth were dissolved into water by the reducing environment in the period.

Euendoliths versus Ambient Inclusion Trails from Early Cambrian Kuanchuanpu Formation, South China

*Xiaoguang Yang^{1,2}, Jian Han², Xing Wang², James Schiffbauer³, Osamu Sasaki⁴, Tsuyoshi Komiya¹

1. Department of Earth Science & Astronomy Graduate School of Arts and Sciences The University of Tokyo, Komaba, 2. Department of Geology and State Key Laboratory of Continental Dynamics, Northwest University, 3. Department of Geological Sciences, University of Missouri, 4. Tohoku University Museum, Tohoku University

Abundant microstructures have been discovered in small skeletal fossils (SSFs) and embryo-like fossils collected from the Lower Cambrian Kuanchuanpu Formation (ca. 535 Ma) in Xixiang County, Shaanxi Province, China. These involve two co-occurring structures: long, unbranched cylindrical filaments, which are comparable with phosphatic casts of microborings constructed by euendolithic cyanobacteria *Endoconchia lata* in morphology and preservation pathway; and meandering micro-tubes or grooves on fossil moulds (and steinkerns) of a wide range of sizes and morphological diversities, discerned as ambient inclusion trails (AITs). Herein, we also report a new type of AIT propelled grains as organic carbon spherules and their implications on morphological diversity of AITs. From the direct comparisons of endolithic traces and AITs, we propose a mechanism to account for their notably different preservation, and further attempt to offer an explanation for their co-occurrence. Their differential preservation suggests a chronological sequence of their formation, such that *E. lata* microborings formed prior to phosphate sedimentation, while the AITs are likely generated in a later phase of (or after) phosphorite precipitation but before calcareous re-cementation. In that sense, we have identified the diagenetic stage of AITs forming in phosphatized fossils. Dissecting the sequence of these structures and detailed morphological observations assists in distinguishing true biologically produced endoliths from otherwise abiogenically produced microstructures.

Keywords: Cambrian, Kuanchuanpu, microfossils, micro-CT, Cambrian Explosion, Biological evolution

Lithological and geochemical features of the Permian-Triassic boundary at the Gujo-Hachiman section in the Mino-Tamba belt, central Japan

*矢野 萌生¹、藤永 公一郎^{2,1}、安川 和孝^{1,2}、野崎 達生^{3,1,4}、黒田 潤一郎⁵、桑原 希世子⁶、加藤 泰浩^{1,2,3}

*Moei Yano¹, Koichiro Fujinaga^{2,1}, Kazutaka Yasukawa^{1,2}, Tatsuo Nozaki^{3,1,4}, Junichiro Kuroda⁵, Kiyoko Kuwahara⁶, Yasuhiro Kato^{1,2,3}

1. 東京大学・工学系研究科、2. 千葉工業大学・次世代海洋資源研究センター、3. 海洋研究開発機構、4. 神戸大学・理学研究科、5. 東京大学・大気海洋研究所、6. 芦屋大学・臨床教育学部

1. School of Engineering, The University of Tokyo, 2. Ocean Resources Research Center for Next Generation, Chiba Institute of Technology, 3. JAMSTEC, 4. Graduate School of Science, Kobe University, 5. Atmosphere and Ocean Research Institute, The University of Tokyo, 6. Ashiya University

The most significant mass extinction event in the Phanerozoic era occurred across the Permian-Triassic boundary (PTB; ca. 252 Ma). However, the entire picture of this global environmental crisis has not reached a consensus, as well as a culprit that invoked the catastrophe. The PTB sections in the Japanese accretionary complex were primarily deposited in the deep-sea of the superocean Panthalassa, hence it is very likely to have recorded signatures of the global-scale environmental change. Therefore, a number of geological, geochemical, and paleontological researches have targeted them to deduce a series of environmental shift during the event (Isozaki, 1997; Kato et al., 2002; Takahashi et al., 2009). The Gujo-Hachiman section located in the Mino-Tamba belt, central Japan, has a continuous outcrop across the PTB (Kuwahara et al., 1998; Yao et al., 2001). In the present study, to decipher the marine environmental change across the PTB, we report the lithological and geochemical features on the basis of thin-section descriptions and chemical analyses of total organic carbon and bulk chemical compositions of the Gujo-Hachiman section.

キーワード：ペルム紀/トリアス紀境界、黒色頁岩、チャート、郡上八幡、海洋無酸素化

Keywords: Permian-Triassic boundary, black shale, chert, Gujo-Hachiman, ocean anoxia

タイ東北部ファイオムセクションにおける79万年前の東南アジア衝突イベント起源の衝撃変成石英の発見

Discovery of Shocked Quartz in a Possible Ejecta Deposit of 0.79 Ma Impact at Huai Om Section, NE Thailand

*多田 賢弘¹、多田 隆治¹、Carling Paul²、Songtham Wickanet³、常 昱¹、田近 英一¹

*Toshihiro Tada¹, Ryuji Tada¹, Paul Carling², Wickanet Songtham³, Yu Chang¹, Eiichi Tajika¹

1. 東京大学大学院理学系研究科地球惑星科学専攻、2. Geography & Environment, University of Southampton、3. Northeastern Research Institute of Petrified Wood and Mineral Resources, Nakhon Ratchasima Rajabhat University
1. Department of Earth and Planetary Science, 2. Geography & Environment, University of Southampton, 3. Northeastern Research Institute of Petrified Wood and Mineral Resources, Nakhon Ratchasima Rajabhat University

天体衝突は地球環境に大きな影響を及ぼすことがあるものの、その詳細は不明な点が多い。オーストラリア・アジアテクトイト分布域は、これまでに報告されている4つのテクトイト分布域の内、東南アジア～南極にわたる最も広域かつ最も年代が若い(0.79Ma)という特徴を持ち、タイ東北部の大規模洪水層やジャワ島サンギランの地層における原人の最終出現との関連が示唆されている(Haines et al., 2004; Hyodo et al., 2011)。

その衝突地点は、陸上のテクトイトや海底堆積物コア中のマイクロテクトイトの分布からインドシナ半島東部と推定されているものの、衝突クレーターは未発見であり、正確な衝突地点や衝突の規模、様式は明らかにされていない(Glass and Koeberl, 2006など)。イジェクタ層はクレーターに近いほど層厚が厚いため、より正確な衝突地点の推定にはイジェクタ層の分布を明らかにすることが重要である。しかし、イジェクタ層は海底堆積物コアのみで同定され、より衝突地点に近いと考えられるインドシナ半島では同定されていない。

本研究は、インドシナ半島でこの衝突のイジェクタ層を同定することを目的とし、タイ東北部において野外調査を行った。タイ東北部ウボン・ラーチャターニー県ファイオムセクションは、Fiske et al. (1996)によって記載され、堆積物中から層状テクトイトが報告されている。本研究では、独自に岩相・層序の記載を行い、シルト基質中に砂岩角礫が含まれることを確認した。また石英粒子を偏光顕微鏡下で観察し、ラメラを持つ石英を見出した。ラメラの方位を、ユニバーサルステージを用いて測定した結果、衝撃変成石英に特徴的な方位が同定されたことから、これらの石英粒子は衝撃変成石英であると考えられる。この衝突に関して陸上の地層中から衝撃変成石英を見出したのは、本研究が初めてである。このことから、Huai Om section にみられる堆積物は、イジェクタ層である可能性が高い。

北西太平洋深海堆積物コアの全岩化学組成に基づくレアース泥起源の解明

Origin of REY-rich mud based on bulk geochemistry of deep-sea sediment cores in the western North Pacific Ocean

*山本 克志¹、見邨 和英¹、中村 謙太郎¹、安川 和孝^{1,2}、大田 隼一郎^{3,1}、藤永 公一郎^{2,1}、町田 嗣樹^{3,1}、加藤 泰浩^{1,2,3}

*Katsushi Yamamoto¹, Kazuhide Mimura¹, Kentaro Nakamura¹, Kazutaka Yasukawa^{1,2}, Junichiro Ohta^{3,1}, Koichiro Fujinaga^{2,1}, Shiki Machida^{3,1}, Yasuhiro Kato^{1,2,3}

1. 東京大学大学院工学系研究科システム創成学専攻、2. 千葉工業大学・次世代海洋資源研究センター、3. 海洋研究開発機構

1. Department of Systems Innovation, School of Engineering, the University of Tokyo, 2. ORCeNG, Chiba Institute of Technology, 3. JAMSTEC

REY-rich mud, a deep-sea sediment containing high concentrations of rare-earth elements and yttrium (REY), is expected to be a new resource for the critical elements due to its multiple advantages such as huge resource potential and paucity of radioactive elements [1]. It has been confirmed that the REY-rich mud also exists in the Japanese exclusive economic zone (EEZ) around Minamitorishima Island [2], followed by the discovery of “extremely REY-rich mud” that contains more than 5,000 ppm of total REY from the Minamitorishima EEZ [3].

For the future development of the REY-rich mud, clarifying the stratigraphy and lateral extent of the extremely REY-rich mud is critically needed. In this respect, continuous cores of deep-sea sediments obtained by deep-sea drilling can provide important information. In the western North Pacific Ocean, only two drilling cores, Ocean Drilling Program (ODP) Sites 1149 and 1179, were almost continuously recovered from the seafloor surface to basement rock (chert). Chemical analyses of these two cores could provide the entire picture of common stratigraphy of deep-sea sediments including the extremely REY-rich mud in the western North Pacific Ocean. The analysis of the Site 1149 has already been implemented by Mimura (2016) [4], which demonstrated that the site has essentially common stratigraphy with the Minamitorishima EEZ. In the present study, we newly analyzed sediment samples of the Site 1179 and determined their bulk chemical compositions.

The sediments in the Site 1179 has been classified into five units: clay- and radiolarian-bearing diatom ooze of Unit I (from the core top to 221.52 mbsf), clay-rich and diatom-bearing radiolarian ooze of Unit II (221.52 to 246.0 mbsf), pelagic clay of Unit III (246.0 to 283.53 mbsf), chert layer of Unit IV (283.53 to 377.15 mbsf) and basaltic crust layer of Unit V [5]. Analytical results of 173 bulk sediment samples from Unit I, Unit II, and Unit III, show that the REY-rich mud occurs only in the Unit III lower than ~260 mbsf. Moreover, whereas the extremely REY-rich mud containing 7,500 ppm of total REY has been discovered at Site 1149, the total REY content at Site 1179 was at most 1,675 ppm. In addition, we clarified the geochemical end-members constituting the deep-sea sediments in the Site 1179: terrigenous, hydrogenous, hydrothermal, and biogenic calcium phosphate components. Based on the results, we further quantified the relative contributions of each component. By comparing the new data from the Site 1179 with previous data from the Site 1149, we found out both a common chemostratigraphy of deep-sea sediments in the western North Pacific Ocean as well as local characteristics overlapping the common features.

References

- [1] Kato, Y. et al. (2011) *Nature Geoscience* **4**, 535-539.
- [2] Kato, Y. et al. (2012) *Abstract with programs, the society of Resource Geology*.
- [3] Iijima, K. et al. (2016) *Geochemical Journal* **50**, 557-573.
- [4] Mimura, K. et al. (2016) *JpGU Meeting 2016*.
- [5] Kanazawa, T. et al. (2001) *Proceedings of Ocean Drilling Program, Initial Reports* **191**, 4. Site 1179, 1-159.

キーワード：レアアース泥、海底鉱物資源、全岩化学組成、ODP Site 1179、北西太平洋

Keywords: REY-rich mud, deep-sea mineral resources, bulk geochemistry, ODP Site 1179, western North Pacific Ocean