

## Integrating Statistical and Expert Knowledge to Develop Phenoregions for the Continental United States


\*Forrest M. Hoffman<sup>1,2</sup>, Jitendra Kumar<sup>1,3</sup>, William W. Hargrove<sup>4</sup>, Steven P. Norman<sup>4</sup>, Bjørn-Gustaf J. Brooks<sup>4</sup>

1. Climate Change Science Institute (CCSI), Oak Ridge National Laboratory (ORNL), 2. Department of Civil & Environmental Engineering, University of Tennessee, 3. The Bredesen Center for Interdisciplinary Research, University of Tennessee, 4. Eastern Forest Environmental Threat Assessment Center (EFETAC), United States Department of Agriculture (USDA) Forest Service

Vegetated ecosystems exhibit unique phenological behavior over the course of a year, suggesting that remotely sensed land surface phenology may be useful for characterizing land cover and ecoregions. However, phenology is also strongly influenced by temperature and water stress; insect, fire, and weather disturbances; and climate change over seasonal, interannual, decadal and longer time scales. Normalized difference vegetation index (NDVI), a remotely sensed measure of greenness, provides a useful proxy for land surface phenology. We used NDVI for the conterminous United States (CONUS) derived from the Moderate Resolution Spectroradiometer (MODIS) every eight days at 250 m resolution for the period 2000–2015 to develop phenological signatures of emergent ecological regimes called phenoregions. We employed a “Big Data” classification approach on a supercomputer, specifically applying an unsupervised data mining technique, to this large collection of NDVI measurements to develop annual maps of phenoregions. This technique produces a prescribed number of prototypical phenological states to which every location belongs in any year. To reduce the impact of short-term disturbances, we derived a single map of the mode of annual phenological states for the CONUS, assigning each map cell to the state with the largest integrated NDVI in cases where multiple states tie for the highest frequency of occurrence. Since the data mining technique is unsupervised, individual phenoregions are not associated with an ecologically understandable label. To add automated supervision to the process, we applied the method of Mapcurves, developed by Hargrove and Hoffman, to associate individual phenoregions with labeled polygons in expert-derived maps of biomes, land cover, and ecoregions. We will present the phenoregions methodology and resulting maps for the CONUS, describe the “label-stealing” technique for ascribing biome characteristics to phenoregions, and introduce a new polar plotting scheme for processing NDVI data by localized seasonality.

Figure: This map shows the 50 phenoregions derived from the MODIS NDVI at 250 m resolution data for years 2000–2012. The phenoregions are colored using a “similarity colors” technique that employs a principle components analysis to produce data-specific combinations of red, blue, and green for every phenoregion in the map.

Keywords: phenology, NDVI, MODIS, phenoregions, label stealing, cluster analysis



## インターネット上で公開されているシチズンサイエンスやアーカイブデータを利用した植物季節観測の高精度化

### Development of plant phenological observation by using citizen science and historical archived data published on the web sites

\*永井 信<sup>1</sup>、齋藤 琢<sup>2</sup>

\*Shin Nagai<sup>1</sup>, Taku Saitoh<sup>2</sup>

1. 海洋研究開発機構地球表層物質循環研究分野、2. 岐阜大学流域圏科学研究センター

1. Department of Environmental Geochemical Cycle Research, Japan Agency for Marine-Earth Science and Technology, 2. River Basin Research Centre, Gifu University

Accurate detection of plant phenology (e.g. timing of flowering, leaf-flush, and leaf-fall) is required to evaluate the spatio-temporal variability of ecosystem functions and service under rapid climate changes. Towards this aim, analysis of daily satellite-observed vegetation index with a coarse spatial resolution (e.g. 500m) and in situ-observed long-term historical biometeorological data set is useful. However, these approaches include many uncertainties and problems mainly caused by heterogeneity of plant species, spatial representativeness, and land cover changes. Here, (1) we examined the relationship between leaf-colouring information published on the meteorology service web site (<http://www.tenki.jp>) and the timing of end of growing season detected by daily Terra and Aqua/MODIS satellite-observed green-red vegetation index in Japan; (2) we evaluated the long-term historical flowering information published on the web sites; and (3) we examined the land cover change in the “Satoyama” landscape area by using aerial photographs published on the geographical survey web site (<http://mapps.gsi.go.jp/maplibSearch.do#1>). In this presentation, we will discuss the usability of citizen science and historical archived data published on the web sites for developing the detection of spatio-temporal variability of plant phenology in Japan.

キーワード：植物季節、リモートセンシング、シチズンサイエンス、ウェブサイト

Keywords: phenology, remote-sensing, citizen science, web site

## Land Surface Phenology Changes in Central Asia

\*Kirsten de Beurs<sup>1</sup>, Geoffrey M Henebry<sup>2</sup>, Braden C Owsley<sup>1</sup>, Irina Sokolik<sup>3</sup>

1. Department of Geography and Environmental Sustainability, University of Oklahoma, 2. Geospatial Sciences Center of Excellence, South Dakota State University, 3. School of Earth and Atmospheric Sciences, Georgia Institute of Technology

Land surface phenology metrics allow for the summarization of long image time series into a set of annual observations that describe the vegetated growing season. These metrics have been shown to respond to both climatic and anthropogenic impacts. In this study we assembled a time series (2001-2016) of Moderate Resolution Imaging Spectroradiometer (MODIS) Nadir BRDF-Adjusted Reflectance (NBAR) data at two spatial resolutions (0.05° and 500m) and land surface temperature data at two spatial resolutions (0.05° and 1000m). We then derived land surface phenology metrics focusing on the peak of the growing season by fitting convex quadratic regression models connecting the NDVI time series with the progression of Accumulated Growing Degree-Days (AGDD) derived from the land surface temperature data. We linked the annual information on (1) peak timing, (2) thermal time to peak and (3) peak magnitude with three important climate oscillations—the Atlantic Multidecadal Oscillation (AMO); the North Atlantic Oscillation (NAO); and the East Atlantic / West Russia pattern (EAWR)—and evaluated the effects of the different spatial resolutions. We discovered several significant correlations between the climate oscillations and the land surface phenology peak metrics for a range of different bioclimatic regions in the drylands of Central Asia, and we linked these correlation results to changes in ambient population modeled by LandScan.

Keywords: Land Surface Phenology, Remote Sensing, Central Asia

## Tracking and characterizing human impacts in tropical forests: Can Landsat go it alone?

\*Ben DeVries<sup>1</sup>, Jan Verbesselt<sup>2</sup>, Johannes Reiche<sup>2</sup>, Arun Kumar Pratihast<sup>2</sup>, Martin Herold<sup>2</sup>

1. University of Maryland, College Park, MD, USA, 2. Wageningen University, The Netherlands

Representing a significant portion of the global carbon store as well as countless other ecosystem services, tropical forests have been under threat due to expansion of human activities in recent decades. Following the public release of the entire Landsat data archive - the longest record of earth observation data in existence - a veritable explosion of innovation in forest monitoring methods using Landsat time series (LTS) has occurred. New approaches have emerged allowing for the tracking of forest changes either retrospectively - supporting carbon accounting for REDD+ and other applications - or in near real-time - supporting operational monitoring and enforcement efforts. Methods exploiting all observations in particular have shown that accounting for seasonal fluctuations (ie. those arising from canopy phenology) can enhance the detection and characterization of human-driven changes. Despite the demonstrated promise of LTS in monitoring changes in tropical forests, limitations in the temporal, spectral and spatial resolution of Landsat data raise questions about extent to which methods based on LTS alone are adequate for addressing monitoring needs in tropical forests.

Here, we show that while LTS provide unprecedented detail in forest change studies, specific monitoring objectives in tropical forests call for the integration of LTS with other data sources. To demonstrate this need, we focus on two monitoring targets: (1) the timely detection of changes and (2) characterization of change dynamics over time. First, given that many tropical regions experience perpetual cloud-cover, generating large temporal gaps in LTS, recent research has shown that fusion of LTS with SAR data can improve the temporal accuracy of forest change alerts by up to 47 days. Fusion of LTS with the ESA-Copernicus Sentinel constellation of SAR and optical satellites thus has the potential to significantly improve the near real-time forest monitoring systems. Second, LTS-based methods face limitations when characterizing forest change processes like deforestation and forest degradation. Recent results integrating forest observations from community-based monitoring (CBM) project sites with LTS-based change indicators show that gradual, small-scale degradation of the forest canopy can be detected using LTS, but require regularly acquired *in situ* observations for adequate calibration and validation of change models. These insights show the promise of fusing LTS with other satellite data streams such as SAR image time series, as well as *in situ* observations and measurements, to enhance forest monitoring capabilities in the tropics to support such objectives as REDD+.

Keywords: Landsat, time series, tropical forests, SAR, deforestation, participatory monitoring

# Interannual Variability in the Length of Growing Season across the Contiguous US as Observed from MODIS, VIIRS, and Tower Vegetation Index Time Series Data

\*Anna Kato<sup>1</sup>, Tomoaki Miura<sup>1</sup>

1. Department of Natural Resources and Environmental Management, University of Hawaii at Manoa

Vegetation phenology is considered one key variable to understand climate change impacts on plant productivity. Spectral vegetation index (VI) time series data from polar-orbiting satellites such as Moderate Resolution Imaging Spectroradiometer (MODIS), have widely been used to characterize vegetation phenology at various spatial and temporal scales. They have shown to be valuable tools in discerning various phenological metrics, such as the start and end of growing seasons (SOS and EOS), and have recently been used to quantify interannual variability in the length of growing season (GSL) at regional and global scales. Assuring the quality of satellite-derived GSL is critical to characterize directions and magnitudes of shifts in GSL, which are linked to the changes in plant productivity. The objective of this study was to assess the quality of satellite-derived GSL in characterizing the relative impacts of the SOS and EOS changes to the GSL interannual variability by an intercomparison with in-situ optical sensor data. The GSL derived from MODIS (2003-2015) and Visible Infrared Imaging Radiometer Suite (VIIRS) (2012-2015) VI time series data were compared with those derived from in-situ (Tower) VI time series data (2003-2015), which were derived from in-situ tower optical sensor data, at 11 AmeriFlux sites located across a range of biomes in the contiguous US (CONUS). The SOS and EOS metrics were obtained from all the three datasets by fitting logistic function for each year, and the GSL was calculated as the difference between the SOS and EOS. Relative contributions of the changes in SOS and EOS to the GSL interannual variability were evaluated using the ratios of the absolute changes in SOS to that of GSL. Year-to-year changes in MODIS and VIIRS GSLs corresponded well with those from the Tower GSL although some biases were observed for some sites ( $R^2 > 0.68$ ,  $p < 0.01$ ). The relative contributions of the SOS and EOS varied across sites (SOS contribution = 20~93%), which were seen in all the three datasets. The stronger SOS contributions (>50%) were seen at many of the sites located in the mid-latitude region (37.5°N~42.5°N), whereas EOS showed stronger contributions (>50%) at those sites located in the southwestern part of the CONUS. These relative contributions changed temporally but the observed spatial patterns were consistent. These results indicate that the satellite-derived GSL is capable of characterizing the interannual variability of GSL in the CONUS. Further analysis at additional locations may help to assure the quality of satellite-derived GSL and to better understand GSL interannual variability under climate change.

Keywords: vegetation phenology, growing season length, MODIS, VIIRS, in-situ radiation data


## Impacts of changes in phenology on land-atmosphere interactions in temperate and boreal regions

\*Jorg D Kaduk<sup>1</sup>, Kirsten Barrett<sup>1</sup>, Andrea Hurtado De Mendoza Rosales<sup>1</sup>, Narumasa Tsutsumida<sup>2</sup>

1. Univ. of Leicester, Dept of Geography and Centre for Landscape and Climate Research, 2. Kyoto Univ., Graduate School of Global Environmental Studies

Recent remote sensing data and ground observations have shown earlier leaf out in spring in the northern hemisphere, which is believed to result from climate warming. The advance of leaf out would be expected to have impacts on the dynamics of the wider ecosystem processes such as primary production and evapotranspiration. Moreover, controlled experiments show that temperate and boreal trees require chilling in winter for rapid leaf out in spring. If the amount of chilling falls below a species specific threshold then an exponentially increasing amount of warming is required to initiate leaf out –potentially actually delaying it in a warmer climate. Implications of these chilling requirements for a delayed greening of vegetation at the biome level are not clear. Impacts of changes in the phenology in the past thirty years are explored by incorporating Leaf Area Index (LAI) data derived from satellite remote sensing observed Normalized Difference Vegetation Index (NDVI) into the Joint UK Land Environment Simulator (JULES), a numerical biosphere-atmosphere exchange simulation system. A 30 year model simulation using daily varying climate and monthly varying LAI is compared to a simulation with the varying climate but a fixed seasonal cycle. So, while the LAI varies between months in the second simulation, the LAI is the same for all 30 Januaries, for all 30 Februaries and so on. The first simulation shows the effects of varying climate and phenology over the last 30 years on the northern hemisphere, the second reflects only the climate variability. The analysis of the simulations shows that there are significant changes due to the changes in the phenology in the biosphere-atmosphere fluxes in some areas of the northern hemisphere. In particular the net primary productivity increases significantly for example in the South Eastern United States. Further changes in the biosphere-atmosphere fluxes are explored. The results highlight the necessity of including appropriate phenology models in climate models for improved predictions of land-atmosphere interactions.

Keywords: Plant Phenology, JULES, Biosphere-atmosphere exchange, Net Primary productivity






# Probing Polarization to Assess the Cuticular Condition of a Terrestrial Canopy

\*Ade Purwanto<sup>1</sup>

1. Hokkaido University

This study presents an empirical study of the polarized-light phenomena on a leaf surface under the laboratory condition, using a Liquid Crystal Tuneable Filter (LCTF) camera with a linear polarizer filter attached on front of its lens and were rotated to obtain the degree of polarization (DoP) of the target. The instrument was capable to capture the spectrum between 460nm to 780nm with the step of 10nm. The experiment was focused to capture the specular reflectance from the surface of the leaf, thus we set the angle between the light source and the sensor always in mirror angle. Several different leaves with distinct different feature on its surface were observed in a dark-room laboratory and outdoor measurement. Results show that a polarization measurement is capable to distinguish changes on the drought-stress plant, thus it remains a key component of the remote sensing probe to understand the vegetation cuticular condition, and in a broader range, terrestrial canopy.

Keywords: Degree of Polarization, Drought, Hyperspectral, Forest Canopy, Specular Reflection



# Long-term monitoring of land surface phenological changes

## Long-term monitoring of land surface phenological changes

\*堤田 成政<sup>1</sup>

\*Narumasa Tsutsumida<sup>1</sup>

1. Graduate School of Global Environmental Studies, Kyoto University

1. Graduate School of Global Environmental Studies, Kyoto University

Land surface phenology (LSP) is associated with climate over space and time, and the monitoring of LSP help understandings of the terrestrial environmental changes. The LSP is often inferred by satellite observation, and long-term and regularly composite satellite imagery is now freely available. In this study, we demonstrate how LSP changes over space and time at the global scale over the last three decades by using GIMMS3g datasets. We focus on the magnitude and the timing of the peak of yearly phenological activity, estimated from a harmonic analysis. The first harmonic curve is regarded as a proxy of the overall productivity of vegetation and the second one is interpreted as a sensitive bimodal system changes. Results show the long-term trend of LSP changes; for example the peak of phenological activity tend to be earlier in high-latitude regions.

キーワード : Phenology、 Remote sensing、 Harmonic analysis

Keywords: Phenology, Remote sensing, Harmonic analysis

## Spatial correlation for phenological responses to climate

\*Joseph Percival<sup>1</sup>, Narumasa Tsutsumida<sup>1</sup>

1. Graduate School of Global Environmental Studies, Kyoto University

Phenological and climate data are highly variable across time and space, and have been shown to display both spatial and temporal non-stationarity. In time series analysis, such variability in the time domain may be described in one variable by autocorrelation, or between two or more variables by cross-correlation and cross-partial correlation expressions using time lags. In spatial analysis, this variability may be captured by creating local models using approaches such as geographically weighted modeling techniques that utilize spatially weighted kernel functions. Although many previous studies regarding phenology focus on how phenological events respond to climate temporally, it is apparent that the spatial relations which may also influence such events have not yet been thoroughly considered. Hence this study demonstrates how spatial and temporal effects may be included in statistical analyses of phenological responses to climate data by including geographically weighted modeling techniques in time series analysis of NDVI values, temperature, and precipitation. This study uses monthly climate and phenological data from the CRUTS and GIMMS3g time series data sets over the last three decades. Cross correlation and cross partial correlation between the three variables are observed to find time lags that display higher positive and negative correlation coefficients; following this, geographically weighted correlation and partial correlation techniques are applied to the lagged data. It is expected that this will allow for a more careful examination of the relationship between temperature, precipitation, and NDVI.

Keywords: phenology, geographically weighted approach, time series, spatial relation

## Afforestation monitoring using long-term remotely sensed data in Chinese semi-arid area

\*原 裕太<sup>1</sup>、堤田 成政<sup>1</sup>、西前 出<sup>1</sup>

\*Yuta Hara<sup>1</sup>, Narumasa Tsutsumida<sup>1</sup>, Izuru Saizen<sup>1</sup>

1. 京都大学大学院地球環境学堂・学舎

1. Graduate School of Global Environmental Studies, Kyoto University

The Conversion of Cropland to Forest Program (CCFP) has been implemented since 1999 to prevent the land degradation in the Loess Plateau, China. Many croplands in this region has been converted into plantation sites according to the CCFP, however the situation (such as growth, stability, or adaptation) of planted trees is not well monitored continuously over space. Satellite sensors have observed terrestrial surface since 1980s which fully covers the pre/post implementation period of the CCFP and available multi-temporal satellite images enable us to analyze vegetation response through the trajectory of Vegetation Index (VI). Therefore we attempt to evaluate the afforestation program by using VI time-series of LANDSAT and MODIS over the long term. The occurrence of afforestation, timing of planting, and temporal greenness trends are analyzed by Breaks For Additive Season and Trend method. As a case study, we choose Wuqi County in the northwest Shaanxi Province, located in the central Loess Plateau. This county is on the semi-arid transit zone between forest and grasslands, and has been designated as the national pilot model county for CCFP. Results showed that VI time-series could detect and measure the plantation forest growth spatially as a consequence of CCFP. It can be said that this study explored the new possibility for understanding the positive/negative effects of CCFP and evaluating how the afforestation by CCFP worked against land degradation.

キーワード：リモートセンシング、時系列植生指数、植林、半乾燥地、黄土高原

Keywords: Remote sensing, Vegetation-Index time series, Afforestation, Semi-arid areas, Chinese Loess Plateau

## Impact of spatial scale for phenological indices derived from remotely sensed data

\*堤田 成政<sup>1</sup>、Kaduk Joerg<sup>2</sup>

\*Narumasa Tsutsumida<sup>1</sup>, Joerg Kaduk<sup>2</sup>

1. Graduate School of Global Environmental Studies, Kyoto University, 2. Department of Geography, University of Leicester

1. Graduate School of Global Environmental Studies, Kyoto University, 2. Department of Geography, University of Leicester

Land surface phenology (LSP) characterizes the vegetated land surface and is practical to understand terrestrial environments at a global scale. Regularly observed remotely sensed data such as Landsat, MODIS, and AVHRR contributes to analyze LSP spatially. However, at least two main challenges should be addressed such that (i) the spatial resolution which attributes to the data source may significantly impact to LSP estimation, and (ii) the estimated LSP may not represent the vegetated land surface well due to the mixed land cover. Previous studies have shown that the estimation of LSP from different data is not consistent due to the spatial scale of data but yet fully linked with the mixed land cover problem. Thus, in this study, we attempt to analyze the impact of spatial scale issue to the estimated LSP in homogenous land cover areas. We use freely available remotely sensed data with different spatial resolution such as Landsat (30m), MODIS (250m, 500m, 1km), and GIMMS3g (8km) and estimate phenological indices for each. As land cover description differs among data products, land cover classes are aggregated into 12 classes globally from major global land cover products (GLCC, GLC2000, and globcover), then spatially homogenous land cover are only picked up. Phenological indices such as the magnitude and the peak of DOY are calculated by harmonic analysis to compare results among different spatial scales. The variability of phenological indices is explored according to the different spatial scale under the condition of homogenous land cover. It is expected to model such variability to overcome the spatial scale impact and such characteristics depending on the spatial scale should be taken into account when considering LSP from satellite.

キーワード : Spatial scale, Land surface phenology, remotely sensed data

Keywords: Spatial scale, Land surface phenology, remotely sensed data

## Remote sensing of reflectance properties related to ecological shifts post-fire in southern Siberia

\*Kirsten M Barrett<sup>1</sup>, Elena Kukavskaya<sup>2</sup>, Joerg Kaduk<sup>1</sup>

1. University of Leicester, 2. V.N. Sukachev Institute of Forest

Rapid ecological shifts such as the replacement of forest ecosystems by grasslands can be detected using remotely sensed reflectance data. In addition to detecting forest losses, a longer time series of reflectance data can provide information about the persistence of these shifts, or alternatively about ecosystem recovery. In addition to changes in the amount of reflected energy, ecological shifts can be detected from the timing of phenological events that may change as the result of the replacement of one plant functional type with another. Our ability to detect forest recovery from disturbance depends on the length of the remotely sensed data record in relation to the recovery signal. Monitoring ecosystem recovery often relies on some proxy for a fully recovered ecosystem, usually the signal from the period prior to disturbance or from an adjacent, undisturbed area. The requirement of a pre-disturbance signal means an even longer time series of data are needed, and finding nearby undisturbed areas for comparison can be unreliable and time-consuming. The pattern of reflectance time series data is also likely vary depending on the spatial and temporal resolution of the data used to analyse them. In the southern boreal forests of Siberia, recovery of reflectance properties post-disturbance may take several decades or more, or it may appear suddenly after a prolonged period without a discernible trend. We analyse several areas of known forest loss post-fire as well as some normally recovering sites to determine typical trajectories of recovery in reflectance properties from several remotely sensed datasets.

Keywords: boreal, wildfire, disturbance, recovery, remote sensing, ecology