

InitMIP-Antarctica experiments with the ice sheet model SICOPOLIS

*Ralf Greve¹, Ben Galton-Fenzi²

1. Institute of Low Temperature Science, Hokkaido University, 2. Australian Antarctic Division

The Ice Sheet Model Intercomparison Project for CMIP6 (ISMIP6) brings together a consortium of international ice sheet and climate models to explore the contribution from the Greenland and Antarctic ice sheets to future sea level rise. For such projections, initialisations are required that provide initial states of the respective ice sheet. InitMIP-Antarctica is an early initiative within ISMIP6 in order to explore this issue for the Antarctic ice sheet across a variety of models and initialisation techniques. We contribute to InitMIP-Antarctica with the ice sheet model SICOPOLIS and a spin-up-type initialisation, that is, a paleoclimatic simulation over 135 ka until the present. A major new component of the model is a physically-based parameterisation of ice shelf basal melting. In this parameterisation, basal melting of ice shelves is computed as a function of both the depth of ice below mean sea level and far-field ocean temperatures. The parameterisation is tuned differently for eight Antarctic sectors in order to achieve reasonable agreement with the modern spatial distribution of ice shelf basal melting. InitMIP-Antarctica also comprises three future climate scenarios, all to be run over 100 a: ctrl (present-day climate), asmb (prescribed schematic surface mass balance anomaly) and abmb (prescribed schematic basal melting anomaly under ice shelves). We present and discuss the performance of the spin-up in terms of agreement between simulated and observed present-day geometry and flow. Further, we investigate the response of the Antarctic ice sheet to the three future climate scenarios.

Keywords: Antarctica, Ice sheet, Ice shelf, Basal melting, Climate change, Modelling

Climate dependent contrast in surface mass balance in East Antarctica over the past 216 ka

*Shuji Fujita^{1,2}, Frédéric Parrenin^{3,4}, Ayako Abe-Ouchi^{5,6}, Kenji Kawamura^{1,2}, V. MASSON-DELMOTTE⁷, Hideaki Motoyama^{1,2}, Fuyuki Saito SAITO⁶, M. Fujita SEVERI⁸, B. STENNI⁹, Ryu Uemura¹⁰, E. W. Wolff¹¹

1. National Institute of Polar Research, Research Organization of Information and Systems(ROIS), 2. Department of Polar Science, The Graduate University for Advanced Studies (SOKENDAI), 3. CNRS, LGGE, F-38041 Grenoble, France, 4. University Grenoble Alpes, LGGE, F-38041 Grenoble, France, 5. Atmosphere and Ocean Research Institute (AORI), University of Tokyo, Chiba, Japan, 6. Japan Agency for Marine–Earth Science and Technology, Yokohama, Japan, 7. Laboratoire des Sciences du Climat et de l' Environnement, Institut Pierre Simon Laplace, UMR CEA-CNRS-UVSQ-UPS 8212, Gif-sur-Yvette, France, 8. Department of Chemistry, University of Florence, Florence, Italy, 9. Department of Environmental Sciences, Informatics and Statistics, Ca' Foscari University Venice, 30123 Venice, Italy, 10. Department Chemistry, Biology and Marine Science, Faculty of Science, University of the Ryukyus, Okinawa, Japan, 11. Department of Earth Sciences, University of Cambridge, UK

Documenting past changes in the East Antarctic surface mass balance is important to improve ice core chronologies and to constrain the ice-sheet contribution to global mean sea-level change. Here we reconstruct past changes in the ratio of surface mass balance (SMB ratio) between the EPICA Dome C (EDC) and Dome Fuji (DF) East Antarctica ice core sites, based on a precise volcanic synchronization of the two ice cores and on corrections for the vertical thinning of layers. During the past 216 000 a, this SMB ratio, denoted SMB_{EDC}/SMB_{DF} , varied between 0.7 and 1.1, being small during cold periods and large during warm periods. Our results therefore reveal larger amplitudes of changes in SMB at EDC compared with DF, consistent with previous results showing larger amplitudes of changes in water stable isotopes and estimated surface temperature at EDC compared with DF. Within the last glacial inception (Marine Isotope Stages, MIS-5c and MIS-5d), the SMB ratio deviates by up to 0.2 from what is expected based on differences in water stable isotope records. Moreover, the SMB ratio is constant throughout the late parts of the current and last interglacial periods, despite contrasting isotopic trends.

Keywords: Antarctica, precipitation, ice core

Improvement in dating of the Dome Fuji ice core using O_2/N_2 (80-165 ka)

*Ikumi Oyabu¹, Kenji Kawamura^{1,2,3}, Kyotaro Kitamura¹

1. National Institute of Polar Research, Research Organization of Information and Systems, 2. SOKENDAI (The Graduate University of Advanced Studies), 3. Japan Agency for Marine Science and Technology

The Dome Fuji ice core preserves valuable information on the climatic changes over the last 720 kyr, which enables to investigate forcings and mechanisms in the Earth's system. Precise ice core chronology is essential to determine sequences and durations of climate events, and to examine the phasing with forcings and other paleoclimatic records. Orbital scale variations in O_2/N_2 ratio of occluded air in the Vostok ice core are similar to those in local summer solstice insolation¹⁾. By using this similarity, Kawamura et al. (2007) synchronized the O_2/N_2 variations in the Dome Fuji and Vostok ice cores with local summer insolation, and established chronology for the ice cores with accuracy generally better than ± 2000 years (DFO-2006). However, it was recently pointed out by using detailed age matching between Dome Fuji, EDC and Chinese speleothem records that the DFO-2006 chronology around 90 kyr BP is too old by ~ 3 kyr³⁾. Possible cause of this error is dislocation of O_2/N_2 peaks because of large noise in the O_2/N_2 record. Recently, O_2/N_2 variations between 300 and 800 kyr has been reconstructed from EPICA Dome C ice core^{4),5)}, however, their records do not always show similar variation with local summer insolation. Thus, their O_2/N_2 record was not used for orbital tuning. These recent studies motivate us to examine the reliability of age markers based on the O_2/N_2 ratio of Dome Fuji ice core. In this study, we reanalyzed O_2/N_2 ratio in the first Dome Fuji ice core for 1200–1974 m, which covers 80–165 kyr BP.

Fractionation of O_2/N_2 ratio occurs from ice surface during storage⁶⁾. Because the first Dome Fuji core has been stored for about 20 years, we expect fractionated O_2/N_2 near the surface of ice. We thus tested different thickness of surface shaving, and found that shaving-off of about 1 cm of surface (and only using the inner part of the ice) is required for precise measurements for the ice samples below 1200 m depth. Because of this careful examination and improvement in methodology, our new O_2/N_2 data set on average do not indicate preferential loss of O_2 . Reproducibility of O_2/N_2 ratio are $\pm 0.425\%$ for 1200–1440 m, $\pm 0.263\%$ for 1440–1640 m and $\pm 0.088\%$ for 1640–1974 m, respectively.

We find large scatter in the new O_2/N_2 data between 1200 and 1440 m (typical amplitude: ~ 6 – 10%). This depth range is just below bubble-clathrate transition zone where both air bubbles and clathrate hydrates were observed by microscopes. To investigate the cause of this scatter, we conducted a high-resolution continuous analysis at 2.5-cm resolution for 1399.030–1399.484 m. The O_2/N_2 shows unexpected wave-like variation from -17 to -9% with a wavelength of ~ 18 cm. The amplitude is comparable to the typical scatter in the O_2/N_2 data set from the discrete 11-cm samples. If we take 11 cm average of the high-resolution data, the average ratio only varies by $\pm 1\%$, which is insufficient to explain the scatter in the discrete data. Our current speculation is that the wavelength and/or amplitude of the O_2/N_2 variations in other depths may be larger than those in the investigated section, and that O_2/N_2 of 11 cm samples can also vary by up to 10% .

Our new Dome Fuji O_2/N_2 record confirms strong correlation with local summer insolation. Assuming no phasing between O_2/N_2 and insolation variations, O_2/N_2 data was smoothed by a low-pass filter with the cut-off period of 16.7–10.0 kyr, and then tuned with local summer solstice insolation by peak-to-peak matching²⁾. Because the scatter in 1200–1440 m does not reflect insolation signal, we rejected data

points as outliers if they deviate from fitting curve by more than 3.3%.

Compared with the DFO-2006 chronology, the new age scale (DF-2016) is younger around 90 and 130 kyr BP, while it is older around 150 kyr BP. There were sharp steps in annual layer thickness (calculated from depth –age relationship) at 94.2 and 150.3 kyr BP from DFO-2006³⁾, but these unnatural steps disappeared in that from DF-2016. The DF-2016 and speleothem (U-Th) age scales agree within 1000 yrs. These results indicate that the revised chronology greatly improved from the DFO-2006 chronology.

To summarize, even though large O_2/N_2 fractionation occurs near the surface of the ice core over two decades of storage, the original O_2/N_2 ratio is preserved in the inner part of ice if it is stored at -50°C , and it can be precisely measured by sufficiently removing the ice surface. Accurate chronology can be constructed by orbital tuning of the high quality O_2/N_2 ratio from the Dome Fuji ice core with local summer insolation.

1) Bender, M. L., EPSL, 2002. 2) Kawamura et al., Nature, 2007. 3) Fujita et al., CP, 2015. 4) Landais et al., CP, 2012. 5) Bazin et al., CP, 2016. 6) Ikeda-Fukazawa et al., EPSL, 2005.

Keywords: ice core, Dome Fuji, chronology, O_2/N_2

Seafloor topography surveys around the East Antarctic continental margin

*Yoshifumi Nogi¹

1. National Institute of Polar Research

Seafloor topography is fundamental information for marine research. However, the detailed topography in the Southern Ocean has not generally been understood yet. Especially, seafloor topography around the Antarctica margin covered by sea ices has been unknown. The Antarctic continental margin is the boundary area between ice sheet and ocean and the bathymetry around this region is essential element to understand the interaction between ice sheet and ocean as well as the tectonic evolutions. Single beam echo soundings have been conducted by old icebreaker Shirase, and multi beam echo-sounder has been installed on new icebreaker Shirase and the swath bathymetry data have been obtained since the 51st Japanese Antarctic Research Expedition (2009-2010). Moreover, sub-bottom profiler has also been equipped on the new icebreaker. Unknown seafloor topography and sub-bottom profiles under sea ices around the Antarctic margin such as continental shelf and sloop are becoming clear, and those data are used as basic data for the Antarctic bottom water channels as well as paleoenvironment studies. Those data combined with magnetic and gravity anomalies have also contributed to the study related to the continental breakup. But there is a limit to observations by the icebreaker. The surveys under sea ices using ROV and/or AUV should be considered. We introduce the present status of multi beam echo-sounder and sub-bottom profiler obtained around the East Antarctic continental margin by icebreaker Shirase and future development of research using ROV and/or AUV are discussed.

Keywords: seafloor topography, East Antarctica, continental margin

Sea ice expansion in Antarctic warming events in the glacial Southern Ocean

*Minoru Ikehara¹, Kota Katsuki², Masako Yamane³, Yusuke Yokoyama³

1. Center for Advanced Marine Core Research, Kochi University, 2. Research Center for Coastal Lagoon Environments, Shimane University, 3. Atmosphere and Ocean Research Institute, University of Tokyo

The Southern Ocean has played an important role in the evolution of the global climate system. Area of sea ice shows a large seasonal variation in the Southern Ocean. Sea ice coverage on sea surface strongly affects the climate of the Southern Hemisphere through its impacts on the energy and gas budget, on the atmospheric circulation, on the hydrological cycle, and on the biological productivity. However, millennial-scale sea ice coverage and its impacts are not well understood. Here we show high-resolution records of sea ice-rafted debris (SIRD) and diatom assemblage to reveal a rapid change of sea ice distribution in the glacial Southern Ocean. The depositions of rock-fragment SIRD excluding volcanic glass and pumice were associated with increasing of sea-ice diatoms, suggesting that the millennial-scale events of cooling and sea-ice expansion were occurred in the glacial South Indian Ocean. The extent of sea ice in the Southern Ocean is occurred during the Antarctic isotope maximum (AIM) events, which is partly linked with the Heinrich Events in the Northern Hemisphere.

Keywords: Southern Ocean, sea-ice, Antarctic warming events, Heinrich Event

Radiolarian changes since the last glacial period in the Conrad Rise and their relation to the oceanic environments

*Takuya Itaki¹, Guiseppe Cortese², Minoru Ikehara³

1. Geological Survey of Japan, AIST, 2. Department of Paleontology, GNS Science, 3. Kochi University

It is well known that the oceanic circulation in the Southern Ocean plays an important role in the global climate changes. For reconstruction of the past ocean circulation in the Southern Ocean, siliceous microfossils such as diatoms and radiolarians preserved abundantly in deep-sea sediments are widely used as paleoceanographic proxies. Fossil assemblages of diatoms (phytoplankton) indicate usually surface water environments, while radiolarians (zooplankton) can be used as indicator for not only surface but also intermediate and deep water conditions because of their discrete habitat depths for each species. In this study, quantitative analysis of radiolarians was conducted for core COR-1bPC (54°S) from Conrad Rise in the Indian Ocean sector of the Southern Ocean.

Keywords: paleoceanography, biogenic productivity, intermediate water

Toward understanding the cause and mechanism of catastrophic collapse of Antarctic ice sheets during the last interglacial

*Osamu Seki¹

1. Institute of Low Temperature Science, Hokkaido University

Ongoing global warming could cause various influences to our lives. One of the most serious concerns caused by the global warming is sea level rise due to large scale collapse and melting of polar ice sheets. Recent observation revealed that the fastest melting rate of polar ice sheets ever observed is ongoing and there is growing concern of greatly rise of sea level in the future. Knowledge of sea level fluctuations in the past warm period provides useful information to better understand future sea level changes. Geological records have shown that sudden and abrupt rises in sea level (6 m rise within hundreds of years) were happened during the last interglacial (13,000-11,500 years ago) when global mean sea surface temperature (SST) was slightly higher (0.5 degree C) than the preindustrial level. If this were really happening, there is a critical ice sheet stability threshold resulting in the catastrophic collapse of polar ice sheets and substantial rapid sea-level rise in the interglacial climate condition. Since the Greenland ice core record showed a 2 m eustatic component from the Greenland ice sheet during the last interglacial, the Antarctic ice sheet greatly contributed to eustatic rise of sea level at that time. Given that the current global mean SST already reached the last interglacial level, it is urgent issue to evaluate whether massive collapse of the Antarctic ice sheets could occur in the future. For that purpose, it is necessary to investigate whether a massive collapse of the Antarctic ice sheet actually happened during the last interglacial or not. However, variability of the Antarctic ice sheets during the period has not been investigated. In this presentation, I will talk about an importance of research on the variability of the Antarctic ice sheets during past warm period such as the last interglacial to better understand the presence of “tipping point” into the new and irreversible melt regime of Antarctic ice sheets.

Keywords: Antarctic ice sheet, last interglacial, sea level rise

Amundsen Sea simulation with optimized ocean, sea ice, and thermodynamic ice shelf model parameters

*Yoshihiro Nakayama^{1,4}, Dimitris Menemenlis¹, Michael Schodlok¹, Ian Fenty¹, Ou Wang¹, An Nguyen², Patrick Heimbach², Pierre Dutrieux³, Eric Rignot^{1,4}

1. NASA Jet Propulsion Laboratory, 2. University of Texas at Austin, 3. Columbia University, 4. University of California, Irvine

The ice shelves and glaciers of the West Antarctic Ice Sheet (WAIS) are melting and thinning rapidly in the Amundsen Sea (AS) and Bellingshausen Sea (BS), with consequences for global sea level rise and ocean circulation. First, approximately 10% of the observed sea level rise has been attributed to the thinning of WAIS between 2005 and 2010. Second, the melting of ice shelves in the AS and BS will freshen the shelf water locally as well as downstream in the Ross Sea (RS), which may lead to a change in the characteristics of Antarctic Bottom Water formed in the RS and thus influence the global thermohaline circulation.

Agreement between model results and observations are crucial for understanding and projecting these impacts on the current and future climate. Thus, we aim to conduct model optimization for a regional Amundsen and Bellingshausen Seas configuration of the MITgcm. Currently, we have adjusted a small number of model parameters to better fit the available observations during the 2007-2010 period using trial-and-error adjustment and a Green's function approach. As a result of adjustments, our model shows significantly better match with observations than previous modeling studies, especially for Winter Water (WW). Since density of sea water depends largely on salinity at low temperature, this is important for assessing the impact of WW on Pine Island Glacier melt rate. We also conduct several sensitivity studies, showing the impact of surface heat loss on the thickness and properties of WW.

Our work is a first step toward improved representation of ice-shelf ocean interactions in the ECCO (Estimating the Circulation and Climate of the Ocean) global ocean retrospective analysis. In this presentation, we briefly explain our overall project and present some preliminary results pertaining to sensitivity simulations using high resolution (2 km) configuration and adjoint sensitivity simulations.

Keywords: Amundsen/Bellingshausen Sea, Ice shelf-ocean interaction, Circumpolar Deep Water

Observations of ice tongue-ocean interaction at Shirase Glacier

*Daisuke Hirano¹, Takeshi Tamura², Shuki Ushio², Kay I. Ohshima¹, Daisuke Simizu², Kazuya Ono¹, Tomohide Noguchi³, Shigeru Aoki¹

1. Institute of Low Temperature Science, Hokkaido University, 2. National Institute of Polar Research, 3. Marine Works Japan

Shirase Glacier Tongue (SGT) is a thick floating slab of ice that forms where the glacier flows down onto the ocean surface at the southern closed-section of Lutzow-Holm Bay (LHB) off Enderby Land, East Antarctica. Compared with other major ice shelves/tongues around Antarctica, SGT is smaller in area but its basal melt rate was estimated to be relatively high at a rate of ~7 m per year (Rignot et al., 2013) based on presence of warm deep water. Although comprehensive hydrographic observations in LHB is indispensable for understanding the SGT-ocean interaction, they are extremely limited, with exception of those conducted by wintering party of the 31st Japanese Antarctic Research Expedition (JARE) in 1990/92. Detailed analysis of the JARE-31 winter hydrographic observations suggests a 3-dimensional circulation, associated with the SGT-ocean interaction, that comprises: (1) warm modified CDW (Circumpolar Deep Water) flows southward at the deep layer of submarine canyon that leads into the region beneath SGT, (2) mCDW meets to melt the base of SGT, and (3) mixture of mCDW and basal melt water exports northward at subsurface layer.

To explore in detail the SGT-ocean interaction, summer comprehensive hydrographic observations in LHB are now in progress during JARE-58 in 2016/17 under the project called ROBOTICA. In this talk, preliminary results from the JARE-58 hydrographic observations are also presented.

Keywords: Shirase Glacier Tongue, basal melt, Circumpolar Deep Water

Breakup of land-fast sea ice in Lutzow-Holm Bay, East Antarctica and its teleconnection to tropical Pacific

*Shigeru Aoki¹

1. Hokkaido University

A large land-fast sea ice breakup occurred in 2016 in Lutzow-Holm Bay, East Antarctica. The breakup caused calving from the Shirase Glacier Tongue (SGT), which is otherwise held back by the ice. Although similar breakups and calving have been observed in the past, the timing and magnitudes are not well-constrained. We analyzed the ice's breakup latitude during 1997-2016 to investigate the variables controlling breakup and examine correlation with local calving for a longer period. The breakup latitude had a persistently high correlation with sea-surface temperature (SST) in the tropical Pacific, which exceeds correlations with local atmospheric variables. The multi-decadal variability of the tropical SST can explain the multi-decadal variation of the calving front of SGT from the 1950s through the breakup of fast ice. The SST-regressed breakup latitude can potentially explain 5 out of 6 SGT calving events from the mid-20th century, including its frontal retreat in the 1980s. Our proposed teleconnection between tropical SST and Antarctic sea ice could lead to better predictions of breakup and might impact the glacier flux for a wider region.

Standing meanders of the Antarctic Circumpolar Current as observed by Argo floats

*Katsuro Katsumata¹

1. JAMSTEC

Eddies in the Southern Ocean have been quantified by such measures as eddy kinetic energy and vertical eddy momentum transfer. Recent studies suggest length of streamlines can be used to characterise the eddy activity. The Antarctic Circumpolar Current, the greatest ocean current in the world, is known to have steady transport under increasing westerly wind stress in interdecadal time scales. From eddy resolving simulations, it has been hypothesised that some adjustment to the wind occurs on and around standing meanders of the ACC. One manifestation of the adjustment is the lengthening of streamlines. I used trajectory and hydrography data from Argo floats to study this behaviour in observed data. Seven major meanders were identified along the ACC. Eddies around the meanders have an effect to increase the radius of curvature, or stretch the streamlines. This effect was found generally on the eastern flanks of meridional ridges, where poleward eddy transport was found. The poleward eddy transport plays important roles in the meridional overturning circulation and momentum budget. Another important role in shaping the streamlines was highlighted.

Keywords: Antarctic Circumpolar Current, Interdecadal variability, Eddy transport, Curvature

Research prospects on variations and interactions of Antarctic ice sheet and climate

*Kenji Kawamura^{1,2,3}, Shin Sugiyama⁴, Ryu Uemura⁶, Hideaki Motoyama^{1,2}, Takanobu Sawagaki⁵, Iizuka Yoshinori⁴, Kazuho Horiuchi⁷, Shuji Aoki⁸, Kumiko Goto-Azuma^{1,2}, Shuji Fujita^{1,2}, Osamu Seki⁴, Motohiro Hirabayashi¹, Ikumi Oyabu¹

1. National Institute of Polar Research, Research Organization of Information and Systems, 2. SOKENDAI, 3. JAMSTEC, 4. Hokkaido University, 5. Hosei University, 6. University of Ryukyus, 7. Hirosaki University, 8. Tohoku University

We introduce our research plan over the next five years on variations and interactions of Antarctic ice sheet and climate. The research aims at better documenting and understanding of the Antarctic ice sheet and climate: their status, variability and interactions in the present and past. We particularly emphasize (1) analyses and proxy developments for reconstructing past, long-term temperature, accumulation rate, sea ice and carbon cycle, and (2) modern observations of the ice sheet margin and ocean by direct observation and remote sensing.

Keywords: Antarctic ice sheet, Climate change, Ice cores, Ice sheet melting

Current climate-ice sheet studies related to Antarctica and Southern ocean using Earth system models

*Ayako Abe-Ouchi¹, Fuyuki SAITO², Masakazu Yoshimori³, Akira Oka¹, Ralf Greve³, Keita Uehara¹, Wing-Le Chan¹

1. Atmosphere and Ocean Research Institute, The University of Tokyo, 2. JAMSTEC, 3. Hokkaido University

Climate-Antarctica ice sheet and Southern ocean interaction not only affects the local climate but also affect the global climate and ocean. We present some of our current modelling studies and plan using coupled atmosphere and ocean model and ice sheet model. In one of our studies, Eocene experiments were carried out to test the model's ability to reproduce proxy data and to investigate the climate system under high CO₂ concentration. Furthermore, in order to investigate to what extent topographical changes are responsible for the difference between the Eocene and present day climates, we created Eocene-like geometries from the present day topography. We present the individual effects of the Drake Passage, Tasman Gateway, Antarctica Ice Sheet on the climate.

Keywords: paleoclimate, Climate modeling, Antarctica ice sheet

A plan for studying the interaction of the solid Earth and the Antarctic ice sheet

*Yoichi Fukuda¹, Jun Nishijima², Takahito Kazama¹, Kazuki Nakamura³, Koichiro Doi⁴, Yusuke Suganuma⁴, Jun'ichi Okuno⁴, Akito Araya⁵, Heitaro Kaneda⁶, Yuichi Aoyama⁴, Hideki Miura⁴

1. Graduate School of Science, Kyoto University, 2. Graduate School of Engineering, Kyushu University, 3. Faculty of Engineering, Nihon University, 4. National Institute of Polar Research, 5. ERI, University of Tokyo, 6. Graduate School of Science, Chiba University

The Antarctic ice sheet, which relates to the global climate changes through the sea level rise and ocean circulation, is an essential element of the Earth system for predicting the future environment changes. Thus many studies of the ice sheet changes have been conducted by means of geomorphological, geological, geodetic surveys, as well as satellite gravimetry and satellite altimetry. For these studies, one of the largest uncertainties is the effects of GIA, which, on the other hand, includes valuable information about the rheological properties of the solid Earth, because GIA is the rheological response of the solid Earth to the ice mass loading. The observational studies of the GIA effects should greatly contribute to investigate the inner structure of the Earth.

GIA as a keyword to investigate the interaction between the solid Earth and the ice sheet changes is an urgent and important research target not only for a practical requirement of predicting global changes but also for a more pure scientific interest to know the structures of the deep Earth's interior.

In view of these points, in addition to the several precise observations at Syowa station and surrounding areas, we plan to conduct geomorphological, geological and geodetic surveys in the inland mountain areas and the coastal areas in East Antarctica, where the in-situ data for constraining GIA models are very few.

Combining these new observations with other in-site data, various satellite data and numerical modeling, we aim to estimating a precise GIA model, constructing a reliable ice melting history after LGM (the Last Glacial Maximum) and obtaining the viscoelastic structure of the Earth's interior.

To achieve the goal, we are planing to conduct the following studies;

- (1) conducting glacial topographic surveys, geological surveys, gravity measurements, and GNSS measurements at the in-land areas of the East Antarctica, where very few observations were conducted so far, and reevaluating the glacial topography using the in-situ observations and recent precise DEM (Digital Elevation Model),
- (2) improving the accuracies of the retreat ages of the ice sheets using the micro glacier topography from the detailed airborne photographic data obtained by unmanned aerial vehicles, the cosmogenic nuclide exposure ages of the basement bowling samples and the moraine rocks,
- (3) monitoring the present day ice sheet movements and sea level changes by combining satellite data and in-situ geodetic and other observations in and around Syowa Station, and
- (4) finally aiming at the quantitative reconstruction of the ice melting history over the last millions years, and the improvement of the models for predicting the future global changes.

Keywords: ice sheet, sea level rise, Glacial Isostatic Adjustment, ice sheet metling history, East Antarctica, viscoelastic structure

Food web in the marginal ice zone: material flow from sea ice through to myctophid fish

*Masato Moteki^{1,2}, Ryosuke Makabe^{2,3}, Kohei Mizobata¹, Shintaro Takao^{2,3}, Keishi Shimada¹, Naho Miyazaki¹, Kunio Takahashi^{2,3}, Tsuneo Odate^{2,3}

1. Tokyo University of Marine Science and Technology, 2. National Institute of Polar Research, 3. The Graduate University for Advanced Studies

The seasonal prevalence of the sea ice zone is a critical element in the Southern Ocean ecosystem structure and dynamics. Sea ice formed in coastal polynyas in autumn and winter is transported northward and covers a vast area of the Southern Ocean. This ice contains a large variety of sea ice biota (SIB), such as ice algae, protozoans, and crustacean larvae. The SIB is released as the sea ice melts from the ice edge in spring and summer. Although a flood of SIB biomass is released into the water column, there is little information on the pathway and dynamics of the SIB released into the Southern Ocean. The sea ice changes that occur with climate change should affect the Southern Ocean ecosystem via this pathway.

We have been investigating the flow of materials derived from SIB through to the Southern Ocean ecosystem after its release into the water column. In a series of studies, we found the nursery grounds of the Antarctic myctophid fish *Electrona antarctica*, which is an important component of the oceanic food web, in the waters influenced by sea ice. This paper introduces a study of 1) the dynamics of SIB inferred from the flora of floating sea ice (ice algae) and the water column at the ice edge and 2) detritus containing diatoms found in the stomachs of *E. antarctica* larvae. The origin of the diatoms is discussed.

A comparison of the numbers of algal cells in sea ice and the water column revealed that more than 90% of the cells of the dominant diatom species were removed from the surface mixed layer. Most of the algal cells were thought to sink into deeper waters or to be grazed by zooplankton close to the sea ice. Detritus served as a food item for early stage larvae of *E. antarctica*, and some of the diatom species in the detritus were usually dominant in the SIB assemblage, suggesting that the *E. antarctica* larvae feed on SIB indirectly, eating the faecal pellets of zooplankton or aggregates.

In January 2017, we conducted drifter observations using a sediment trap in the vicinity of the ice edge; these should provide more direct information on the dynamics of sinking material from the SIB. Regarding the food habits of *E. antarctica* larvae, a metagenome analysis of the diatoms in the stomach and a biomarker analysis will be conducted to determine the detailed pathway of material flow from sea ice.

Keywords: sea ice biota, ice algae, food web, marginal ice zone, marine snow, detritus

Paleo-environmental changes at coastal lakes along the Soya Coast, East Antarctica during the Holocene

*IJIN KANG¹, Kaoru Kashima², Koji Seto³, Yukinori Tani⁴, Takahiro Watanabe⁵, Toshio Nakamura⁶, Satoshi Imura⁷, Genki I. Matsumoto⁸

1. Department of Earth and Planetary Sciences, Graduate School of Sciences, Kyushu University, 2. Department of Earth and Planetary Sciences, Faculty of Sciences, Kyushu University, 3. Research Center for Coastal Lagoon Environments, Shimane University, 4. Institute of Environmental Sciences, University of Shizuoka, 5. Japan Atomic Energy Agency, 6. Center for Chronological Research, Nagoya University, 7. National Institute of Polar Research, 8. School of Social Information Studies, Otsuma Women's University

The East Antarctic Ice Sheet (EAIS) is the largest glacial system on Earth, and documenting its changes is important to understand and estimate its future behavior. Antarctic coastal lakes are invaluable archives of paleo-climate and paleo-environment changes caused by the retreat of Antarctic Ice Sheet. In Soya Kaigan (Coast) of Lutzow-Holm Bay region, many coastal lakes are located in ice-free areas. This study discussed the environmental change inferred from microscopic observation of fossil diatom assemblages in a sediment cores from coastal freshwater lakes, Lake Oyako-ike, Lake Maruwanminami-ike and Lake Maruwan-Oike, in Soya Coast along with biomarkers and microscopic observation of microalgae and cyanobacteria, sedimentary facies and AMS 14C dating.

Three lakes are considered as marine relict lakes resulted from the recession of glaciers and subsequent isostatic uplift. The Ok4C-01 core (length 135 cm) from Lake Oyako-ike was divided in 4 zones according to the diatom assemblage changes. This lake has changed from costal marine to freshwater lake at ca. 1100 cal yr BP (core depth 60 cm). The MwS4C-01 core (length 147 cm) from Lake Maruwanminami-ike was also divided in 4 zones. The transition timing from marine to freshwater lake was ca. 2400 cal yr BP (core depth 65 cm). The Mw4C-01 core (length 226 cm) from Lake Maruwan-Oike was divided in 4 zones as well. This lake has changed at ca. 2800 cal yr BP (core depth 22 cm). Diatom assemblage changes in these sediment cores show similar pattern with other analyses results such as sediment facies and elemental analyses (TC, TS, TN contents).

Diatom assemblages in these sediment samples implied that more specific environmental changes not only transition from marine to freshwater but also an existence of sea-ice (*Fragilariopsis curta*, *F. cylindrus*), a desalination process (*Psammothidium papilio*) and an oligotrophication of lake water.

Keywords: Antarctica, Paleoenvironment, Lake sediment, Diatom assemblage

Pliocene seasonal sea-ice history around the Kerguelen Plateau in the Southern Ocean based on diatom analysis

*Saki Ishino¹, Yuji Kato¹, Itsuki Suto¹

1. Graduate School of Environmental Studies, Nagoya University

Fossil diatoms have been regarded as a useful proxy for paleoceanographic reconstruction such as past sea-ice distribution in the Southern Ocean (Armand et al., 2005). Several studies have attempted to reveal sea-ice history based on fluctuation of the sea-ice related diatoms, however there is few paleoceanographic studies focusing on changes of summer and winter sea-ice distribution. We categorized the sea-ice related diatoms reported in Armand et al. (2005) into two groups, summer sea-ice diatom species and winter sea-ice diatom species. The former group is abundantly observed within summer sea-ice zone and the latter group is widely found within winter sea-ice zone. In this study, late Pliocene summer and winter sea-ice distributions in the Indian sector of the Southern Ocean are estimated based on temporal changes in abundance of the summer and winter sea-ice diatom species. Analyzed samples are ODP Leg 188 Hole 1165B and Leg 119 Hole 745B, which are located near recent summer and winter sea-ice edge respectively.

Keywords: Southern Ocean, paleoenvironment, sea ice, ODP, diatom

The middle Miocene paleoceanography based on diatoms and chrysophyte cysts in the Atlantic sector of the Southern Ocean

*Yuji Kato¹, Saki Ishino¹, Itsuki Suto¹

1. Graduate School of Environmental Studies, Nagoya University

The Southern Ocean, which is composed of the Antarctic Circumpolar Current (ACC) and several water masses, is an important element as having great influence on global climate change. Therefore, it is significant to reconstruct the paleoceanographic history in the Southern Ocean including the ACC, and many researchers have been approaching this issue using various methods. For example, changes in sediment structures derived from seismic profiling of deep-sea sediments have been widely accepted as an evidence for the past ACC fluctuations. However, there are almost no previous studies that show long-term and detailed ACC changes, while many of paleo-ACC studies have been made with order of “epoch” or have been focused on much shorter time-range.

In the Southern Ocean, abundant and continuous occurrences of siliceous microfossils such as diatoms and chrysophyte cysts are often found from the Neogene sediments. Generally, diatoms thrive in high-latitude or upwelling regions and they show clear segregation of their habitat corresponding to the distribution of the water masses. Therefore, it is likely that the changes in abundance of subtropical diatom taxa, which inhabit north of the ACC, reveal the long-term migration history of the ACC from north to south vice versa. In addition, as most chrysophytes inhabit freshwater environment, the occurrences of fossil chrysophyte cysts in the Southern Ocean are mainly originated from Antarctic terrestrial freshwater (i.e., melt-water of the ice sheet). Therefore, fossil chrysophyte cyst can be treated as a useful tool to reconstruct the changes of the Antarctic ice sheet in the geological past. In this presentation, we would like to present and discuss on several paleoenvironmental events in the Southern Ocean during the late Miocene–Pliocene (ca. 9–3 Ma) based on fossil diatom and chrysophyte cyst analysis preserved in sediment core samples from the Atlantic sector of the Southern Ocean (ODP Leg 113 Site 689 and DSDP Leg 71 Site 513), which include repetitive north-south migrations of the ACC in a 100–200 kyr cycle observed in 7–4.8 Ma.

Keywords: Diatom, Chrysophyte cyst, Southern Ocean, Antarctic Circumpolar Current, DSDP, ODP

Climate model experiments using state-of-the art boundary conditions for the Mid to Late Pliocene

*Wing-Le Chan¹, Ayako Abe-Ouchi^{1,2}

1. Atmosphere and Ocean Research Institute, The University of Tokyo, 2. Dept. of Integrated Climate Change Projection Research, JAMSTEC, Yokohama

The Mid to Late Pliocene (3.3-3.0 million years ago) represented a period during which atmospheric CO₂ concentrations were similar to that of present day. Globally, the climate was warmer by about 2-3°C, and warmer for prolonged periods. The idea of an analogue to future global climate change, albeit with caveats, has created much interest in this period, with focus on both climate proxy data and model simulations. At a time when there was reduced glaciation in Antarctica, there is evidence from benthic foraminifera to suggest that North Pacific deep waters were much colder than North Atlantic Deep Water and that subsequent glaciation of Antarctica had global ramifications by altering the deep ocean circulation and contributing to the intensification of glaciation in the opposite hemisphere.

The Pliocene Model Intercomparison Project (PlioMIP) was established by bringing together the paleodata analysis group, PRISM, and various climate modelling groups to further our knowledge of this period. Specific protocols have been set up for climate model experimental design, utilizing the latest paleoenvironmental reconstruction datasets which include Pliocene vegetation, soils, ice distribution and ocean bathymetry. In particular, Pliocene ice sheet reconstructions depict a West Antarctic seaway, no ice over West Antarctica, small increases in the elevation in the interior of Antarctica and retreat of ice sheet in the low-lying Wilkes and Aurora subglacial basins, in accordance to proxy evidence. In the present study, we ran experiments using the atmosphere-ocean coupled model, MIROC4m, to investigate their effects on the climate by incorporating all these latest boundary conditions from PlioMIP2. Related sensitivity experiments help to quantify the relative contribution to Pliocene warmth from individual boundary conditions and to investigate the climate and Earth system sensitivity.

Keywords: Pliocene, Climate modelling, Climate change, Antarctic ice sheet

Sensitivity studies of ice divide position using a numerical ice-sheet/shelf model

*Fuyuki SAITO¹, Ayako Abe-Ouchi^{2,1}

1. Japan Agency for Marine-Earth Science and Technology, 2. Atmosphere Ocean Research Institute, University of Tokyo

Ice divides are important locations for drilling on ice-sheets. Since the ice flow pattern, which affects dating, is significantly different between an ice divide and the other areas, sensitivity of the ice-divide position to changes in various boundary conditions should be investigated. In this study, numerical experiments under synthetic configurations are systematically conducted using a numerical ice-sheet/shelf model IcIES, to evaluate how local bedrock topography and/or ice-sheet extent affects the ice-divide position.

Keywords: ice-sheet

Responses of marine ice sheet to basal melting of ice shelves

*Takashi Obase¹, Ayako Abe-Ouchi^{1,2}, Fuyuki SAITO²

1. The University of Tokyo, Department of Earth Planetary Science, 2. Japan Agency for Marine-Earth Science and Technology

Integrated study of combining climate models and ice sheet models are required to understand the response of Antarctic ice sheet to climate changes, and basal melting beneath ice shelves is a key player (Mengel et al. 2015; Deconto and Pollard 2016; Kusahara 2016). Simulations by high-resolution ocean models that resolves ice shelf cavity circulations show that basal melt rate of ice shelves increase drastically in a warm climates (Timmermann et al. 2013; Obase et al. in press), and could cause collapse of marine ice sheet.

Previous studies investigated the model dependency on the response of marine ice sheet to changing ice accumulation, ice temperature, and basal sliding, but not for changing basal mass balance (Vieli and Payne 2005; Pattyn et al. 2012; Pattyn et al. 2013). Recent study with explicit treatment of basal melt rate suggests that careful treatment of basal melt rate near grounding line and higher horizontal resolution are required for simulating grounding line retreat because of abrupt change in basal mass balance near grounding lines (Gladstone et al. 2017).

In this study, we investigate the responses of marine ice sheet to basal melt rate beneath ice shelves under an idealized flow-line system. We use a numerical ice sheet-shelf model IcIES for flow-line calculation. Ice flow is approximated by Shallow Ice Approximation for grounded ice and Shallow Shelf Approximation for floating ice, and sub-grid scale grounding line migration is parameterized with ice flux at the grounding line (Schoof 2007). In the flow-line calculation, ice flux that is orthogonal to ice flow and lateral resistive stress are set to zero. The ice sheet and ice shelf are assumed to be isothermal, therefore thermodynamics is not included. Bedrock topography is taken from several ice sheet drainages from Antarctic ice sheet. We systematically simulate steady-states of ice sheet shape under a given basal melt rate and bedrock topography. We discuss the determining factors of changing basal melt rate, changing sea level, and bedrock topography on the stability of marine ice sheet.

West Antarctic ice shelf melting causes Ross Sea freshening and Circumpolar Deep Water warming.

*Yoshihiro Nakayama^{1,2}, Ralph Timmermann¹, Hartmut H Hellmer¹

1. Alfred Wegener Institute, 2. NASA Jet Propulsion Laboratory

Ross Sea (RS) freshening and Circumpolar Deep Water (CDW) warming have been unveiled from oceanographic observations, yet responsible mechanisms remain uncertain. Using a sea-ice/ice-shelf/ocean model, we show that enhanced ice shelf melting in West Antarctica explains the observed changes. The RS continental shelf freshening is caused by an increase in ice shelf meltwater from the Amundsen/Bellingshausen Sea. This weakens the Antarctic Bottom Water formation in the RS, which reduces the density of mid-depth and deep water, allowing CDW to shift further south and causing open ocean warming. The warming signal is transmitted onto the Amundsen/Bellingshausen Sea continental shelves including the ice shelf cavities, implying a positive feedback. Good agreement between simulations and observations suggests that the proposed mechanism is able to explain the ongoing RS freshening and CDW warming. Warmer water on the shelf likely enhances ice shelf melting and may amplify the rate of sea level rise.

Keywords: Ice shelf-ocean interaction, Amundsen-Bellingshausen Sea, Ross Sea , Circumpolar Deep Water

Possibility of AABW source originating from meddle size of Polynya along the coast of Australian-Antarctic Basin

*Kitade Yujiro¹, Keishi Shimada¹, Yuki Ogata¹, Shigeru Aoki², Taiyo Kobayashi⁴, Kohei Mizobata¹, Takeshi Tamura³, Toshio Suga⁵, Kay I. Ohshima²

1. Tokyo University of Marine Science and Technology, 2. University of Tokyo, 3. National Institute of Polar Research, 4. Japan Agency for Marine-Earth Science and Technology, 5. Tohoku University

Antarctic Bottom Water (AABW) originating from Vincennes Polynya (VP) was discovered recently (Kitade et al. 2014). The fact that middle size of Polynya also produces AABW suggests the possibility that the unknown formation area still exists along the coast of Australian-Antarctic Basin (AA basin). From the viewpoint of sea ice production, the amount of sea ice production of Sackleton Polynya (SP) in the west of VP is about 1.4 times that of VP (Tamura et al. 2016), implying that SP is a possible candidate for AABW formation area.

A deep float, called “Deep NINJA” which is able to observe temperature and salinity at depths up to 4,000 m, was developed by Japan Agency for Marine-Earth Science and Technology and Tsurumi-Seiki Co. (Kobayashi et al., 2015). Five deep floats were deployed along 110°E in Jan. 2014. One of them drifted westward almost along the continental rise and had been observing 40 profiles within two years. However, no signal of newly formed AABW was observed except off VP, and such property is consistent with The Baseline Research on Oceanography, Krill and the Environment (BROKE) survey. Therefore, as a result of investigating the salinity of Dense Shelf Water (DSW) on the shelf in the elephant seal bio-logging data, it was found that the salinity of DSW’ s core in offshore SP is 0.1 or lower than that in offshore VP. Although this cannot be explained by sea ice production difference, it can be explained by considering AVISO absolute dynamic topography data, ocean climatology data and sea ice melting amount. Moreover, it was thought that AABW was not formed offshore of the SP’ s shelf because the salt content of DSW was insufficient for AABW formation. Although these facts do not completely negate the additional formation of AABW originating from middle size of polynya located at west of VP, their formation volume of AABW is suggested to be much smaller than that off VP. As a result of investigating the possibility of AABW formation in other medium size polynya using the same algorithm, Dibble Polynya was considered as the most influential unconfirmed candidate in the polynya facing the AA basin.

Keywords: Antarctic Bottom Water, Dense Shelf Water, Australian-Antarctic Basin, Formation process of shelf water , Vincennes Polynya, Sackleton Polynya

Recent quick changes of Antarctic Bottom Water off the Adélie Coast, Antarctica

*Taiyo Kobayashi¹

1. Japan Agency for Marine-Earth Science and Technology

In the study, changes of Antarctic Bottom Water (AABW) off the Adélie Coast, Antarctica, were examined mainly with observations of deep floats for December 2012 to August 2014. AABW was observed to have disappeared rapidly in the order of the densest part and its thickness had decreased quickly by around 45 m yr^{-1} , several times of the rate for the recent decades. Temperature and salinity on isopycnals showed seasonal changes, but there were no clear trends. The results of the repeat hydrographies clarified that the rapid deepening would have begun around 2010/11, at the latest and that AABW were largely freshened by around 0.005 in 2011. The changes of AABW ought to have raised the sea level by around $4.7 \text{ (3.1-6.5) mm yr}^{-1}$ for 1900-4000 dbar, which agreed well with the independent observations within errors; 5.8 mm yr^{-1} of Aviso, 0.5 mm yr^{-1} of Argo for 0-1900 dbar, and 1.8 mm yr^{-1} of the mass component at averages for 2011-2014. The collapse of Mertz Glacier Tongue in February 2010 was expected to lead the larger changes of AABW. Its rapid disappearance would be caused by the smaller supply of ALBW due to the smaller sea ice production there and it would continue for a longer time because the revolutionary change of the ocean conditions would hinder the supply of a locally formed coastal dense water from recovering to the similar level before the collapse. However, the “long-term” freshening might be changed less by the collapse; the “large” freshening in 2011 would correspond to the freshening due to the global warming for about 5-6 years, and then very small changes would follow for several years.

Keywords: deep and bottom water, float obseravation

Increasing fresh water impact on Sea Level Rise in Australian-Antarctic Basin

*Keishi Shimada¹, Yujiro Kitade¹, Shigeru Aoki², Kay I. Ohshima², Kohei Mizobata¹, Takeshi Tamura³

1. Tokyo University of Marine Science and Technology, 2. Institute of Low Temperature Science Hokkaido University, 3. National Institute of Polar Research

Significant warming and freshening of Antarctic Bottom Water (AABW) in the Australian-Antarctic Basin are reported in previous studies. The reported changes are, however, based on repeated hydrographic data with 5-10-year interval, and hence, may be subject to temporal aliasing due to shorter time scale variability. In this study, we have assessed temperature and salinity using repeated hydrographic data with 1-2-year interval. The Conductivity-Temperature-Depth (CTD) data are obtained by the training and research vessel Umitaka-maru along 110°E, where locates slightly west of WOCE section 109s. In the region south of 60°S, both warming and freshening trends in deep layers (e.g., below 2000 m) are clear since 1990s and they are consistent with previous studies. However, it is found that temperature and salinity changes are not strongly correlated. Linear freshening trend is clearly intensified to bottom (~ -0.01 PSS-78/decade). However, warming trend includes shorter vertical scale variations; taking maximum at 3000-3500 m (~ 0.025 °C/decade), but weakened toward bottom. In deep layers below 2000 m, as for freshening, linear trend component explains over 90% of standard deviation. As for warming, however, it explains only about 50 % on average, meaning that shorter time scale variability effects only for temperature variations. Finally, impact of linear warming and freshening trends on Sea Level Rise (SLR) are estimated. Reflecting the dominant linear trend component, impact of freshening in deep layers explains 60% of overall SLR trend. This implies that deep freshening can be main component that induces SLR in near future.

Keywords: Antarctic Bottom Water, warming, freshening, sea level rise

Temporal and spatial variability of atmosphere and ocean in the Southern Ocean

*Masafumi Yagi¹, Kunio Kutsuwada², Mikio Naganobu³

1. School of Geo-environmental Science, Tokai University, 2. School of Marine Science and Technology, Tokai University, 3. NRIFS/Fisheries Research Agency

The variability of atmospheric circulation in the Southern Ocean (SO) is dominated by a zonally symmetric pattern, that is the Southern Annular Mode (SAM) (Thompson and Wallace, 2000), which is characterized by a dipole in the westerly winds with opposing centers between mid- and high-latitudes. Especially, the westerly winds prevail in the Drake Passage, suggesting key area of oceanic and atmospheric variations. Our previous study focusing on structures in surface oceanic layer near the Drake Passage clarified that their year-to-year variations have relationships with the DPOI which is a good indication of surface winds across the Passage and related to the Krill recruitments (Naganobu et al., 1999, 2008). In addition, we examine year-to-year variations in the surface wind field over the SO characterized by the DPOI, and detected another variation associated with the meridional shift of the westerly wind area over the Southern Ocean (Indian-Pacific-Atlantic sector) and having high correlation with the DPOI (Yagi et al., 2017). However, the influence of DPOI-related variation in the surface winds on the oceanic field still remains unclear. In this study, we investigate relationships between the sea-surface-temperature (SST) and surface wind variations over the SO.

The leading mode, derived from the empirical orthogonal function (EOF) analysis for the monthly-mean zonal wind field, has a contribution of 55.5% with maxima on the Drake Passage. The score of this mode has a high correlation with the DPOI (0.94), so is related to the DPOI. Its spectrum has significant peaks at the periods of 6-month and 12-months, and another peak at the periods of 30-months. We focus on variabilities with time scales of about 3 years. To examine relationship between the DPOI-related surface wind and SST over the SO, we derive time series of the score of the 1st EOF mode for the zonal wind which are band-pass filtered for periods around 3 years, and examine its spatial correlations with the SST field. Results reveal that high correlations are found over the entire SO region. Spatial correlations indicate that there are negative areas to the west of the Drake Passage and positive ones to the east of it, and maxima in the Ross and Weddell seas. Relationships will be examined between surface wind and SST fields in these regions in detail.

Keywords: Westerly Wind, DPOI, Air-Sea interaction

Variability of Antarctic coastal polynyas and their linkage with fast ice revealed from AMSR-E and AMSR2 data

*Sohey Nihashi¹, Kay I. Ohshima², Takeshi Tamura³

1. National Institute of Technology, Tomakomai College, 2. Institute of Low Temperature Science, Hokkaido University, 3. National Institute of Polar Research

A coastal polynya is newly-forming thin sea-ice areas formed by divergent ice motion driven by prevailing winds and/or ocean currents. In coastal polynyas, huge amounts of heat flux from the ocean to the atmosphere occur because the heat insulation effect of sea-ice is greatly reduced in the case of thin ice, and accordingly sea ice is formed actively. Dense water formed in Antarctic coastal polynyas with the intense sea-ice production is a major source of Antarctic Bottom Water (AABW), which is a key player in the global climate system.

In Antarctic coastal polynya areas, algorithms that detect the polynya areas and estimate the thin ice thickness from passive microwave satellite data (SSM/I-SSMIS: 1992—present or AMSR-E: 2003—2011) had been developed to estimate the ice and dense water production. Fast ice areas were also detected using the passive microwave data. The spatial (grid) resolution of AMSR-E (about 6 km at 89 GHz) is four times higher than that of SSM/I-SSMIS in the pixel density. This advantage of AMSR-E is critical for the monitoring of the coastal polynyas and fast ice because of their small areal extent (i.e., from 10 to 100 km at most).

The coincident circumpolar mapping of Antarctic coastal polynyas and fast ice from AMSR-E had revealed that most of the polynyas are formed on the western side of fast ice or glacier tongue, indicating an important role of fast ice and glacier tongue in the polynya formation. Because the fast ice and glacier tongue are particularly vulnerable to oceanic and atmospheric conditions, their extent can be changed drastically and suddenly. The change in the fast ice or glacier tongue can cause dramatic changes in sea-ice production in the adjacent polynya and possibly AABW formation, as in the case of the Mertz Glacier Tongue (MGT) calving in 2010. This can potentially contribute to further climate change. Although AMSR-E failed in October 2011, AMSR2, the successor to AMSR-E, was launched in May 2012. The spatial resolution of AMSR2 is improved about 17% from AMSR-E.

This study developed an algorithm which can detect the polynya area and can estimate the thin ice thickness from AMSR2 data, based on a similar method to the AMSR-E algorithm development. Fast ice areas were also detected using AMSR2 data. Ice production in the polynyas was estimated from heat flux calculation using the sea-ice data from AMSR2. In the major polynyas, the AMSR2 ice production was compared with the AMSR-E ice production though a comparison with the SSM/I-SSMIS ice production. The comparison confirmed that the AMSR2 and AMSR-E data with higher spatial resolution can be used for time series analysis of the relationship between coastal polynyas and fast ice for >10 years. For example, maps of annual ice production and fast ice from AMSR2 and AMSR-E can reveal the details of the Mertz Polynya change before and after the MGT calving. Continuous monitoring of the coastal polynyas by the AMSR series is essential for climate-change-related analyses in the Antarctic Ocean.

Keywords: AMSR2, coastal polynyas, fast ice

Circumpolar mapping of the Antarctic coastal polynyas with discrimination of ice type

*Kazuki Nakata¹, Kay I. Ohshima¹, Sohey Nihashi²

1. Institute of Low Temperature Science, Hokkaido University, 2. National Institute of Technology, Tomakomai College

In Antarctic coastal polynyas, high production of sea ice occurs due to huge heat loss to atmosphere, resulting in the formation of dense water, precursor of Antarctic Bottom Water. Sea-ice production within polynyas is directly related to polynya extent and thin-ice thickness within the polynya. Thus, it is important for the calculation of sea-ice production to estimate thin ice thickness accurately. Several studies have developed algorithms for estimation of the thin ice thickness from brightness temperature (TB) of satellite passive microwave sensor [e.g., Martin et al., 2004; Nihashi and Ohshima, 2015]. In these algorithms, ice thickness of less than 20 cm is empirically estimated by utilizing negative correlation between the ice thickness and a ratio of the horizontally to vertically polarized TBs (PR). Several studies have also extended these algorithms to mapping of sea-ice production for Antarctic coastal polynyas. However, ice type, which has an influence on microwave characteristic of thin-ice, has not been considered in these algorithms.

Thin ice (polynya) areas are classified roughly into two ice types. One is active frazil type: a mixture of open water and frazil/grease ice areas. The other is thin solid ice type: nearly uniform thin ice covered area. Nakata et al. [in prep.] indicated that PR-thickness relationships are different clearly between these two ice types: active frazil type has much smaller thickness than solid ice for the same PR value. Based on the result, Nakata et al. [in prep.] also developed a thin ice algorithm in which ice thickness for each ice type is estimated from the corresponding empirical equation after discrimination of ice type. This improved algorithm provides more accurate estimation of sea-ice production. In addition, the algorithm can specify a predominant ice type for each Antarctic coastal polynya, which is useful for examination of the polynya dynamics. In this study, we apply the new algorithm to the entire Southern Ocean and carry out mapping of the Antarctic coastal polynya and its ice production.

We used AMSR-E/Aqua Level 2A (L2A) global swath spatially resampled TBs at 36 and 89GHz. We first mapped all AMSR-E L2A data obtained within a day onto the NSIDC polar stereographic grid (the spatial resolution of 6.25 km), with the land and fast ice mask by Nihashi and Ohshima [2015]. Then, we use the algorithm to obtain ice-type and thin ice thickness. Sea-ice production is calculated from heat flux calculation using the obtained thin ice thickness and ERA-Interim atmospheric data. From the above procedure, we create a data set of ice type, thin ice thickness and ice production for the entire Southern Ocean on a daily basis, during winter (April-October) for the period 2003-2010.

The climatological mapping shows that the active frazil type is more predominant in polynyas in the East Antarctica, where the strong offshore wind is prevailing. Thin solid ice type is predominant in polynyas with relative weak wind, such as in the Ross Ice Shelf polynya. These suggest that the difference in predominant ice type is mainly caused by the difference in strength of offshore wind.

In the previous algorithms, ice thickness was overestimated because the PR-thickness relationship is similar to that of the thin solid ice type. In the new algorithm, sea-ice production in the polynyas along the East Antarctica is corrected. Especially, sea ice production in the Cape Darnley polynya with the highest occurrence frequency of active frazil type is calculated to be about 1.5 times as that of the previous studies.

Keywords: coastal polynya, ice type, sea-ice production, AMSR-E

