

## 太陽型星における恒星黒点とスーパーフレア

### Starspot activity and superflares on solar-type stars

\*前原 裕之<sup>1</sup>、野津 湧太<sup>2</sup>、野津 翔太<sup>2</sup>、行方 宏介<sup>2</sup>、本田 敏志<sup>3</sup>、石井 貴子<sup>2</sup>、野上 大作<sup>2</sup>、柴田 一成<sup>2</sup>  
\*Hiroyuki Maehara<sup>1</sup>, Yuta Notsu<sup>2</sup>, Shota Notsu<sup>2</sup>, Kosuke Namekata<sup>2</sup>, Satoshi Honda<sup>3</sup>, Takako T. Ishii<sup>2</sup>, Daisaku Nogami<sup>2</sup>, Kazunari Shibata<sup>2</sup>

1. 自然科学研究機構国立天文台、2. 京都大学、3. 兵庫県立大学

1. National Astronomical Observatory of Japan, NINS, 2. Kyoto University, 3. University of Hyogo

Recent space-based observations (e.g., Kepler mission) enable us to investigate the nature of “superflares” on solar-type stars (G-type main sequence stars). The bolometric energy of superflares ranges from  $10^{33}$  erg to  $10^{36}$  erg which is  $10$ - $10^4$  times larger than that released by a typical X10 class solar flare. Most of the stars with superflares show large-amplitude photometric variations associated with the stellar rotation which suggest that the stars with superflares have large starspots. Spectroscopic studies of superflare stars revealed that the chromospheric activity correlates with the amplitude of brightness variations.

We analyze the correlation between starspots and superflares on solar-type stars using the data from the Kepler mission. Our analysis shows that the fraction of the stars showing superflares decreases as the rotation period increases and as the amplitude of photometric variations, which is thought to correlate with the area of starspots, decreases. We found that the fraction of superflare stars among the stars with large starspots also decreases as the rotation period increases. This suggests that some of the slowly-rotating stars with large starspots show a much lower flare activity than the superflare stars with the same spot area and rotation period.

Assuming simple relations between spot area and life time and between spot temperature and photospheric temperature, we compared the size distribution of large starspots with the area of  $>10^4$  MSH (micro solar hemispheres;  $1 \text{ MSH}=3 \times 10^{16} \text{ cm}^2$ ) on slowly-rotating solar-type stars with that of sunspot groups. The size distribution of starspots shows the power-law distribution and that of larger sunspots lies on the same power-law line. The size distribution of spots from the Kepler data suggests that the average appearance frequency of the starspots with the area of  $>3 \times 10^4$  MSH on the solar-type stars with the rotation period similar to that of the Sun is once in a few hundred years.

We also found that the frequency-energy distributions for flares originating from spots with different sizes are the same for solar-type stars with superflares and the Sun. These results suggest that the magnetic activity on solar-type stars with superflares and that on the Sun is caused by the same physical processes.

キーワード：フレア、黒点

Keywords: flare, starspot

## 太陽フレアのimpulsivityと白色光フレア・CMEとの関係

### Solar flare impulsivity and its relationship with white-light flares and with CMEs

\*渡邊 恭子<sup>1</sup>、有馬 伸<sup>1</sup>、堀 巖允<sup>1</sup>、増田 智<sup>2</sup>

\*Kyoko Watanabe<sup>1</sup>, Shin Arima<sup>1</sup>, Yoshimasa Hori<sup>1</sup>, Satoshi Masuda<sup>2</sup>

1. 防衛大学校、2. 名古屋大学宇宙地球環境研究所

1. National Defense Academy of Japan, 2. Institute for Space-Earth Environmental Research (ISEE), Nagoya University

There are many types of classification in solar flares. One of them is a classification by flare duration in soft X-rays; so-called impulsive flare and long duration event (LDE). Typically, the duration of an impulsive flare is shorter than 1 hour, and that of an LDE is longer than 1 hour. These two types of flare show different characteristics. In soft X-rays, impulsive flares usually have a compact loop structure. On the other hand, LDEs show a relatively large-scale loop, sometimes a large arcade structure. In hard X-rays, the difference appears clear, too. The former shows a strong and short-time (<10 minutes) emissions and intense double footpoint sources and sometimes a weak loop-top source. LDEs have relatively weaker and longer (>10 minutes) emissions and show a large coronal source. These facts suggest that hard X-ray observation becomes one of a good indicator to classify solar flares, especially for the study on the particle acceleration and the related phenomena. However, hard X-ray data do not always exist due to the satellite orbit and the small sensitivity of hard X-ray instruments. So, in this study, based on the concept of the Neupert effect (Neupert, 1968), we use soft X-ray derivative data as the proxy of hard X-ray. From this data, we define impulsivity (IP) for each flare. Then we investigate solar flares using this new index. First we apply IP index to our white-light flare research. We have already performed a statistical analysis of white-light flare to reveal the physical conditions to generate white-light enhancements. We investigate how white-light enhancement depends on IP, then it is found that white-light flares tend to have large IP values. So the flare impulsivity (IP) is one of the important factors if white-light enhancement appears or not in a solar flare. Next we investigate how CME physical parameters depend on IP index. It has been believed that most of CMEs are associated with LDEs, but we found that there is only a weak correlation between the existence of CME and IP index. We couldn't find any relationship between CME physical parameters (speed, mass, energy) and IP. Finally, we also search for the relationship between white-light flare and CME as a function of IP and discuss the physical condition of white-light flare.

キーワード：太陽フレア、白色光、CME

Keywords: solar flare, white-light, CME

# The Relation between Time-Series Characteristics of Solar Active Region Magnetic Field Properties and Flare Activity

\*Eo-Jin Lee<sup>1</sup>, Yong-Jae Moon<sup>1,2</sup>, Sung-Hong Park<sup>3</sup>

1. School of Space Research, Kyung Hee University, 2. Department of Astronomy and Space Science, Kyung Hee University, 3. Trinity College Dublin

Solar flares suddenly release an immense amount of energy mainly in the form of electromagnetic radiation, and consequently they can cause major disturbances in the Sun-Earth environment. An accurate prediction of flares is therefore crucial for protecting our space assets and the safety of astronauts and navigation systems. In this study, we investigated a statistical data set of 8-day time series of 18 magnetic field parameters in 94 flaring solar active regions (ARs) in 2010 to 2016. The time series data were derived from 12-min cadence photospheric vector magnetograms observed by SDO/HMI. For each AR under investigation, we calculated the mean values of the magnetic parameters from their corresponding time series data, as well as the fluctuations using detrended fluctuation analysis. Then, the mean values and fluctuations were compared with the sum of peak soft X-ray intensities of flares produced in the investigated AR (called flare index) during the eight days. As a result, we find the mean of time series of parameters measuring magnetic non-potentiality of ARs, such as vertical current, current helicity and free magnetic energy density, have a good correlation with the flare index: i.e., their correlation coefficients (CCs) are higher than 0.58. Especially, the mean of AR vertical current time series has a fairly good correlation with the flare index (CC=0.64). It is also found that the fluctuation of AR current helicity time series is well correlated with the flare index (CC=0.68). These results show that the mean and fluctuation of time series of AR non-potential magnetic parameters can be considered as a useful measure for estimating AR' s flaring activity over the entire time series. Moreover, it is expected that further time series analysis can help improve flare prediction.

Keywords: Sun, Solar Flares, Time Series Analysis

# Solar Flare Prediction using the Machine-learning and Operational Evaluation Method

\*西塚 直人<sup>1</sup>、杉浦 孔明<sup>1</sup>、久保 勇樹<sup>1</sup>、石井 守<sup>1</sup>

\*Naoto Nishizuka<sup>1</sup>, Komei Sugiura<sup>1</sup>, Yuki Kubo<sup>1</sup>, Mamoru Ishii<sup>1</sup>

1. 国立研究開発法人 情報通信研究機構

1. National Institute of Information and Communications Technology

We have developed a flare prediction model using solar observation data and machine-learning techniques. From the full-sun images, we extracted features such as photospheric magnetic field, chromospheric brightening and X-ray activities in each active region, and then we predicted the maximum class of flares occurring in the following 24 hours (Nishizuka et al., 2016, oral presentation in JpGU). However, a standard evaluation method of flare prediction models has not been established. Moreover, under the sever condition that the test dataset is completely independent from the training dataset in an operational setting, we could not predict solar flares with high accuracy.

In this presentation, we introduce a method of the time-series cross-validation (CV) to evaluate flare prediction models in an operational setting, though the k-fold (10-fold) CV has ever been used in the previous studies. In some sense, these two methods are reasonable and available. However, when we focus on the operational usage, the time-series CV is superior to the k-fold CV. Furthermore, we used a machine-learning algorithm called the Gradient Boosted Trees for the first time. The boosting is a method to minimize the loss function by sequentially adding weak classifiers, or decision trees in our model. This is used to achieve a better prediction, by repeating learning of the calculation of the gradient when optimizing parameters in each step. We applied this algorithm to the flare prediction and performed the time-series CV. As a result, we succeeded in improving our prediction score, a skill score called the true skill statistic, from 0.2 to 0.6 for X-class flares and to 0.8 for M-class flares. We also compared the performance of other five different machine-learning algorithms to predict flares, and we found that the ranking of the performance of the algorithms completely differs according to the CV method.

キーワード：太陽フレア、予測、モデリング、機械学習、評価手法

Keywords: Solar Flare , Prediction, Modeling , Machine-learning, Evaluation Method

# Energy Budget in Cold Solar Flares Observed with Nobeyama Radioheliograph

\*増田 智<sup>1</sup>

\*Satoshi Masuda<sup>1</sup>

1. 名古屋大学宇宙地球環境研究所

1. Institute for Space-Earth Environmental Research, Nagoya University

The energy source of solar flare is magnetic energy in the solar corona. When a solar flare takes place, this energy is impulsively released, and it is converted to other kinds of energy such as thermal energy (plasma heating), kinetic energy (mass ejection), and energy of high-energy particles (particle acceleration). The ratio of these three energies seems different in each solar flare. Some solar flares show thermal energy is dominant and the others do non-thermal energy of accelerated particles is dominant. What controls the energy conversion ratio? This is so-called energy budget problem in solar flares.

A hint to solve this problem could be provided by some extreme flares such as thermal-rich flares and purely nonthermal flares. In this study, we focused on so-called cold flare which shows significant nonthermal emissions with relatively very small amount of thermal emissions. We have already analyzed a cold solar flare occurring on 10 March 2011. This flare was well observed with Nobeyama Radioheliograph (NoRH). It shows a significant microwave enhancement (about 210 SFU at 17GHz). This microwave enhancement is produced via gyro-synchrotron emissions by high-energy electrons. However, any significant enhance was not found in the GOES X-ray light curve during the flare period. RHESSI did not observe this flare, but Suzaku detected hard X-rays in the energy range above 100 keV. So it was confirmed that electrons were actually accelerated while plasma heating process did not work well in this flare. The characteristics of this flare were short duration and very compact size. From these facts, we concluded that the magnetic field of the loop top region might be very intense. This might be a key how a cold flare is produced. To confirm this, we need more analyses of similar events. From the event list of NoRH, we pick up candidates of cold flares which show a significant microwave enhancement without any counterpart in soft X-rays and analyze them. The characteristics seem to be the same as the previous event. Then we discuss why thermal emissions (plasma heating) are suppressed in these flares.

キーワード : 太陽フレア、電波、粒子加速

Keywords: Solar flare, radio, particle acceleration

## Statistical analysis of solar flare multi-wavelength observation data

\*西本 将平<sup>1</sup>、渡邊 恭子<sup>1</sup>、今田 晋亮<sup>2</sup>、川手 朋子<sup>3</sup>、Lee Kyoung Sun<sup>4</sup>

\*Shohei Nishimoto<sup>1</sup>, Kyoko Watanabe<sup>1</sup>, Shinsuke Imada<sup>2</sup>, Tomoko Kawate<sup>3</sup>, Kyoung Sun Lee<sup>4</sup>

1. 防衛大学校、2. 名古屋大学宇宙地球環境研究所、3. 宇宙科学研究所、4. 国立天文台

1. National Defense Academy, 2. Institute for Space-Earth Environmental Research, Nagoya University, 3. Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 4. National Astronomical Observatory of Japan

Impulsive energy release phenomena such as solar flares, sometimes affect to the solar-terrestrial environment. Usually, we use soft X-ray flux as the index of flare scale. However, the magnitude of effect to the solar-terrestrial environment is not proportional to the GOES X-ray class. To identify the relationship between solar flare phenomena and influence to the solar-terrestrial environment, we need to understand the full spectrum of solar flares. For this purpose, we are performing statistical analysis of electromagnetic data of solar flares. In this study, we use solar flare events larger than C3-class from the Hinode flare catalogue (Watanabe et al., 2012). We use the soft X-ray flux, flare duration, location, and type and size of flaring active region as the fundamental parameters of a solar flare. For the first step, we focus on the flux of EUV emission, because the sudden ionospheric disturbance(SID) is caused by a solar flare especially due to soft X-ray and UV emissions. We use EUV data from the Extreme Ultraviolet Variability (EVE) onboard the Solar Dynamics Observatory (SDO). We examined the EUV lines (Fe VIII, Fe XII, Fe XV, Fe XVIII, Fe XX) for 111 events larger than M3-class flares, and found positive correlation between the “soft X-ray flux” and the “EUV peak intensity” for all lines. The “EUV peak time” of hot lines are earlier than that of cool lines. Next, we examined the relationship between the size of flaring active regions and the flare intensity. We found very weak correlation between them when we plot 5578 flare events larger than C-class from November 2006 to July 2016. In addition, we also check the occurrence timing of flare events during evolution or dissipation of active region, and found that there is no relationship between them. Then, we also examined the hard X-ray data obtained by Reuven Ramaty High Energy Solar Spectroscopic Imager (RHESSI). We also found correlation between the hard X-ray intensity and the soft X-ray intensity, especially in low energy. In this paper, we also show the spectral index of hard X-ray spectra, and discuss their relationship for other electromagnetic emissions.

キーワード : 太陽フレア、太陽フレアスペクトル

Keywords: solar flare, solar flare spectra

## Study of double arc instability causing the onset of solar eruption

\*石黒 直行<sup>1</sup>、草野 完也<sup>1</sup>

\*Naoyuki Ishiguro<sup>1</sup>, Kanya Kusano<sup>1</sup>

1. 名古屋大学 宇宙地球環境研究所 総合解析研究部

1. Division for Integrated Studies, Institute for Space-Earth Environmental Reserch, Nagoya University

The stability of magnetic flux rope in the solar corona is an important issue for the understanding what causes solar eruptions, such as solar flares and coronal mass ejections (CMEs). Although various scenario has been suggested for this problem, the tether-cutting reconnection scenario suggested by Moore et al. (2001) is widely accepted. However the stability of double arc loop, which can be formed by tether-cutting reconnection of sheared magnetic field before eruption, has not been analyzed yet. To explain such a solar eruption based on the tether-cutting reconnection model, recently we suggested Double Arc Instability (DAI). In this theory, we numerically analyzed the stability of double arc electric current loop in the same method as the axisymmetric torus instability developed by Demoulin & Aulanier (2010). As a result, we found that the double arc loop is more easily destabilized than the axisymmetric torus, and that it becomes unstable even if the external field is not decayed with altitude in contrast to the axisymmetric torus instability. It suggests that the tether-cutting reconnection can efficiently work as the onset mechanism of solar eruption, and that the decay index does not necessarily need to be larger than a criteria to trigger solar eruptions, while the torus instability can play a role in the full eruption of the flux rope to inter-planetary space. We discuss the critical conditions for the DAI to grow in the force-free magnetic field and derive that the critical parameter, which is defined as the product of the magnetic twist and the normalized tether-cutting reconnecting flux, must be larger than a certain threshold to destabilize DAI. Finally, we develop three-dimensional MHD simulation of the DAI to analyze the onset process of solar eruption in more realistic configuration of magnetic field.

キーワード：太陽、太陽フレア、不安定性

Keywords: Sun, flares, instability

## The statistical analysis of correlation between solar flares and photospheric magnetic field

\*旭 友希<sup>1</sup>、草野 完也<sup>1</sup>、伴場 由美<sup>2</sup>、塩田 大幸<sup>1</sup>、柴山 拓也<sup>1</sup>

\*Asahi Yuki<sup>1</sup>, Kanya Kusano<sup>1</sup>, Yumi Bamba<sup>2</sup>, Daikou Shiota<sup>1</sup>, Takuya Shibayama<sup>1</sup>

1. 名古屋大学宇宙地球環境研究所、2. 宇宙航空研究開発機構

1. Institute for Space-Earth Environmental Research , 2. The Japan Aerospace Exploration Agency

A solar flare is caused by the explosive release of magnetic energy and sometimes greatly disturbs the Earth's electromagnetic environment and may impact socio-economic system. For that reason, the prediction of flare occurrence is important for space weather forecast. However, the accurate prediction of flare occurrence has not been realized yet, because the mechanism of flare trigger is not well understood yet. According to previous studies, some parameters, such as the area of active region, total magnetic flux, the magnetic shear on PIL and the magnetic gradient have certain relation with the flare activities. Recently, Kusano et al. (2012) indicated that the magnetic reconnection in the small magnetic structures that appear in the strong-sheared magnetic field near the magnetic polarity inversion lines can trigger solar flares. In addition, Ishiguro & Kusano (M25a for ASJ meeting 2016 in March) found a possibility that the magnetic twist causes instability which is able to cause flares and CMEs. The object of this study is to give a new parameter related to flare activity on the basis of these previous studies. In order to achieve it, we have performed the statistical analysis of magnetic field data on photosphere surface. For 294 ARs which sunspot area is relatively large in 2012 to 2016, we took the correlation analysis on the total flux and various magnetic parameter in each AR using the magnetic field data of SDO/HMI. The results suggest that not only magnetic free energy but also the shape of distribution of magnetic free energy correlates with the flare activity. We report the preliminary results of the analyses and discuss about the application to space weather forecast.

キーワード : 太陽フレア、宇宙天気、SDO/HMI

Keywords: Solar flare, Space weather, SDO/HMI



# Investigation of meridional flow pattern from magnetic elements motion

\*Yusuke Iida<sup>1</sup>

1. Kwansai Gakuin University

Meridional flow, which is considered a persistent flow pattern along North-South direction, is important for solar dynamo problem. Despite its importance, observation of the meridional flow pattern is difficult because the amplitude is just  $\sim 10$  m/s, which is just a few percent of that of convective flow pattern covering the solar surface. Hathaway and Lightmire (2010) succeeded in detecting the meridional flow pattern as an advection of magnetic region patterns between consecutive magnetograms obtained by SoHO/MDI. However, their study is limited to investigate the averaged pattern of magnetic region, not each magnetic element. Hence we try detecting the meridional flow pattern in the magnetic elements' motion in this study, which is more direct detection and enable us to investigate deeper character of the flow.

North-south anisotropy in magnetic elements' motion is investigated in the magnetograms obtained by Helioseismic and Magnetic Imager (HMI) onboard Solar Dynamics Observatory (SDO). The magnetic elements are tracked by feature-tracking method of magnetic concentrations which is developed by the author. Our feature-tracking method and one-month magnetograms of SDO/HMI enable us to analyze huge number of magnetic elements' motion. More than 5,000 elements are recognized in one magnetogram with our detecting threshold and  $10^8$  displacements are obtained as total in our analysis. We found a small anisotropy along north-south direction and it shows similar latitudinal dependence obtained in Hathaway and Lightmire (2010), namely its amplitude increases up to  $\sim 14$  m/s from the equator to  $\sim 60$  degree in north and south hemisphere. Further we investigate the dependence of flow strength on magnetic elements character and newly found that larger elements have larger flow strength.

Keywords: Sun, Magnetic Field, Surface Flow

## 活動領域NOAA12297における黒点の逆回転とX2.1フレア Reversed Rotation of the Sunspot and the X2.1 Flare in the Active Region NOAA12297

\*長谷川 隆祥<sup>1,2</sup>、清水 敏文<sup>2,1</sup>

\*Takahiro Hasegawa<sup>1,2</sup>, Shimizu Toshifumi<sup>2,1</sup>

1. 東京大学、2. 宇宙航空研究開発機構宇宙科学研究所

1. The University of Tokyo, 2. Institute of Space and Astronautical Science, JAXA

我々は、活動領域NOAA12297におけるX2.1クラスフレア前後の磁場構造の、2015年3月10日から13日までの時間発展について研究した。本活動領域の主要な黒点は、3/10の0時ごろの段階では時計回りの回転をしていたが、3/10、22時ごろに現れた急速な磁気浮上によるシア流れによって反時計回りの回転が促された。磁気浮上が進むにしたがって黒点に発生するローレンツトルクも増加し、黒点の回転速度は上昇していき、最大で $\sim 2.5$ deg/hに達した。これに伴い、磁場の非ポテンシャル性を表す物理量(平均フォースフリー $\alpha$ )も増大して行った。さらに、黒点の逆回転は、大局的に正の磁気ヘリシティを持つ本活動領域に、大きな負の磁気ヘリシティをもたらしていた。3/11の16時ごろ、この黒点のすぐ北でX2.1クラスフレアが発生した。そのすぐ後、黒点の回転速度は減少を始め、3/13の0時ごろには時計回りに戻っていた。この解析結果から、我々は、Xクラスほどの巨大フレアの自由エネルギー蓄積には、黒点付近の磁気浮上に加えて、黒点の回転が重要な役割を持つことを主張する。またこの結果は、系のもつ大局的な磁気ヘリシティに対して逆向きの磁気ヘリシティがもたらされることが、フレアのオンセットに寄与する可能性を示唆する。

キーワード：太陽、フレア、黒点、エネルギー蓄積、磁気ヘリシティ

Keywords: Sun, Flare, Sunspot, Energy build-up, Magnetic helicity

## Quantitative relationship between Mid-latitude ionospheric ExB drift and equatorial electrojet as a function of solar zenith angle

\*橋本 久美子<sup>1</sup>、菊池 崇<sup>2</sup>、富澤 一郎<sup>3</sup>、長妻 努<sup>4</sup>

\*Kumiko Hashimoto<sup>1</sup>, Takashi Kikuchi<sup>2</sup>, Ichiro Tomizawa<sup>3</sup>, Tsutomu Nagatsuma<sup>4</sup>

1. 吉備国際大学、2. 名古屋大学宇宙地球環境研究所、3. 電気通信大学宇宙・電磁環境研究センター、4. 情報通信研究機構  
1. Kibi International University, 2. Institute of Space-Earth Environmental Research, 3. University of Electro-Communications, 4. National Institute of Information and Communications Technology

In order to estimate the penetration electric fields in the mid-latitude ionosphere from the equatorial electrojet (EEJ), we analyzed stormtime global Pc5 magnetic pulsations. During the recovery phase of the geomagnetic storm on 31 October 2003, the HF Doppler sounder detected large amplitude oscillations in the ionospheric electric field at mid-latitude (Oarai and Sugadaira, Japan) over 10 hours from 11 to 21 LT. Prolonged excitation of Pc5 pulsations were recorded on the magnetometer data at high-to-equatorial latitudes with significant amplitude enhancement at the dayside equator. We deduced the EEJ by subtracting the low latitude Pc5 (Okinawa) from the equatorial Pc5 (Yap) and found that the ExB drift velocity ( $V_{\text{vert}}$ ) in the mid-latitude is well correlated with the EEJ with correlation coefficients of 0.80-0.95 and that the EEJ to  $V_{\text{vert}}$  ratio reached maximum at 11 LT and dramatically decreased until 18 LT as a function of  $\cos(\text{solar zenith angle})$  for the zenith angle less than 92 degrees. The quantitative relationship is evaluated by estimating the root mean square error of 2.5 m/s for afternoon sector and 1.8 m/s for dusk sector, which is small enough compared to the observed  $V_{\text{vert}}$  of 15-40 m/s. On the other hand, the electric field at mid-latitude is well correlated with the EEJ even during the night after 19 LT (zenith angle > 100 degrees), showing that the EEJ to mid-latitude  $V_{\text{vert}}$  ratio is almost constant, on average 1.65. This result indicates that the Pc5 electric field is so strong as to drive equatorial electrojet in the nighttime ionosphere. By applying the quantitative relationship to other longitudes, such as Sao Luis in Brazil, Addis Ababa in Ethiopia, we estimated a local time distribution of the electric field at the mid-latitude and obtained the day-night asymmetry of the electric field.

キーワード：電離圏電場、赤道ジェット電流、ExBドリフト速度、中緯度電離圏、HFドップラー

Keywords: ionospheric electric field, equatorial electrojet, ExB drift velocity, mid-latitude ionosphere, HF Doppler

## On the relationships between EEJ distribution and plasma bubble occurrences

\*秋山 鷹史<sup>1</sup>、吉川 顕正<sup>2</sup>、松下 拓輝<sup>1</sup>、藤本 晶子<sup>3</sup>、魚住 禎司<sup>3</sup>

\*Takafumi Akiyama<sup>1</sup>, Akimasa Yoshikawa<sup>2</sup>, Hiroki Matsushita<sup>1</sup>, Akiko Fujimoto<sup>3</sup>, Teiji Uozumi<sup>3</sup>

1. 九州大学大学院理学府地球惑星科学専攻宇宙地球電磁気学分野、2. 九州大学大学院理学研究院地球惑星科学部門、3. 九州大学国際宇宙天気科学・教育センター

1. Field of Space and Earth Electromagnetism, Department of Earth and Planetary Sciences, Graduate School of Sciences, Kyushu University, 2. Department of Earth and Planetary Sciences, Kyushu University, 3. International Center for Space Weather and Education, Kyushu University

To understand the relationship between distributions of Equatorial Electro Jet (EEJ) along dayside dip-equator and generation characteristic of plasma bubble, we investigate how Local Time (LT) dependence of EEJ amplitude is relating to the plasma bubble occurrences. The LT dependence of the amplitude of EEJ is characterized by using integrated value of EUEL of EE-index (produced by ICSWSE, Kyushu University) at daytime (07:00~17:00LT), noon time (09:00~15:00LT) and evening time (17:00~19:00LT). While plasma bubble occurrence is determined by S4 index of ionospheric scintillation produced by ISEE, Nagoya University. In this study, EUEL data at equatorial station: Langkawi, Malaysia (GG Lon. =99.78, Dip Lat. =-1.07), and S4 data at Kototabang, Indonesia (GG Lon. =100.32, Dip Lat. =-10.1) from 1 January 2011 to 8 November 2014 are used to investigate the relationship between EEJ and the scintillation. Our result shows that there are relationship between plasma bubble occurrence and integrated EUEL value in all the above period; larger integrated EUEL value, larger occurrence rate of plasma bubble. Moreover it turned out that plasma bubble tends to be suppressed when integrated EUEL value in the evening is negative. Our result for evening side is consistent with previous study of Uemoto et al., [2010], while the one for daytime does not. This might be due to different way of EEJ amplitude estimation was used. In our study, EUEL value at Langkawi not only reflects EEJ amplitude but also background Sq disturbances. While in the study of Uemoto et al., [2010], EEJ amplitude is estimated by purely equatorial enhancement and/or equatorial depression component. Our results strongly suggested that plasma bubble occurrence is not only controlled by the Equatorial enhancement but also Sq structure near dip equator.

キーワード：赤道ジェット電流、プラズマバブル

Keywords: Equatorial ElectroJet, Plasma Bubble

## Solar cycle variation and its impact on Critical Frequency of F layer

\*Pramod Kumar Purohit<sup>1</sup>, Roshni Atulkar<sup>1</sup>, Azad A. Mansoori<sup>2</sup>, Parviaz A. Khan<sup>3</sup>

1. National Institute of Technical Teachers' Training and Research, Bhopal -462002, MP, India, 2. Department of Electronics, Barkatullah University, Bhopal -462026, MP, India, 3. Department of Electronics and Communication Engineering, Islamic University of Science and Technology, Pulwama, J & K, India

The period of approximately 11 year cycle of solar activity is characterized by the rise and fall in the numbers and surface area of sunspots. We observed a number of other solar activity indices, including the 10.7 cm radio flux, solar Mg II core to wing ratio, relative sunspot number  $R_z$  and solar flare index and geomagnetic activity that vary in association with the sunspots for solar cycles 21, 22 and 23 (1976–2008). This paper presents an analysis of the F-region variability of the ionospheric parameter foF2 at mid latitude station Hobart (Hobart is a town in County Australia) Latitude:- 42.8806° S and Longitude: 147.3250° E during in the whole period (1976–2008) of solar cycle-21, 22, and 23. The diurnal, monthly, yearly and cycle to cycle characteristics of these ionospheric F-region parameter foF2 have been studied in detail. We also compared the dependence of foF2 on solar activity indices by using a correlation analysis, and showed that a significant linear relationship between the foF2 values and Solar indices. The foF2 variation is strongly influenced by solar activity with about an 11-year solar cycle from the solar maximum to solar minimum.

Keywords: Solar cycle, foF2, , geomagnetic indices, correlation

## Analysis of the ISEST/MiniMax24 WG4 campaign events on the linkage between CMEs and solar wind disturbances

\*丸橋 克英<sup>1</sup>、Cho Kyungsuk<sup>2</sup>、石橋 弘光<sup>1</sup>

\*Katsuhide Marubashi<sup>1</sup>, Kyungsuk Cho<sup>2</sup>, Hiromitsu Ishibashi<sup>1</sup>

1. 情報通信研究機構、2. 韓国天文学宇宙科学研究所

1. National Institute of Information and Communications Technology, 2. Korea Astronomy and Space Science Institute

The ISEST/MiniMax24 is one of the four projects of VarSITI. Its objectives and goals are to understand the propagation of solar transients through the space between the Sun and the Earth, and develop space weather prediction capability. Toward the goals the ISEST/MiniMax24 Working Group 4 organized a campaign study by selecting 11 solar-terrestrial events. We examined the solar wind data during each period of the campaign events and identified flux rope structure in 9 cases. Then geometries of 9 interplanetary flux ropes (IFRs) were determined by a model-fitting method and compared them with magnetic structures in their solar source regions. As a result, we could confirm the general coincidence between the IFR orientations and the orientations of the polarity inversion lines (PILs) in the corresponding solar source regions. The results are summarized in Table 1. This analysis result suggests a possibility for predicting variations of solar wind magnetic fields associated with flux ropes basically by observing the causative solar eruptions. However, we found several cases in which prediction needs the precise knowledge about the flux rope structure and where the Earth encounters it with a flux rope. We report the relationships between the IFRs and the corresponding PILs with emphasis upon how the observed magnetic fields are determined by the encountering geometries.

キーワード：磁気ロープ、コロナガス噴出、磁気極性反転線

Keywords: magnetic flux rope, coronal mass ejection, polarity inversion line

Table 1

Event ID	model	IP Flux Rope		Solar Source		S/C
		R/L	IFR tilt	N/S	PIL tilt	
1. 2012 Jul 12-14	cylinder	R	320°	S	325°	WIND
2. 2012 Oct 04-08	torus	R	323°	S	330°	WIND
3. 2013 Mar 15-17	torus	L	227°	N	230°	ACE
4. 2013 Jun 01	torus	L	272°	N	—	ACE
5. 2015 Mar 15-17	torus	R	173°	S	165°	ACE
6. 2015 Jun 22-24	Special Analysis underway					A/W
7. 2012 Mar 07-09	torus	L	37°	N	42°	WIND
8. 2012 Jul 23-24	cylinder	R	258°	S	(260°)	STEREO-A
9. 2014 Jan 06	No flux rope signature (Limb event)					A/W
10. 2014 Jan 07-09	No flux rope signature (Deflected, Möstl)					A/W
11. 2014 Sep 10-13	torus	L	247°	N	245°	

## Statistical characteristics of interplanetary magnetic field near the Earth

\*Kyung-Eun Choi<sup>1</sup>, Dae-Young Lee<sup>1</sup>, Kyu-Cheol Choi<sup>2</sup>, Jaehun Kim<sup>3</sup>

1. Chungbuk National University, Department of Astronomy and Space Science, Cheongju, Chungbuk, Korea, 2. Research Institute, SELab, Inc., Korea, 3. Korean Space Weather Center, National Radio Research Agency, Korea

Space weather creates geomagnetic disturbances that can impact our life. There are many reasons that cause space weather changes, and the interplanetary magnetic field (IMF) is the most important one. Compared with other components, IMF Bz is an immediate cause as it is a parallel component to the geomagnetic field. Intense geomagnetic disturbance can be made by magnetic reconnection which can occur more strongly when the magnitude of southward IMF Bz is stronger. Nevertheless, until to date, it has not been trivial to predict IMF Bz. In this work we analyze the statistical properties of IMF Bz measured at L1 point using the data from 1996 to 2015 corresponding to a solar cycle and a half. In this study, we classify the IMF data into three groups. They are IMF conditions associated with CME, those with CIR+HSS (high-speed stream) and the remaining ordinary conditions. We find that a majority (84%) of the IMF data are during the ordinary conditions, only 5% are related to CME structure and ~11 % are associated with CIR+HSS. In order to study geoeffectiveness, we have examined relativistic electron (>2MeV) flux response at geosynchronous orbit and compared them among the three groups. We have also investigated AL index. We find that the geomagnetic response in terms of AL and geosynchronous electron flux is non-negligible during the ordinary solar wind condition intervals compared to those of CME and CIR+HSS. In addition, we examine the correlations of IMF Bz with various solar wind parameters to determine the most responsible factors for southward IMF Bz.

# Numerical Study of Double-Cell-Type Solar Meridional Circulation Based on a Mean-Field Hydrodynamic Model

\*戸次 宥人<sup>1</sup>、横山 央明<sup>1</sup>

\*Yuto Bekki<sup>1</sup>, Takaaki Yokoyama<sup>1</sup>

1. 東京大学大学院理学系研究科

1. Graduate School of Science, The University of Tokyo

Large-scale flow structures of the Sun, differential rotation and meridional circulation, play crucial roles in generating and sustaining the solar magnetic fields through the dynamo mechanism. In the framework of flux-transport dynamo model, most of the previous kinematic simulations have assumed a single-cell meridional circulation so that equator-ward migration of sunspot groups could be attributed to the equator-ward transport of toroidal magnetic fluxes by the meridional flow at the base of the convection zone. However, recent helioseismic observational results suggest the possibility of a double-cell structure for the meridional circulation with the pole-ward flow at the base, demanding some modifications for the conventional flux transport dynamo model. Therefore, the theoretical investigations on the maintenance mechanism of this double-cell meridional circulation is regarded as of a great importance.

By conducting mean-field hydrodynamic simulations where the effect of the angular momentum transport by the Reynolds stress is parameterized, we calculate the structures of differential rotation and meridional circulation self-consistently and investigate whether or not double-cell meridional circulation could be achieved along with the solar-like differential rotation. As a result, we find out that the double-cell meridional circulation can be achieved when the Reynolds stress transports angular momentum upward in the lower convection zone and downward in the upper layer. We confirm that, in the steady state, the accumulated angular momentum via the Reynolds stress in the middle layer is advected to both the upper and lower part of the convection zone by each of the upper and lower meridional circulation cells, respectively.

キーワード：太陽、対流

Keywords: The Sun, Convection



## Results from the daily solar segmentation (SOLARSEG) of EUV images over a 5-years period: correlation and periodicity of the EUV irradiance.

\*Gabriel GIONO<sup>1</sup>, Joe ZENDER<sup>2</sup>, Rangaiah KARIYAPPA<sup>3</sup>, Matthias BERGMANN<sup>4</sup>, Veronique DELOUILLE<sup>5</sup>, Luc DAME<sup>6</sup>, Jean-François HOCHEDÉZ<sup>6</sup>, S.T. KUMARA<sup>7</sup>


1. KTH, 2. ESA/ESTEC, 3. Vemana Institute of Technology, 4. Julius-Maximilians-Universität Würzburg, 5. Royal Observatory of Belgium,, 6. LATMOS, 7. APS College of Engineering

Understanding the variability of the EUV and UV irradiance on time scales from hours to months is of critical importance in the context of the Sun-Earth connection. A promising method to study the impact of the different coronal features, such as active regions and coronal holes, is by using segmentation algorithm on full-disc EUV images. In this respect, we created the Solar Segmentation (SOLARSEG) pipeline, based on the Spatial Possibilistic Clustering Algorithm (SPoCA) segmentation algorithm, which has been continuously producing daily maps (4h cadence) of the active regions (ARs) and coronal holes (CHs) using EUV images from the Atmospheric Imaging Assembly (SDO/AIA) starting from January 2011. The segmented maps were used to extract the EUV irradiance from these regions of interest using the different wavelengths provided by the AIA instrument. Initial comparison with the full-disk EUV irradiance recorded by the Large Yield Radiometer (PROBA2/LYRA) indicated a strong correlation with the irradiance extracted from EUV images, proving the usefulness of image segmentation to estimate the EUV irradiance from the different features. Maps were also be applied to other imaging instrument, namely the Sun Watcher with Active Pixel System detector and Image Processing (PROBA2/SWAP) and the Helioseismic and Magnetic Imager (HMI), to respectively compare the recorded irradiance around 17.1 nm and discuss the underlying magnetic field.

Although the end-goal of the project is to discuss the variability of the EUV irradiance over a full solar cycle period, current results for a 5-years period already contain interesting results. For example, a strong correlation was found between the coronal irradiance and the photospheric magnetic field. This correlation was expected for active regions, but was also found in the quiet sun region (i.e. region not included in either the ARs maps or the CHs maps). Such result, combined with one of our previous results estimating the contribution of the quiet sun region to more than 60% of the total EUV irradiance, suggests that the photospheric magnetic field has a more global impact on the EUV irradiance than expected, indicating its importance in understanding and predicting the EUV irradiance variability.

The present poster discusses in more details the correlation and periodicity found between the irradiance from the different atmospheric layers observed by the AIA EUV bands and the underlying magnetic field over a 5-years period, from 2011 to 2016.

Keywords: Solar physics, EUV irradiance variability



## Identification of Active Longitude from the Solar Magnetograms

\*近藤 克哉<sup>1</sup>、今田 晋亮<sup>1</sup>、藤山 雅士<sup>1</sup>、町田 忍<sup>1</sup>

\*Katsuya Kondo<sup>1</sup>, Shinsuke Imada<sup>1</sup>, Masashi Fujiyama<sup>1</sup>, Shinobu Machida<sup>1</sup>

1. 名古屋大学宇宙地球科学研究所

1. Institute for Space-Earth Environmental Research

It is well known that the sun has the cycle of 11 years for its activity. In the active time, we can see lots of sunspots on the solar surface and sometime it cause large flares which largely affect the Earth's environments. For example, the release of large-scale plasma called Coronal Mass Ejection (CME) hits the Earth's magnetosphere and causes some effects, e.g. the Earth's magnetic field fluctuation or trouble of the satellite. The appearance of sunspots has several characteristics. For example, sunspots appear around in the higher latitudes at the beginning of the cycle and in the lower latitudes at the end of the cycle on the sun. However, recent studies have shown that the appearances of sunspots also depend on longitudinal direction. The solar longitude where the sunspots are frequently observed is called Active Longitude (AL). The presence of active longitude has been discussed long time. But it is still not clear whether there is an AL or not. In this study, we identified AL using magnetic field observation on the solar surface, not sunspots data. As a result, we can study AL for the magnetic bipoles which do not develop to sunspots. The results show that the phase of AL in the northern hemisphere and the southern hemisphere are different in cycles 23 and 24. Further, we found that there is clear AL in the strong magnetic field data ( $|B| > 500\text{G}$ ), although there is no clear AL in the weak magnetic field data ( $|B| < 40\text{G}$ ).

キーワード：黒点、太陽

Keywords: Active Longitude, Sunspot

## Cosmogenic $^{10}\text{Be}$ in endogenic travertine deposits at Baishuitai, China: A pilot study

\*宮原 ひろ子<sup>1</sup>、坂下 渉<sup>1</sup>、堀内 一穂<sup>2</sup>、周 立旻<sup>3</sup>、徐 洪陽<sup>3</sup>、松崎 浩之<sup>4</sup>

\*Hiroko Miyahara<sup>1</sup>, Wataru Sakashita<sup>1</sup>, Kazuho Horiuchi<sup>2</sup>, Limin Zhou<sup>3</sup>, Xu Hongyang<sup>3</sup>, Hiroyuki Matsuzaki<sup>4</sup>

1. 武蔵野美術大学、2. 弘前大学、3. 華東師範大学、4. 東京大学

1. Musashino Art University, 2. Hirosaki Univ., 3. East China Normal Univ., 4. The Univ. of Tokyo

We report on the preliminary experiment for measurement of beryllium-10 in travertine samples from Baisuitai, China. Beryllium-10 is one of the cosmic-ray induced isotopes and thus can be used as a proxy for the flux of galactic cosmic rays incident on the Earth. For example, the content of beryllium-10 in ice cores from polar region is often used for examining the past cosmic ray flux and solar variations. However, the data from ice cores is often accompanied by dating uncertainty. In this study, we seek for the possibility to use travertine which has recognizable annual layers.

キーワード：宇宙線生成核種、太陽活動周期、トラバーチン堆積物

Keywords: Cosmogenic nuclide, Solar cycle, Travertine deposit

## Solar 27-day rotational period detected in a wide-area lightning activity in Japan

\*宮原 ひろ子<sup>1</sup>、寺澤 敏夫<sup>2</sup>、片岡 龍峰<sup>3</sup>、佐藤 光輝<sup>4</sup>、高橋 幸弘<sup>4</sup>

\*Hiroko Miyahara<sup>1</sup>, Toshio Terasawa<sup>2</sup>, Ryuho Kataoka<sup>3</sup>, Mitsuteru Sato<sup>4</sup>, Yukihiro Takahashi<sup>4</sup>

1. 武蔵野美術大学 教養文化・学芸員課程研究室、2. 理化学研究所、3. 国立極地研究所、4. 北海道大学

1. Humanities and Sciences/Museum Carriers, Musashino Art University, 2. RIKEN, 3. National Institute of Polar Research, 4. Hokkaido Univ.

A signal of the 27-day solar rotational period is often observed in cloud and lightning activities over the globe. Here we provide evidence of the 27-day periodicity of lightning activity in Japan using daily observational records of lightning for AD1989–2015. The 27-day period is detected only in a wide-area lightning activity over more than a few hundred thousand square kilometers. The 27-day signal is more prominent around the maxima of solar decadal cycles.

キーワード：太陽気候影響、雷活動、太陽自転周期

Keywords: Solar forcing of climate, Lightning activity, Solar rotational period

## 久住におけるシューマン共鳴の長期変動

### Long-term variation of Schumann resonance parameters at Kuju station

\*池田 昭大<sup>1</sup>、魚住 禎司<sup>2</sup>、吉川 顕正<sup>3</sup>、藤本 晶子<sup>2</sup>、阿部 修司<sup>2</sup>、野澤 宏大<sup>1</sup>

\*Akihiro Ikeda<sup>1</sup>, Teiji Uozumi<sup>2</sup>, Akimasa Yoshikawa<sup>3</sup>, Akiko Fujimoto<sup>2</sup>, Shuji Abe<sup>2</sup>, Hiromasa Nozawa<sup>1</sup>

1. 鹿児島工業高等専門学校、2. 九州大学国際宇宙天気科学・教育センター、3. 九州大学理学府

1. NIT, Kagoshima College, 2. ICSWSE, Kyushu Univ., 3. Department of Earth and Planetary Sciences, Kyushu Univ.

The Schumann resonance (SR) is the global resonance of electromagnetic waves generated by global lightning activity. The resonance is formed by the Earth-ionosphere cavity and the specific resonance frequency appears in ground magnetic field variation. Expectations for the SR have increased recently as an indicator of global and regional lightning activity. In order to use the SR parameters for studying such earth's climate, we need a better understanding of the long-term variations of the SR. In this study, we focused on the long-term variation of the SR at Kuju, Japan (KUJ; M.Lat. = 23.4 degree, M. Lon. = 201.0 degree).

The ground magnetic field variation in the extremely low frequency (ELF) range has been measured by an induction magnetometer at KUJ since 2003. The observation is a part of activities by International Center for Space Weather Science and Education Kyushu University.

The fundamental mode of the Schumann resonance (SR1) around 8 Hz can be seen at KUJ. The peak frequency of SR1 correlates with F10.7. Also the correlation between SR1 amplitude and F10.7 reveals. The SR parameters seem to be affected by the ionospheric conditions which change depending on solar activity.

キーワード：シューマン共鳴、太陽活動

Keywords: Schumann resonance, solar activity

# Development of an automated prediction method of CME arrival with SUANOO-CME

\*塩田 大幸<sup>1</sup>、草野 完也<sup>1</sup>

\*Daikou Shiota<sup>1</sup>, Kanya Kusano<sup>1</sup>

1. 名古屋大学 宇宙地球環境研究所

1. Institute for Space-Earth Environmental Research, Nagoya University

CMEs, especially their internal southward magnetic fields, are one of the main drivers of geomagnetic disturbances and hence the prediction of CME arrival to the Earth. The prediction of CME arrival to the Earth and the southward magnetic flux brought by the CMEs are one of crucial tasks for space weather forecast of magnetic storms.

Recently, we have developed a new MHD simulation capable of reproducing the interplanetary propagation of multiple CMEs with internal magnetic flux rope (Shiota & Kataoka 2016) called as SUANOO-CME. The simulation solves propagation of solar wind and CMEs in the inner heliosphere outer than the inner boundary at 30 Rs where the speed of all the bulk flow exceeds fast mode speed. The information of solar wind and CME is specified at the inner boundary with empirical and analytical models. The CME model has many free parameters such as the orientations of the internal magnetic flux rope, etc. Hence, in order to execute this simulation for use in real time forecast, we need a method to specify the free parameter that we cannot get any clue in the real time observations.

In this study, we present test results of MHD simulation (SUANOO-CME) for from May to September of 2005, applying a new method to specify the parameters of the CME model derived from only real time observations (SDO, GOES). We will evaluate the scores for forecast and discuss the current status of our capability for use in real time forecast.

キーワード：宇宙天気、コロナ質量放出、太陽風、MHD シミュレーション

Keywords: space weather, coronal mass ejection, solar wind, MHD simulation

# An investigation of numerical techniques in an MHD relaxation method for NLFFF extrapolation

\*三好 隆博<sup>1</sup>、草野 完也<sup>2</sup>、井上 諭<sup>2</sup>

\*Takahiro Miyoshi<sup>1</sup>, Kanya Kusano<sup>2</sup>, Satoshi Inoue<sup>2</sup>

1. 広島大学大学院理学研究科物理科学専攻、2. 名古屋大学宇宙地球環境研究所

1. Department of Physical Science, Graduate School of Science, Hiroshima University, 2. Institute for Space-Earth Environmental Research, Nagoya University

Prediction of explosive phenomena in the solar corona such as solar flares and coronal mass ejections (CMEs) is one of the most important components for space weather forecast. These phenomena are considered to be processes that abruptly liberate the magnetic energy stored in the solar corona due to photospheric motions. In order to clarify these processes, three-dimensional information of the magnetic field in the solar corona is needed. The data, however, cannot be directly obtained in the solar corona though that on the photosphere can be observed. Therefore, various reconstruction methods from the vector magnetic field on the photosphere to a three-dimensional magnetic field have been proposed so far [1]. Particularly, a nonlinear force-free field (NLFFF) is often suited to model low beta coronal plasmas and seems to reconstruct the solar coronal magnetic field.

A magnetohydrodynamic (MHD) relaxation method is one of the promising methods for the NLFFF extrapolation [1]. The NLFFF obtained by the MHD relaxation method is directly usable as an initial condition of nonlinear MHD simulations because available discretization methods of the governing equations, MHD relaxation equations, are similar to the MHD simulations. However, modern numerical techniques for MHD such as upwind-type [2] and higher-order schemes [3] have not been investigated well in the MHD relaxation method. Therefore, in this paper, we study the numerical techniques for the MHD relaxation equations. We will also discuss an application to an extended MHD relaxation method for a non-force-free field (NFFF).

[1] S. Inoue, Prog. Earth Planet Sci., 3:19, 2016.

[2] e.g., T. Miyoshi, K. Kusano, J. Comput. Phys., 208, 315, 2005.

[3] e.g., G.-S. Jiang, C.-W. Shu, J. Comput. Phys., 126, 202, 1996.

キーワード：太陽コロナ、非線形フォースフリー磁場、MHD緩和法

Keywords: Solar corona, NLFFF, MHD relaxation method



## Radiative cooling of Nitric Oxide emission observed by TIMED/SABER over asian sector during severe geomagnetic storms

\*Gaurav Bharti<sup>1</sup>, Sunil Krishna MV<sup>1</sup>

1. Indian Institute of Technology Roorkee

Nitric oxide (NO) is a very important trace species in the mesosphere-lower-thermosphere (MLT) region. It plays a significant role in the MLT region due to its low ionization energy and the presence of IR-active vibrational bands. The radiative emission due to NO at  $5.3 \mu\text{m}$  is the main cooling mechanism in the thermosphere. It is very well known that geomagnetic storms strongly alter the energetics and chemistry of thermosphere. The effect of geomagnetic storms on the nitric oxide (NO) radiative cooling at  $5.3 \mu\text{m}$ , in MLT region, has been studied over Asian sector during 26-29 September 2011 and 18-21 February 2014 using measurements made by SABER onboard the NASA's TIMED satellite. The SABER retrieved data along with the densities from NRLMSISE-00 model have been used to study the latitudinal variation of nitric oxide radiative cooling during the storm period. The variations induced in the radiative cooling are understood with the help of fluctuations in neutral species and the resulting changes in chemistry. It has been found that the NO radiative emission is strongly influenced by the storm conditions. The altitude of peak emission rate shows an upward movement in the higher latitude regions during day and night time. Similarly, the magnitude of peak emission rate is also observed to show enhancement in the high latitude sectors during the main phase of storm in day as well as night time. The atomic oxygen number density and O/N<sub>2</sub> ratio calculated using NRLMSISE-00 model, shows depletion in the higher latitude region. Consequently, there is a negative correlation among the nitric oxide volume emission rate, atomic oxygen density and O/N<sub>2</sub> ratio over Asian sector during the geomagnetic storm period. The various factors influencing the correlation between density ratios and radiative cooling will be presented.

Keywords: space weather, radiative cooling, thermosphere,

# Construction of global chorus wave intensity distribution from precipitating electron flux measurement and its application

\*栗田 怜<sup>1</sup>、掘 将大<sup>1</sup>、三好 由純<sup>1</sup>

\*Satoshi Kurita<sup>1</sup>, Masahiro Hori<sup>1</sup>, Yoshizumi Miyoshi<sup>1</sup>

1. 名古屋大学宇宙地球環境研究所

1. Institute for Space-Earth Environmental Research, Nagoya University

Whistler-mode chorus emissions are frequently observed in the inner magnetosphere during geomagnetically disturbed periods on the dawn and noon side magnetosphere. The waves have been considered as an important driver of the dynamics of inner magnetosphere. Resonant interactions of energetic electrons with chorus emissions result in flux enhancement of relativistic electrons in the outer radiation belt. On the other hand, the waves responsible for pitch angle scattering of electrons in the wide energy range from keV up to MeV range via pitch angle scattering. Previous studies show that chorus wave activities are correlated well with amount of energetic electron precipitations at low altitudes [Lam et al., 2010], and derive global intensity distributions of chorus waves during some events [Li et al., 2014; Chen et al., 2014]. It is possible that chorus wave intensity is constructed in global sense from the global map of precipitating electron fluxes. In this study we statistically investigate global distributions of > 30 keV precipitating electron fluxes observed by low-altitude POES satellites as a function of L, MLT, and Kp index. The distributions are compared with those of chorus wave intensity derived from the THEMIS satellites, and relationship between chorus wave intensity is parameterized as a function of L, MLT, and Kp index. Based on the parameterization, we successively construct global distributions of chorus wave intensity, and prediction of global chorus wave intensity variations is possible in cooperation with forecast of Kp index. In this presentation, we show the methodology of our analysis, obtained results, their performance to construct global distributions of chorus wave intensity, and its application to future research in the inner magnetosphere.

キーワード：放射線帯、ホイッスラーモード・コーラス、降下電子

Keywords: radiation belt, whistler mode chorus, precipitation

## 高緯度と低緯度コロナホール起源の太陽風変動と地球磁気圏応答 Solar wind variations originating from the high-latitude and low-latitude coronal holes and their response to the Earth's magnetosphere

\*中川 裕美<sup>1</sup>、新堀 淳樹<sup>2</sup>、野澤 恵<sup>1</sup>

\*Yumi Nakagawa<sup>1</sup>, Atsuki Shinbori<sup>2</sup>, Satoshi Nozawa<sup>1</sup>

1. 茨城大学大学院理工学研究科宇宙地球システム科学専攻、2. 名古屋大学宇宙地球環境研究所

1. Ibaraki University, 2. Institute for Space-Earth Environment Research (ISEE), Nagoya University

Coronal holes are frequently observed in the declining phase of the solar cycle and have open magnetic fields expanding to the interplanetary space. The solar atmospheric plasma flows out along the magnetic field line from the coronal hole, and causes a major disturbance of the Earth's magnetosphere such as geomagnetic storm and substorm. Tsurutani et al.(2006) proposed that analyses of the temporal area of polar coronal holes over the solar cycle [Harvey et al., 2000; Harvey and Recely, 2002] provide a good idea of the geoeffectiveness of high speed streams over the solar cycle, but the effects of isolated equatorial coronal holes has yet to be evaluated. In order to clarify the effects of the Earth's magnetosphere and ionosphere associated with solar wind disturbances originating from the equatorial coronal hole, we conducted the superposed epoch analysis of the variations of coronal hole area, solar wind, interplanetary magnetic field, and geomagnetic indices (AE and SYM), and investigated a difference of the solar wind variations from between the high-latitude and low-latitude coronal holes and their response to the Earth's magnetosphere. In the present analysis, we used the Sun whole two-dimension images taken by the solar whole the extreme ultraviolet imaging telescope (EIT) onboard the Solar and Heliospheric Observatory (SOHO). Solar wind data are obtained from the advanced composition explorer (ACE), Wind, and OMNI2 data provided by NASA CDAWeb. The data period is October 1996 –May 2013. For the coronal holes area, we defined the threshold of the solar brightness in the EUV range as a half of the median value of the intensity in a whole area and divided the solar surface in four regions: (-60 - -30, -30 - 30), (-30 - 0, -30 - 30), (0 - 30, -30 - 30), and (30 - 60, -30 - 30) (degrees) in the solar latitude and longitude, respectively. Moreover, we determined the coronal holes area as a ratio of pixel numbers less than the threshold to each region. As a result, we found 5 low-latitude coronal holes. For the response to the Earth's magnetosphere, we used the SYM-H and AE indices provided by World Data Center (WDC) for Geomagnetism, Kyoto University. The superposed epoch analysis results showed that when the coronal hole area become maximum, the solar wind density increased from 3 to 13 /cc, rapidly. At this time, the solar wind speed was minimum, and the Bx, By, and Bz were directed sunward, dawnward, and northward, respectively. After that, the solar wind density decreased to 3 /cc, and the solar wind speed increase from 350 km/sec to 600 km/ sec within 3 days. At the same time when the solar wind speed increased rapidly, the Bx, By, and Bz were directed earthward, duskward, and southward, respectively. On the other hand, the AE and SYM indices showed a significant increase and decrease within 3 days after the solar wind speed increased and the IMF Bz component became negative. However, the magnitude of the AE and SYM variations was small in a case of the solar wind originating from the high-latitude coronal hole. Therefore, it can be concluded that the effects of the Earth's magnetosphere are larger for the equatorial coronal hole than the high-latitude one.

キーワード：共回転相互作用領域、コロナホール、太陽風変動

Keywords: Co-rotation Interaction Region, Coronal holes, Solar wind variations

## High Intensity Long Duration Continuous Auroral-Electrojet Activity(HILDCAA) in relation to Geomagnetically Induced Current(GIC)

\*Krishna Khanal<sup>1,2</sup>, Binod Adhikari<sup>1,2</sup>, Narayan Pd Chapagain<sup>1</sup>, Binod Bhattarai<sup>1,2</sup>

1. Tribhuwan University, Nepal, 2. St. Xavier's College, Nepal

GIC in the buried pipelines during intense geomagnetic storms has long been a subject of research because of its corroding potentiality to the pipeline material accompanied by the production of telluric current. In this work, for the first time, we have analyzed GIC statistics recorded in Finnish Natural Gas pipeline during three HILDCAA events with different interplanetary sources: CIR storm preceded, ICME storm preceded and Isolated. Despite the weak nature of HILDCAAs in terms of Dst reduction, their cumulative contribution in corrosion process is conjectured to be significant as they occur continuously several days to weeks. Analyzing the result, it is found that continuous fluctuation in GIC is observed throughout the AE intensification during all three events, which corresponds to the alfvanic wave fluctuation during HILDCAAs. However, there is no peak to peak relationship between AE index and GIC. Equivalently, every AE intensification does not necessarily account for the higher GIC amplitudes. We have adopted wavelet analysis and cross correlation methods to study the nature of GIC and its possible correlation with other parameters during HILDCAAs. From discrete wavelet transform we analyzed the singularities associated with the discontinuities present in GIC signal up to the three decomposition levels. Similarly, the result of discrete wavelet transform showed spectral variabilities in GIC with different periodicities.

Keywords: HILDCAA, GIC, Wavelet Analysis, Cross Correlation

## 磁気擾乱時における中低緯度領域電磁誘導応答の研究 Research for electromagnetic induction response in the low-and-mid-latitudinal region at geomagnetic disturbances

\*中原 美音<sup>1</sup>、松下 拓輝<sup>1</sup>、吉川 顕正<sup>2</sup>、魚住 禎司<sup>3</sup>、藤本 晶子<sup>3</sup>、阿部 修司<sup>3</sup>

\*Mio Nakahara<sup>1</sup>, Hiroki Matsushita<sup>1</sup>, Akimasa Yoshikawa<sup>2</sup>, Teiji Uozumi<sup>3</sup>, Akiko Fujimoto<sup>3</sup>, Shuji Abe<sup>3</sup>

1. 九州大学大学院理学府地球惑星科学専攻宇宙地球電磁気学分野、2. 九州大学大学院理学研究院地球惑星科学部門、3. 国際宇宙天気科学・教育センター

1. Spade and Earth Electromagnetism Laboratory, Department of Earth and Planetary Sciences, Graduate School of Science, Kyushu university, 2. Department of Earth and Planetary Sciences, Kyushu University, 3. International Center for Space Weather Science and Education Kyushu University

地上の磁場は様々な宇宙天気現象と連動して、日々変動している。そのような地磁気擾乱現象が引き起こす現象の一つとしてGIC(Geomagnetically Induced Current)が知られている。GICの推定には地表の電場の水平成分が有用[Pulkkinen et al., 2007]と言われており、地表に誘導される電場については現実的な磁場変動を用いることによる推定が議論されている。GICによる被害報告は高緯度領域に多いが、地磁気的な緯度が低い日本でもGICの検討が開始されている[Watari et al., 2015]。本研究では中低緯度領域に位置する日本で、磁気擾乱現象を定量化した指数からGICの観測、予測に到るまでの過程理解を目的とし、磁気擾乱指数と磁場の関連性についての調査を行った。本研究ではGICの主要な起電力となりうる地表の誘導電場の水平成分の推定に着目する。磁場鉛直(Z)成分の変動成分、Faradayの電磁誘導の法則に基づき、誘導電場の水平成分分布を求めることが出来るであろうという観点から、磁場Z成分時間微分について考察し、オーロラ嵐の大きさと磁場Z成分の相関についての解析を行った。また、磁場変動から地表の誘導電場を見積もるための初段階として、磁場と電場の変動の相関関係についての考察についても紹介する予定である。

キーワード：電磁気、宇宙天気、GIC

Keywords: Electromagnetism, Space weather, GIC

# Simulation of Geomagnetically Induced Electric Field Originating from Field-Aligned Current

\*栗栖 一樹<sup>1</sup>、海老原 祐輔<sup>1</sup>、中村 紗都子<sup>1</sup>

\*Kazuki Kurisu<sup>1</sup>, Yusuke Ebihara<sup>1</sup>, Satoko Nakamura<sup>1</sup>

1. 京都大学生存圏研究所

1. Reserach Institute for Sustainable Humanosphere, Kyoto University

Solar wind has a large impact on the Earth's magnetosphere. When the solar wind with southward Interplanetary Magnetic Fields (IMF) comes, magnetic storms and substorms occur in the near-Earth space. Due to this change in the space environment, field-aligned current (FAC) generated in the magnetosphere flows into the ionosphere. In addition to FAC, the magnetspheric and ionospheric currents induce geomagnetically induced electric field (GIE) on the surface of the Earth. GIE generates geomagnetically induced current (GIC) flowing in the conducting electrical wires. GIC causes severe damages on our life. In 1989, for example, large amplitude GIC flowed in the province of Quebec, Canada, and interrupted electric services over nine hours. The purpose of our study is to understand the relationship between the ionospheric current and GIE. Here, We focused on GIE in the polar region, and conducted finite-difference time-domain (FDTD) simulations. We used the 3-dimensional model that contains the ionosphere, the atmosphere, land and/or sea. We set FAC and Pedersen current as current sources. In order to evaluate the validity of our simulation, we used the Häkkinen method. This method calculates the electromagnetic field induced on the surface in frequency domain. We compared the electromagnetic field calculated by FDTD simulation with the one by the Häkkinen method. We will investigate the simulation results in future for a better understanding and modeling of GIC.

キーワード：地磁気誘導電流、地磁気誘導電界、沿磁力線電流

Keywords: geomagnetically induced current, geomagnetically induced electric field, field-aligned current

## Multi-instrument observations of periodic poleward moving polar cap arcs

\*Zan-Yang Xing<sup>1</sup>, Qing-He Zhang<sup>1</sup>, Desheng Han<sup>2</sup>, Yongliang Zhang<sup>3</sup>, Jun Liang<sup>4</sup>, Joran Moen<sup>5</sup>, Zejun Hu<sup>2</sup>, P. Jayachandran<sup>6</sup>, Natsuo Sato<sup>7</sup>, Shunrong Zhang<sup>8</sup>, Keisuke Hosokawa<sup>9</sup>, Y. Wang<sup>1</sup>, Yu-Zhang Ma<sup>1</sup>

1. Shandong University, 2. Research Institute of China, 3. Johns Hopkins University, 4. University of Calgary, 5. University of Oslo, 6. University of New Brunswick, 7. National Institute of Polar Research, 8. Massachusetts Institute of Technology, 9. University of Electro-Communications

Polar cap arcs (PCAs) are one of the outstanding phenomena in the polar cap region. We examined a case of periodic poleward moving arcs observed on January 3, 2014 by dual all-sky imagers, one at the Chinese Yellow River Station (YRS) and the other at Resolute Bay (RSB), the spaced-based SSUSI imager onboard DMSP spacecraft and Resolute Bay Incoherent Scatter Radar (RISR) during quiet geomagnetic conditions. We found that some poleward moving arcs observed at RSB were repeatedly detached from the dawnside auroral oval, which is consistent with the IMF-By polarity, and some arcs observed at YRS were likely less poleward moving from the duskside auroral oval. We also observed some periodic spatial arcs by DMSP SSUSI imagers and strong plasma velocity shears around these arcs. At the same time, the precipitating particles observed in the ionosphere associated with these PCAs showed magnetosheath-like properties. Moreover, the RISR data show that the F-region plasmas above these arcs were structured, with significant E-region ionization above an intense arc. Based on these observational results, we suggest that these periodic poleward moving PCAs may be triggered by bursts of dayside magnetic reconnection.

Keywords: periodic poleward moving arcs, plasma velocity shears, magnetic reconnection



## Comparison of magnetospheric magnetic field variations at quasi-zenith orbit based on Michibiki observation and REPPU global MHD simulation

\*久保田 康文<sup>1</sup>、長妻 努<sup>1</sup>、田 光江<sup>1</sup>、中溝 葵<sup>1</sup>、松本 晴久<sup>2</sup>、田中 高史<sup>3</sup>

\*Yasubumi Kubota<sup>1</sup>, Tsutomu Nagatsuma<sup>1</sup>, Mitsue Den<sup>1</sup>, Aoi Nakamizo<sup>1</sup>, Haruhisa Matsumoto<sup>2</sup>, Takashi Tanaka<sup>3</sup>

1. 情報通信研究機構、2. 宇宙航空研究開発機構、3. 九州大学

1. National Institute of Information and Communications Technology, 2. Japan Aerospace Exploration Agency, 3. Kyushu University

We are developing a numerical simulator for future space weather forecast using magnetosphere-ionosphere coupling global MHD simulation called REPPU (REProduce Plasma Universe) code. We investigate the validity of the MHD simulation result as compared with observation. In this study we simulate some events including both quiet and disturbed geomagnetic conditions using OMNIWeb solar wind data. The simulation results are compared with magnetic field observations from Michibiki satellite, which is on the quasi-zenith orbit (QZO). In quiet geomagnetic condition, magnetic field variations at QZO obtained from simulation results have good consistency as compared correspondence with those from Michibiki observation. In disturbed geomagnetic condition in which the Dst < -50 nT, however, V component of magnetic field variations from simulation results tend to deviate from observations especially at the night side. We consider that this deviation during disturbed geomagnetic condition might be due to tail and/or ring current enhancement which is already suggested by many other MHD simulation studies as compared with the magnetic field observation at geosynchronous orbit. In this presentation, we will discuss the cause of this discrepancy in more detail with studying the relationship between the magnetic field deviation and some parameters such as Dst and solar wind.

## Reconstructing polar asymmetry of GLE69 via WASAVIES simulation

\*PARK INCHUN<sup>1</sup>、三好 由純<sup>1</sup>、佐藤 達彦<sup>2</sup>、片岡 龍峰<sup>3</sup>、久保 勇樹<sup>4</sup>、三宅 晶子<sup>5</sup>

\*PARK INCHUN<sup>1</sup>, Yoshizumi Miyoshi<sup>1</sup>, Tatsuhiko Sato<sup>2</sup>, Ryuho Kataoka<sup>3</sup>, Yuki Kubo<sup>4</sup>, Shoko Miyake<sup>5</sup>

1. 名古屋大学宇宙地球環境科学研究所、2. 日本原子力研究開発機構、3. 国立極地研究所、4. 情報通信研究機構、5. 茨城工業高等専門学校

1. Nagoya University Institute for Space-Earth Environmental Research, 2. Japan Atomic Energy Agency, 3. National Institute of Polar Research, 4. National Institute of Information and Communications Technology, 5. National Institute of Technology, Ibaraki College

Solar Energetic Particle(SEP) is one of the most important components for predicting space weather.

When a powerful SEP event occurs, neutron observations on the ground can detect the increase of flux.

This event is called Ground Level Enhancement(GLE).

On January 20, 2005, a powerful GLE (GLE69) occurred, causing significant flux enhancement on south pole region. The McMurdo neutron monitor in Antarctica recorded 18 times more fluxes compared to Thule station in Greenland. Previous studies suggested that this asymmetry phenomenon is related to IMF direction, which has never been confirmed by the simulation.

In this research, we investigate this asymmetry of neutron flux using WASAVIES (Warning System for Aviation Exposure to SEP [Kataoka et al., 2014]) which consists of SEP simulation in the heliosphere, particle trajectory tracing in the magnetosphere, and the air shower simulation. Using WASAVIES, we investigate propagation direction of SEPs and compare the results with neutron monitor observations. Initial results of this study indicate that SEPs come mainly from southern-hemisphere, which is consistent with previous studies. The results also suggest importance of transportation of SEP in the azimuthal direction, which has not included in this study.

キーワード : WASAVIES、太陽高エネルギー粒子、宇宙天気

Keywords: WASAVIES, Solar Energetic Particle, Space Weather

# 国際宇宙ステーションにおける高エネルギー電子・陽子の長期的観測

## The long term observation of the high energy electrons and protons on the Internatilnal Space Station

\*上野 遥<sup>1</sup>、松本 晴久<sup>1</sup>、片岡 龍峰<sup>2</sup>

\*Haruka Ueno<sup>1</sup>, Haruhisa Matsumoto<sup>1</sup>, Ryuho Kataoka<sup>2</sup>

1. 宇宙航空研究開発機構、2. 国立極地研究所

1. Japan Aerospace Exploration Agency, 2. National Institute of Polar Research

JAXAでは2009年より高エネルギー粒子モニタ (Standard Dose Monitor: SDOM) を運用し、部品材料の劣化や電子部品の誤動作等の原因となる電子、陽子、 $\alpha$ 線等の高エネルギー軽粒子の粒子別エネルギー分布を計測している。また近年ではISS船外での線量評価のために、NASAへのデータ提供も実施しており、被ばく線量評価のデータとして活用されている。

本講演ではSDOMデータの太陽周期変化と地磁気活動依存性を示し、また、Kataoka et al. (2016) で報告されたように、太陽活動に関連した電子の降込みがISS軌道で観測されており、その時のSDOMの観測データにもとづいて、放射線帯の変動や船外活動における被ばく量の議論をする。

キーワード：国際宇宙ステーション、宇宙放射線、放射線被ばく、高エネルギー粒子

Keywords: the International Space Station, Space radiation, Radiation exposure, High energy particles

## Cross-calibration of high-energy electron observations at geostationary earth orbit

\*長妻 努<sup>1</sup>、坂口 歌織<sup>1</sup>、越石 英樹<sup>2</sup>

\*Tsutomu Nagatsuma<sup>1</sup>, Kaori Sakaguchi<sup>1</sup>, Hideki Koshiishi<sup>2</sup>

1. 国立研究開発法人 情報通信研究機構、2. 国立研究開発法人 宇宙航空研究開発機構

1. National Institute of Information and Communications Technology, 2. Japan Aerospace Exploration Agency

Cross-calibration of high-energy electron observations obtained from individual satellites are quite essential procedure for reconstruction of high-energy electron distribution at geostationary earth orbit (GEO). To compare high-energy electron data from individual satellite, we should find the period where the L\*-value (drift shell) is the same. Because the magnetic dipole axis is not aligned with the rotational axis of the Earth, L\*-value of each GEO satellite is changing depending on the longitude of GEO and magnetic local time. In addition, L\*-value of each GEO satellite also changes depending on geospace disturbances. We have compared Himawari-8 high-energy electron (SEDA-e) data with GOES 15 high-energy electron (MAGED and EPEAD) data during the same L\*-value of both satellites with quiet geomagnetic condition. This result shows quite good correspondence of both observation, and factor of sensitivity can be estimated.

Our previous study showed seasonal dependence of sensitivity difference between high energy electron sensor onboard Himawari and that onboard Kodama. We found that this seasonal dependence might be caused by the north-south drift of the Kodama's orbit. Previous study, we assumed that the orbit of Kodama is aligned with GEO. This result suggests that the detailed orbit information is important for cross-calibration. Based on this cross-calibration procedure, we can produce particle distribution at GEO. In our presentation, some sample events of particle distribution at GEO will be shown, and the future plan for producing particle distribution in the inner magnetosphere using energetic electron data from ARASE(ERG), Van-Allen Probes, and GPS will be introduced.

キーワード：宇宙天気予報、磁気圏粒子分布、プラズマ粒子データ校正

Keywords: space weather forecast, magnetospheric particle distribution, cross-calibration of particle data

## On two-dimensional distribution of trapped protons from solar cell degradation of the Akebono satellite

\*三宅 亙<sup>1</sup>、戸田 穂乃香<sup>1</sup>、三好 由純<sup>2</sup>、松岡 彩子<sup>3</sup>

\*Wataru Miyake<sup>1</sup>, Honoka Toda<sup>1</sup>, Yoshizumi Miyoshi<sup>2</sup>, Ayako Matsuoka<sup>3</sup>

1. 東海大学工学部、2. 名古屋大学宇宙地球環境研究所、3. JAXA宇宙科学研究所

1. Tokai University, 2. ISEE, Nagoya University, 3. ISAS/JAXA

We have been studying on L-shell distribution of energetic (>10 MeV) protons from solar cell degradation of Akebono satellite orbiting in the inner magnetosphere. We obtained more compact distribution of the trapped protons than given by the AP8 and AP9 models. In our previous study (Miyake et al., 2014; 2015), we assumed that proton flux varies along the field line in the same rate as in the AP8 model. If the flux is more confined around the equator, the L-shell distribution may be possibly widen and the difference from the previous models can be smaller. Recent observation of the Van Allen probe has showed that quite anisotropic distribution of pitch angle which leads to spatial distribution confined near the equator (Selesnick et al., 2014). Thus, we introduce some different variation along the field line and seek the best-fit to the observed degradation of solar cells.

キーワード：あけぼの衛星、プロトン放射線帯、太陽電池劣化

Keywords: Akebono satellite, proton radiation belt, solar cell degradation

## Development of information system of spacecraft surface charging potential and discharging alert

\*中村 雅夫<sup>1</sup>、川内 諒太<sup>1</sup>、寺岡 毅<sup>1</sup>

\*Masao Nakamura<sup>1</sup>, Ryota Kawachi<sup>1</sup>, Tsuyoshi Teraoka<sup>1</sup>

1. 大阪府立大学

1. Osaka Prefecture University

Spacecrafts can be damaged by surface charging and resultant discharging arcs. We are developing information system of surface charging potential and discharging alert for individual spacecraft. We create lookup tables of surface potential for many combinations of plasma temperatures and densities using charging simulation for the spacecraft engineering model with a spacecraft charging analysis software and/or spacecraft observational results. Using interpolation with the lookup tables, we can estimate surface potential on demand for any given on-orbit plasma environment. The discharging alert also can be accomplished by additional information about the estimated differential potential and the experimental discharge potential of the spacecraft. We will introduce current status of our system.

キーワード：宇宙機表面帯電、衛星放電警報、宇宙天気

Keywords: Spacecraft surface charging, Spacecraft discharging alert, Space weather