

## Software-type Wave-Particle Interaction Analyzer on board the ARASE satellite

\*Hirotugu Kojima<sup>1</sup>, Yuto Katoh<sup>2</sup>, Mitsuru Hikishima<sup>3</sup>, Takeshi Takashima<sup>3</sup>, Kazushi Asamura<sup>3</sup>, Yoshizumi Miyoshi<sup>4</sup>, Yoshiya Kasahara<sup>5</sup>, Satoshi Kasahara<sup>6</sup>, Takefumi Mitani<sup>3</sup>, Nana Higashio<sup>7</sup>, Ayako Matsuoka<sup>3</sup>, Iku Shinohara<sup>3</sup>

1. Research institute for sustainable humanosphere, Kyoto University, 2. Graduate school of science, Tohoku university, 3. Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 4. Institute for Space-Earth Environmental Research, Nagoya University, 5. Information Media Center, Kanazawa University, 6. Graduate school of science, University of Tokyo, 7. Research and Development Directorate, Japan Aerospace Exploration Agency

The Wave-particle interaction analyzer (WPIA) is a new method of observing wave-particle interactions in space. Based on the cutting-edge of technologies in the onboard instruments, the Software-type WPIA (S-WPIA) is installed in the ARASE satellite, which was successfully launched on December 20, 2016. The present paper introduces the principles of the WPIA and describes the detailed design of the S-WPIA on board the ARASE satellite. Understanding wave-particle interactions is essential in the study on space plasma environments, because space plasmas are collisionless and their kinetic energies are transferred through wave-particle interactions. In the conventional way of the study on wave-particle interactions via satellites, we have compared features of plasma waves with velocity distribution functions. However, that conventional way is not appropriate for identifying wave-particle interactions quantitatively. The nature of wave-particle interactions lies in the phase difference between electric field vectors ( $\mathbf{E}$ ) and velocity vectors of particles ( $\mathbf{V}$ ). This appears as the inner product form as  $\mathbf{E} \cdot \mathbf{V}$ . The conventional way using velocity distribution functions misses the information of this phase difference, because the velocity distribution function is obtained on the time integration basis. The WPIA overcomes the above problem in the conventional method by handling each detected particle and instant electric field intensity with keeping the enough accuracy in the relative time difference between them. The ARASE satellite and its onboard S-WPIA instrument should be frontiers in the study of wave-particle interactions. The leading edge of the system in the ARASE satellite allows us to collect whole information of particles at every particle detection timing and instant electric fields at every sampling timing. The collected particle and waveform data are stored on the onboard data storage called Mission Data Recorder (MDR). The S-WPIA calculates the phase difference and other quantities onboard reading out the data from the MDR and send the results as well as raw data of particles and plasma waves to the ground.

The main objective of the S-WPIA on board the ARASE satellite is to detect quantitatively the wave-particle interaction related to the generation of the Chorus emissions. The SWPIA also targets the quantitative detection of accelerations of electrons due to plasma waves. The function of the S-WPIA has been already confirmed during the initial operation of the ARASE satellite. The present paper introduces the details of the S-WPIA and discuss the strategy how we meet the objective of the S-WPIA in the operation.

## Onboard Processing on PWE OFA/WFC (Onboard Frequency Analyzer/Waveform Capture) aboard the ERG(ARASE) Satellite

\*Shoya Matsuda<sup>1</sup>, Yoshiya Kasahara<sup>2</sup>, Hirotsugu Kojima<sup>3</sup>, Yasumasa Kasaba<sup>4</sup>, Satoshi Yagitani<sup>2</sup>, Mitsunori Ozaki<sup>2</sup>, Tomohiko Imachi<sup>2</sup>, Keigo Ishisaka<sup>5</sup>, Mamoru Ota<sup>2</sup>, Atsushi Kumamoto<sup>4</sup>, Fuminori Tsuchiya<sup>4</sup>, Yoshizumi Miyoshi<sup>1</sup>, Ayako Matsuoka<sup>6</sup>, Iku Shinohara<sup>6</sup>

1. Nagoya University, 2. Kanazawa University, 3. Kyoto University, 4. Tohoku University, 5. Toyama Prefectural University, 6. ISAS/JAXA

Exploration of energization and Radiation in Geospace (ERG) is a mission for understanding particle acceleration, loss mechanisms, and the dynamic evolution of space storms in the context of cross-energy and cross-regional coupling [Miyoshi et al., 2012]. The ERG (ARASE) satellite was launched on December 20, 2016, and successfully inserted into an orbit.

The Plasma Wave Experiment (PWE) is one of the science instruments on board the ERG satellite to measure electric field and magnetic field in the inner magnetosphere. PWE consists of three sub-components, EFD (Electric Field Detector), OFA/WFC (Onboard Frequency Analyzer and Waveform Capture), and HFA (High Frequency Analyzer). Especially, OFA/WFC measures electric and magnetic field spectrum and waveform from a few Hz to 20 kHz. OFA/WFC processes signals detected by a couple of dipole wire-probe antenna (WPT) and tri-axis magnetic search coils (MSC) installed onboard the satellite. The PWE-OFA subsystem calculates and produces three kind of data; OFA-SPEC (power spectrum), OFA-MATRIX (spectrum matrix), and OFA-COMPLEX (complex spectrum). They are continuously processed 24 hours per day and all data are sent to the ground. OFA-MATRIX and OFA-COMPLEX are used for polarization analyses and direction finding of the plasma waves. The PWE-WFC subsystem measures raw (64 kHz sampled) and down-sampled (1 kHz sampled) burst waveform detected by the WPT and the MSC sensors. It activates by a command, automatic triggering, and scheduling.

The initial check-out process of the PWE successfully completed, and initial data has been obtained. In this presentation, we introduce onboard processing technique on PWE OFA/WFC and its initial results.

Keywords: ERG/ARASE, Plasma wave, Chorus wave, EMIC wave

## Charged particle measurements in the radiation belts by ERG

Kazushi Asamura<sup>1</sup>, Nana Higashio<sup>3</sup>, Masafumi Hirahara<sup>4</sup>, \*Satoshi Kasahara<sup>2</sup>, Yoichi Kazama<sup>7</sup>, haruhisa matsumoto<sup>3</sup>, Takefumi Mitani<sup>1</sup>, Wataru Miyake<sup>5</sup>, Yushi Suto<sup>2</sup>, Takeshi Takashima<sup>1</sup>, Bo-Jhou Wang<sup>7</sup>, Shiang-Yu Wang<sup>7</sup>, Kazuhiro Yamamoto<sup>6</sup>, Shoichiro Yokota<sup>1</sup>

1. ISAS, 2. The university of Tokyo, 3. JAXA, 4. Nagoya University, 5. Tokai University, 6. Kyoto University, 7. ASIAA

Radiation belts show mysterious and dynamic variability during geospace storms. The ERG spacecraft aims to observe the cross-energy coupling plasma physics behind the decay and enhancement of the radiation belts. In order to cover the broad energy range from 10 eV up to 20 MeV, ERG is equipped with 6 particle instruments (XEP, HEP, MEP-e, MEP-i, LEP-e, and LEP-i). Here we review the specifications of these sensors.

## Observations of low-energy ions with Arase/LEPi

\*Kazushi Asamura<sup>1</sup>, Yushi Suto<sup>1,3</sup>, Yoichi Kazama<sup>2</sup>, Shoichiro Yokota<sup>1</sup>, Satoshi Kasahara<sup>3</sup>, Tomoaki Hori<sup>3</sup>, Yoshizumi Miyoshi<sup>4</sup>

1. Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 2. Academia Sinica Institute of Astronomy and Astrophysics, 3. University of Tokyo, 4. Nagoya university

Arase satellite has been successfully launched into the orbit. The Arase project is focuses on understandings of plasma acceleration / loss / transport mechanisms taking place in the inner magnetosphere. Since energy of particles is ranges over several orders in geospace, Arase carries several particle instruments in order to cover wide energy range, from 10eV/q to 200keV/q (ions), and from 10eV to 10MeV (electrons). LEPi (Low-energy particle experiments - ion mass analyzer) is one of the instruments onboard Arase, which is an energy-mass spectrometer designed to measure the ions with energies from ~0.01keV/q up to 25keV/q. When the instrument passes the initial checkout phase after launch, it will start regular observations.

We will present the current status of LEPi.

Keywords: Arase, low-energy ion, observation

## Ground calibration experiments of Magnetic field experiment on the ERG satellite

\*Mariko Teramoto<sup>1</sup>, Ayako Matsuoka<sup>2</sup>, Reiko Nomura<sup>2</sup>

1. Institute for Space-Earth Environmental Research, Nagoya University, 2. Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency

The Arase (ERG) satellite was launched on 20 December 2016 to study plasma process in the inner magnetosphere. The Magnetic field experiments (MGF), which is one of the scientific experiments onboard the Arase satellite, observes the background magnetic field and its low frequency fluctuations. The MGF has a set of tri-axis ring-core type fluxgate sensors (MGF-S) to observe the magnetic field in the inner magnetosphere. For accurate measurements of the magnetic field vector along the Arase orbits, ground calibration experiments of MGF-S are needed.

We have been performed in order to determine the sensitivity and alignment via ground calibration experiments. From response of MGF-S to known applied magnetic field, we determined the sensitivity of each axis and found that the error of the sensitivity is less than 0.06%. The axis of the sensor is orthogonal to each other within 0.95 degrees. The estimated error of alignment is within 0.07 degrees. We also have examined the temperature dependence of the sensitivity and offset. The sensitivities relative to the room temperature have linearity with the standard error less than 0.0016, while the offset of the sensors have no clear linearity but reproducibility against temperature. From these ground examinations, the determination accuracies of the amplitude and direction of the magnetic field observed by the MGF will satisfy the science requirements for the Arase observations. In this presentation, we will also evaluate and show measurement error of MGF-S along the Arase orbit.

Keywords: ERG satellite, Magnetic Filed experiment

## Modeling energetic particles generated by substorm dipolarizations.

\*Konstantin Kabin<sup>1</sup>, German Kalugin<sup>1</sup>, Eric Donovan<sup>2</sup>, Emma Spanswick<sup>2</sup>

1. Royal Military College of Canada, 2. The University of Calgary

We describe a novel model of magnetotail which is easily controlled by several adjustable parameters, such as the thickness of the tail and the location of transition from dipole-like to tail-like magnetic field lines. The model is fully three-dimensional and includes the day-night asymmetry of the terrestrial magnetosphere, although the field lines are contained in the meridional planes. This model is well suited to studies of the magnetotail dipolarizations which we consider to be associated with the movements of the transition between dipole-like and tail-like field lines. Induced electric fields generated by this reconfiguration of the magnetotail are capable of energizing electrons and ions. In some cases, the energy of the particles can increase by a factor of 25 or more. These electric fields are also responsible for transport of the energized particles closer to the Earth where they can be observed, either in-situ by the satellites, or indirectly by ground-based instruments, such as riometers. Results of our calculations suggest that this scenario provides a plausible explanation of substorm particles injections.

Keywords: energetic particles in the inner magnetosphere, magnetotail modeling, substorm dipolarizations, substorm particle injections

## Relativistic effect on energetic particle injections during the 14 July 2013 Substorm

\*Tzu-Fang Chang<sup>1,2</sup>, Chih-Yu Chiang<sup>1</sup>, Chio-Zong Cheng<sup>1</sup>, Sunny Wing-Yee Tam<sup>1</sup>

1. Institute of Space and Plasma Sciences, National Cheng Kung University, Taiwan, 2. Institute of Astronomy and Astrophysics, Academia Sinica, Taiwan

Substorm energetic particle injections observed by multiple satellites during a small storm on 14 July 2013 are studied by considering the relativistic effect. We combine the ground observations and in situ magnetic field and particle data from geosynchronous satellites and Van Allen Probes in the inner magnetosphere to investigate the current systems and the energetic particle injections associated with the substorm. Based on a classical electromagnetic field pulse model, we propose a relativistic model to simulate the evolution of energetic particle injections during the particular substorm event. Detailed discussion of the differences in relativistic calculation from non-relativistic ones and satellite observations will be presented.

Keywords: energetic particle injection, inner magnetosphere, substorm

## Observations of Electric Fields at Dipolarization Sites from THEMIS mission: Preliminary Results

\*Kaiti Wang<sup>1</sup>, Ching-Huei Lin<sup>2</sup>, Po-Yen Chen<sup>1</sup>, Shi-Hung Chuang Yang<sup>1</sup>, Tai-Yuan Wang<sup>1</sup>, Yi-Ju Chen<sup>1</sup>, Guo-Cing Su<sup>1</sup>, Zhuo-Ren Lang<sup>3</sup>

1. Tamkang University, 2. Chien-Hsin University of Science and Technology, 3. National Cheng Kung University

The electric fields at dipolarization sites in the Earth's tailside have been found to be disturbed when geomagnetic activities occurs (i.e., AL index decreases). These fields can accelerate electrons so that they are possible to be associated with electron injection or changes in electrons' pitch-angles. Therefore, it is essential to understand the variations of these environmental electric fields when dipolarization occurs. In this study, observational data of electric fields from the EFI (Electric Field Instrument) on board of THEMIS mission are analyzed. The database are selected based on dipolarization events around 10 Earth radii identified according to THEMIS observations from year of 2008 to 2011. Both the large-scale and wave-scale features of these dipolarization electric fields will be investigated. The preliminary results of this analysis will be shown in the poster.

Keywords: THEMIS mission, Electric Fields, Dipolarization

## Response of relativistic electron microbursts to the arrival of high speed solar wind streams and its relation to flux variation of radiation belt electrons

\*Satoshi Kurita<sup>1</sup>, Yoshizumi Miyoshi<sup>1</sup>, J. B. Blake<sup>2</sup>, R. H. W. Friedel<sup>3</sup>

1. Institute for Space-Earth Environmental Research, Nagoya University, 2. The Aerospace Corporation, CA, USA, 3. Los Alamos National Laboratory, NM, USA

Relativistic electron microbursts are short-lived bursty precipitations of relativistic electrons observed by low-altitude satellite in the radiation belt. They are considered as a consequence of pitch angle scattering of radiation belt electrons by discrete whistler-mode emissions known as chorus. Microbursts are frequently observed during geomagnetic storms and previous studies show that atmospheric loss through microbursts appears to contain enough electrons to deplete the radiation belt. They suggest that microburst is an important loss process of radiation belt electrons during the main phase of geomagnetic storms. Microbursts are also frequently observed during high-speed solar wind stream (HSS) events, while important solar wind parameters for the frequent microburst precipitations have not been well understood. We perform a superposed epoch analysis of the microburst occurrence during HSS events, considering the polarity of interplanetary magnetic field (IMF) and solar wind speed according to the method used by Miyoshi and Kataoka (2008). We find the most frequent microburst precipitations during the highest-speed solar wind streams with a southward offset of IMF (SBZ-fast HSS events), indicating that both the southward IMF and fast solar wind are important for enhanced microburst precipitations. We also demonstrate that fluxes of radiation belt electrons with energies from hundreds keV up to 7 MeV preferentially increase during the SBZ-fast HSS events. The result suggests that loss through microbursts is not major loss process of radiation belt during the HSS events. We conclude that relativistic electron microbursts can be a proxy of acceleration of MeV electrons by chorus.

Keywords: radiation belt, high speed solar wind streams, radiation belt dropout

## Transport and acceleration of electrons trapped in the inner magnetosphere in response to interplanetary shock

\*Takuya Ikeda<sup>1</sup>, Yusuke Ebihara<sup>1</sup>

1. Kyoto University

Interplanetary (IP) shock is known to have a large effect on the electrons trapped in the inner magnetosphere. Observations have shown that the enhancement of the electron flux depends on the pitch angle and energy. It is also proposed that when the IP shock impinges on the magnetosphere, the electrons in the radiation belts are energized not only by induced electric field but also waves excited by low-energy electrons. Therefore, we conduct simulations for acceleration processes of both energized and low-energy electrons by using the global magnetohydrodynamics(MHD) simulation and Comprehensive Inner Magnetosphere-Ionosphere Model (CIMI). In MHD simulation, 12 solar wind conditions are imposed on the upstream boundary condition by changing solar wind velocity, solar wind density and  $B_z$  of the Interplanetary magnetic field (IMF). We examine the results of response of the electron flux, temperature anisotropy, the ratio of cyclotron frequency and plasma frequency, the ratio of hot and cold electron density and cold electron density.

We obtained the simulation results as follow. 1) Generally, when the IP shock arrives, energetic electrons ( $>50$  keV) in the dayside magnetosphere are accelerated by the sudden enhancement of the electric field associated with a compressional wave. On the nightside when southward IMF is imposed electrons are transported inward due to  $E \times B$  drift because the convection electric field is developed. 2) Temperature anisotropy is increased on the nightside by the  $E \times B$  drift. The value is more than 1 when southward IMF is imposed. 3) Plasmapause is slightly compressed by the compressional wave. Plasmapause is contracted by the convection electric field when southward IMF is imposed. 4) The ratio of plasma frequency and cyclotron frequency is decreased because the magnetic field is increased.

## Rapid acceleration of outer radiation belt electrons associated with solar wind pressure pulse : A coupling simulation of GEMSIS-RB and GEMSIS-GM

\*Masahiro Hayashi<sup>1</sup>, Yoshizumi Miyoshi<sup>1</sup>, Shinji Saito<sup>1</sup>, Yosuke Matsumoto<sup>2</sup>, Kunihiro Keika<sup>3</sup>, Tomoaki Hori<sup>3</sup>, Takanobu Amano<sup>3</sup>, Kanako Seki<sup>3</sup>, Shinobu Machida<sup>1</sup>

1. Institute for Space-Earth Environmental Research, Nagoya University, 2. Chiba University, 3. University of Tokyo

Relativistic electron fluxes of the outer radiation belt dynamically change in response to solar wind variations. There are several time scales for the particle acceleration in MeV energy range. One of the shortest acceleration processes is wave-particle interactions between drifting electrons and fast-mode waves induced by compression of the dayside magnetopause through interplanetary shocks (e.g., Li et al., 1993). In order to investigate how relativistic electrons are accelerated by fast-mode waves driven by solar wind pressure pulse, we perform a code-coupling simulation using the GEMSIS-RB test particle simulation (Saito et al., 2010) and the GEMSIS-GM global MHD magnetosphere simulation (Matsumoto et al., 2010). As a case study, the interplanetary pressure pulse with the dynamic pressure of  $\sim 5$  nPa is used as an up-stream condition. In the magnetosphere, the fast mode waves with the azimuthal electric field (negative  $E_{\phi}$  :  $|E_{\phi}| \sim 10$  mV/m) propagates from the dayside and then extends to the entire dayside magnetosphere from 0600 to 1800 MLT. Using the electric/magnetic fields simulated by the GEMSIS-GM, we calculate the electron motion with different initial conditions (energy, and pitch angle). As a result, the increase of electron fluxes occurs for a wide energy range and energy spectrum become hard. The acceleration depends on the initial energy of electrons. We also investigate initial pitch angle dependence of acceleration and find that the fluxes of electron whose initial pitch angle closer to  $90^\circ$  are largely enhanced. The pitch angle dependence may be a result of the latitudinal structure of the induced electric fields and the pitch angle dependence of the drift velocity.

The results of investigation for initial energy and pitch angle imply that the acceleration condition of electrons is related to propagation speed of fast-mode waves, drift velocity of electrons and the spatial structure of electric field.

Keywords: radiation belt

## Impact of interplanetary shock on ring current ions

Hiroki Tsuji<sup>1</sup>, \*Yusuke Ebihara<sup>1</sup>, Takashi Tanaka<sup>2</sup>

1. Research Institute for Sustainable Humanosphere, Kyoto University, 2. Kyushu University

An interplanetary (IP) shock is known to have a large impact on magnetospheric ions. We have performed test particle simulation under the electric and magnetic fields provided by the global magnetohydrodynamics (MHD) simulation developed by Tanaka et al. (2010). In this particular simulation, the solar wind speed was increased from 372 to 500 km/s in order to reproduce the IP shock. The number density in the solar wind was set to a constant to be 5 cm<sup>-3</sup>, and the Z component of the interplanetary magnetic field (IMF) was turned from +5 to -5 nT. Just after the arrival of the IP shock, a fast mode wave propagated tailward in the magnetosphere. Dawnward electric field comes in first, followed by duskward electric field. The amplitude of the electric field exceeded 20 mV/m. We reconstructed the evolution of phase space density of H<sup>+</sup>, He<sup>+</sup>, O<sup>+</sup> ions by tracing trajectories of the ions backward on the basis of Liouville's theorem. The trajectory and the phase space density are found to be drastically modified by the fast mode wave in different ways. (1) The ion flux increases at entire energy range. This effect has been traditionally considered. (2) Multiple energy-time dispersion appears in energy-time spectra of the ions with energy less than ~100 keV due to adiabatic acceleration at high latitudes. This is associated with bounce motion (bounce phase bunching), and consistent with the Cluster satellite observation reported by Zong et al. (2012). (3) The ion flux depends on gyro phase due to gyro phase bunching. The gyro phase bunching is prominent for ions with initial speed being comparable to, or less than the ambient ExB drift speed. These results imply that the guiding center approximation is invalid for the ring current ions when large-amplitude fast mode wave is propagating associated with the IP shock. We also calculated temperature anisotropy. The temperature anisotropy increases near the leading edge of the wave where the dawnward electric fields is strong. The increase in the temperature anisotropy may favor the excitation of electromagnetic ion cyclotron (EMIC) waves, and may lead to rapid precipitation of ions and electrons. We evaluated growth of EMIC waves by using KUPDAP (Kyoto University Plasmas Dispersion Analysis Program). We will discuss the effect of IP shock on growing EMIC waves.

Keywords: Ring current, Interplanetary shock

## Deeper and earlier penetrations of oxygen ions than protons into the inner magnetosphere observed by Van Allen Probes

\*Kenji Mitani<sup>1</sup>, Kanako Seki<sup>2</sup>, Kunihiro Keika<sup>2</sup>, Matina Gkioulidou<sup>4</sup>, Louis Lanzerotti<sup>3</sup>, Donald Mitchell<sup>4</sup>, Craig Kletzing<sup>5</sup>

1. Institute for Space-Earth Environmental Research, Nagoya University, 2. Graduate School of Science, University of Tokyo, 3. New Jersey Institute of Technology, 4. Applied Physics Laboratory, Johns Hopkins University, 5. Department of Physics and Astronomy, University of Iowa

It is observationally known that protons and oxygen ions are the main components of the ring current during magnetic storms and are considered to have different source and supply mechanisms. In order to characterize the ion supply to the ring current during magnetic storms, we study the properties of energetic proton and oxygen ion phase space densities (PSDs) during the 23-25 April 2013 geomagnetic storm observed by the Van Allen Probes spacecraft. We calculated ion PSDs for specific first adiabatic invariants (for proton; for oxygen ion) and the local pitch angles near 90 degrees. The PSD profiles as a function of L show that both proton and oxygen ions penetrated to  $L < 5$  during the main phase of the magnetic storm. The timing of oxygen ion penetration was approximately the same for all values. The observations also show that oxygen ions penetrated more deeply in L and earlier in time than protons for the same value. The early penetration of oxygen ions suggest that the source of the transported oxygen ions was located closer to the Earth than the inner edge of plasma sheet protons. We also discuss the possibility that the interaction between  $>200$  keV oxygen ions and Pc3 ULF waves in the inner magnetosphere causes selective transport of oxygen ions. Our results imply the importance of the contribution from  $>200$  keV oxygen ions to the storm-time ring current.

Keywords: Van Allen Probes, Ring Current, Oxygen ion

## Oxygen impulsive energization during the storm main phase and its contribution to the ring current buildup

\*Kunihiro Keika<sup>1</sup>, Kanako Seki<sup>1</sup>, Yoshizumi Miyoshi<sup>2</sup>, Masahito Nose<sup>3</sup>, Louis J. Lanzerotti<sup>4</sup>, Donald G. Mitchell<sup>5</sup>, Matina Gkioulidou<sup>5</sup>, Andrew Gerrard<sup>4</sup>, Harlan Spence<sup>6</sup>, Brian A. Larsen<sup>7</sup>, Jerry W. Manweiler<sup>8</sup>

1. Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo, 2. Institute for Space-Earth Environmental Research, Nagoya University, 3. World Data Center for Geomagnetism, Graduate School of Science, Kyoto University, 4. New Jersey Institute of Technology, 5. Applied Physics Laboratory, Johns Hopkins University, 6. University of New Hampshire, 7. Los Alamos National Laboratory, 8. Fundamental Technologies, LLC

We investigate proton and oxygen ion energization during the main phase of magnetic storms. This study addresses one of the unresolved issues: supply, transport, and acceleration of ionospheric oxygen ions in the inner magnetosphere during magnetic storms. It is also yet to be determined whether oxygen ion contribution to the storm-time plasma pressure (and the ring current) is spatially global or localized. Plasma pressure in the inner magnetosphere is dominated by proton and oxygen ions with energies of a few to a few hundreds of keV. It is well known that such energetic oxygen ions increase drastically on a short time scale (< a few tens of minutes). We thus examine impulsive flux enhancements of protons and oxygen ions observed by the RBSPICE and HOPE instruments on board the Van Allen Probes spacecraft.

Van Allen Probes observed oxygen ion enhancements during the main phase of the 17 March 2015 storm. The oxygen energy density showed different temporal variations and radial profile from the proton energy density. It was enhanced during the early main phase ( $Dst \sim -120$  nT) up to the proton energy density level in an L range of 3 to 5. However, it decreased by about an order of magnitude around the beginning of the later main phase. It was increased again during the later phase ( $Dst \sim -220$  nT) particularly at  $L \sim 3$ , while it did not reach the early phase level. The radial profile was affected by temporarily impulsive enhancements more significantly than the proton energy density. The difference between the outbound (pre-midnight) and inbound (around midnight) paths is much clearer for oxygen ions than protons.

In this poster, we show the results of multi-event studies on such mass-dependent features during magnetic storms that occurred in 2013 to 2016. Our analysis is particularly focused on changes of energy spectra and pitch angle distributions, and spatial distributions of the oxygen ion contribution to the ring current. We discuss when and where ionospheric oxygen ions are energized to make a significant contribution to the ring current.

Keywords: Magnetic storms, Ring current, Oxygen ion outflow, Plasma transport and acceleration

# Modeling Results of the Two Largest Geomagnetic Storms of Solar Cycle 24

\*Yihua Zheng<sup>1</sup>, Lutz Rastaetter<sup>1</sup>, Maria M Kuznetsova<sup>1</sup>

1. NASA Goddard Space Flight Center

In this paper, radiation belt response to the two largest geomagnetic storms of Solar Cycle 24 (17 March 2015 and the 22 June 2015) is investigated in detail. Even though both storms are primarily CME driven, each has its own complexities [Liu et al., 2015, Kataoka et al., 2015]. Using the CCMC's run-on-request system, modeling results using the RBE (Radiation Belt Environment) model within the SWMF (Space Weather Modeling Framework) and the RBE model coupled with the SWMF and RCM (Rice Convection Model, which takes the ring current's contribution into consideration) will be examined. Comparative and comprehensive analyses of the same event from two different models and of two events from the same model/model suite will be provided. Focus will be specially given to impacts of different solar wind drivers on radiation belt dynamics and to the coupling and interactions of different plasma populations/physical processes within the region.

Keywords: radiation belt dynamics, largest geomagnetic storms, coupling

## Statistical Analysis of the Spatial Distribution of Low Frequency Magnetosonic Waves and Proton Ring-like Distribution


\*Kazuhiro Yamamoto<sup>1</sup>, Masahito Nose<sup>2</sup>, Craig A Kletzing<sup>3</sup>, Charles W Smith<sup>4</sup>, Robert J MacDowall<sup>5</sup>, George B Hospodarsky<sup>3</sup>, Harlan E Spence<sup>4</sup>, Geoff D Reeves<sup>6,7</sup>, Brian A Larsen<sup>6,7</sup>

1. Graduate School of Science, Kyoto University, 2. Data Analysis Center for Geomagnetism and Space Magnetism, Graduate School of Science, Kyoto University, Kyoto, Japan, 3. Department of Physics and Astronomy, University of Iowa, Iowa City Iowa, USA, 4. Institute for the Study of Earth, Oceans and Space, University of New Hampshire, Durham, New Hampshire, USA, 5. Solar System Exploration Division, Goddard Space Flight Center, Greenbelt, Maryland, USA, 6. Space Sciences and Applications Group, Los Alamos National Laboratory, Los Alamos, New Mexico, USA, 7. Space Sciences Division, The New Mexico Consortium, Los Alamos, New Mexico, USA

We statistically investigate the spatial distribution of magnetosonic waves at  $f < 32$  Hz and proton ring-like distribution observed by Van Allen Probes from September 2012 to December 2016. The spatial distribution of magnetosonic waves has an occurrence peak at  $L = 4 - 6$  and  $13 - 16$  MLT and that of proton ring-like distribution has an occurrence peak at  $L = 4 - 7$  and  $13 - 17$  MLT. The coincidence of the occurrence frequency peaks suggests that proton ring-like distribution is likely to be an energy source of magnetosonic waves. We reveals that the proton ring-like distribution with  $V_r > 2V_A$  has potential to excite magnetosonic waves at  $f < 32$  Hz, where  $V_r$  and  $V_A$  are ring velocity and Alfvén velocity, respectively. Case studies of convective growth rate analysis confirms the possibility of wave excitation by the proton ring-like distribution near the frequency of waves observed by satellites in these cases. Under the disturbed magnetospheric condition, the occurrence rate of magnetosonic waves increase up to 10 % and the ring energy increases up to  $\sim 20$  keV. This is consistent with an idea that and the high ring energy satisfies the wave growth condition of  $V_r > 2V_A$ . The condition of wave excitation at low frequency is attributed of a weighting function included in the calculation of the convective growth rate. A statistical analysis of the wave frequency reveals that magnetosonic waves in plasma trough are observed around the multiples of local proton cyclotron frequency except the first harmonics and most of them are considered to be excited locally, while some of magnetosonic waves observed inside the plasmopause seems to propagate from the other region.

Keywords: inner magnetosphere, plasma waves, magnetosonic waves, equatorial noise, ring like distribution, Van Allen Probes

Probe B/2015-11-18



L			
MLT [hr]	5,7	5,1	4,1
GMLAT [°]	14,0	14,7	15,7
hhmm	0,7	-0,7	-3,6
2015 Nov 18	1100	1200	1300

## Nonlinear generation mechanism of EMIC falling tone emissions

\*Masafumi Shoji<sup>1</sup>, Yoshiharu Omura<sup>2</sup>

1. Institute for Space-Earth Environmental Research, Nagoya University, 2. Research Institute for Sustainable Humanosphere, Kyoto University

We have conducted a self-consistent hybrid simulation, successfully reproducing EMIC emissions with falling-tone frequencies. The hybrid simulation is implemented with a parabolic ambient magnetic field. In the simulation, strong oxygen band EMIC emissions are generated through nonlinear wave growth. The cold ion density is modulated by electrostatic structures which are induced by the forward and backward propagating oxygen band EMIC waves. Along with the growth of the oxygen band, the helium band waves also grow because of the linear growth and the nonlinear growth. The nonlinear growth of the helium band waves is affected by the cold plasma density modulation, and there appear short wave packets of helium band emissions. The short wave packets entrap energetic protons efficiently, resulting in electromagnetic proton hills in the velocity phase space. The proton hill forms a nonlinear resonant current causing the falling frequency of the EMIC waves. We find strong deformation of the velocity distribution function of the energetic protons due to the proton hill being guided by the increasing resonance velocity.

## A statistical examination of favorable plasma conditions concerning inner magnetosphere EMIC wave excitation

Anthony A. Saikin<sup>1</sup>, \*Jichun Zhang<sup>1</sup>, Charles W. Smith<sup>1</sup>, Harlan E. Spence<sup>1</sup>, Brian A. Larsen<sup>2</sup>, Geoff D. Reeves<sup>2</sup>, Roy B. Torbert<sup>1</sup>, Craig A. Kletzing<sup>3</sup>, Irina S. Zhelavskaya<sup>4</sup>, Yuri Y. Shprits<sup>4</sup>

1. University of New Hampshire, 2. Los Alamos National Laboratory, 3. University of Iowa, 4. Helmholtz-Zentrum Potsdam - Deutsches GeoForschungsZentrum GFZ

We examine plasma conditions associated with the excitation of electromagnetic ion cyclotron (EMIC) waves in the inner magnetosphere ( $L < \sim 7$ ). Measurements from the Van Allen Probes ( $r = 1.1 - 5.8 R_E$ ) are used. EMIC wave events were identified from the polarization analysis of high-resolution magnetic field data from the Electric and Magnetic Field Instrument Suite and Integrated Science (EMFISIS) onboard the Van Allen Probes. The time span is from November 2012 to August 2014, which is one complete Van Allen Probes magnetic local time (MLT) precession. Plasma measurements were obtained from the Helium, Oxygen, Proton, Electron (HOPE) instrument. We calculate the observational growth parameter ( $\Sigma_h$ ) of the EMIC waves and the theoretical EMIC instability threshold ( $S_h$ ) to determine if the plasmas are favorable for EMIC wave excitation, i.e.,  $\Sigma_h - S_h > 0$ .  $\Sigma_h$  and  $S_h$  are calculated using measurements of the hot ( $>1$  keV) proton anisotropy, parallel hot proton plasma beta, the hot proton density, and the density of electrons. We examine occurrence rates and spatial distributions for the wave-favorable plasma conditions and the ratios of wave vs. non-wave occurrences under these conditions. Plasmas most favorable for EMIC wave generation are primarily observed at the probe apogee ( $L = \sim 6$ ). Peak EMIC wave occurrence is found in the afternoon - midnight (1600 - 0100) MLT sectors. This same region coincides with the enhancements of parallel hot proton plasma beta and hot proton density. Hot proton anisotropy measurements peak in the midnight - noon (0 - 1200) MLT sectors.

Keywords: Plasma waves and instabilities, Wave/particle interactions, Magnetospheric configuration and dynamics, Magnetosphere - inner, Ring current

# EMIC waves-driven radiation belt electron precipitation into the atmosphere with ground-based observations in the subauroral region

\*Hirai Asuka<sup>1</sup>, Fuminori Tsuchiya<sup>1</sup>, Takahiro Obara<sup>1</sup>, Hiroaki Misawa<sup>1</sup>, Kazuo Shiokawa<sup>2</sup>, Yoshizumi Miyoshi<sup>2</sup>

1. Planetary Plasma and Atmospheric Research Center, Graduate School of Science, Tohoku University, 2. Institute for Space-Earth Environmental Research, Nagoya University

Energetic electron losses from the outer radiation belt occur during magnetic storm and substorm. One of the mechanisms is precipitation into the atmosphere and electromagnetic ion cyclotron (EMIC) waves are one of candidates to cause pitch angle scattering of energetic electron. EMIC waves, which are observed in the Pc1–Pc2 frequency range (0.1–5Hz) are excited by the ion cyclotron instability in the equatorial region of the magnetosphere during the main and the recovery phase of magnetic storms. It has been theoretically studied that EMIC waves play an important role in energetic electron precipitation into the atmosphere, but there have been limited experimental observations to support this idea.

Here, we investigated relation between occurrence of EMIC waves and energetic electron precipitation by means of ground-based magnetometers and low frequency (LF) radio wave propagation observation and confirmed EMIC waves to be driving electron precipitation.

We use induction magnetometer data in North America (ISEE and CARISMA stations) to investigate occurrence of EMIC waves. LF radio wave signals transmitted from WWVB, United States (40.7°N, 255.0°E, L=2.28), are observed at Athabasca, Canada (54.7°N, 246.7°E, L=4.35) to investigate precipitation of energetic electron (>100keV) into the atmosphere. LF radio waves propagate, reflecting between earth's surface and the lower ionospheric boundary (altitude ≈ 70–90km). Ionization caused by precipitating electron in the lower ionosphere changes altitude of the reflection height, resulting in a deviation of the LF wave phase from that in undisturbed conditions.

We detected energetic electron precipitation from the LF radio wave observation in 07:00–09:20 UT on July 7, 2011 and EMIC waves were observed by the induction magnetometer at Athabasca in 05:35–10:55 UT on the same day. At the almost same time, EMIC waves were observed at some CARISMA stations. These observations indicate that EMIC waves are expected to cause detected electron precipitation. Polarization characteristics of EMIC waves which reflect locations where the waves inject into the ionosphere and direction of subsequent horizontal propagation in the F-region were examined by cross-spectrum analysis of EMIC waves.

Based on time variations in intensity, frequency, polarization sense, and angle of the major axis, the period of EMIC appearance could be divided into six sequential events. This suggests that source of the EMIC waves observed in 05:35–10:55 UT was not a single but consisted of multiple locations.

We found that time variation of the LF wave phase corresponds to that of EMIC waves, and the deviation of the LF wave phase only occurred during the 2nd, 3rd and 4th EMIC events. This result implies that the source locations of the three EMIC events were close to the Athabasca–WWVB radio wave propagation path and the EMIC-driven energetic electron precipitation caused the phase deviation of WWVB signal. Identification of actual source locations of the EMIC events is a future work.

## Acknowledgement

The authors thank I.R. Mann, D.K. Milling and the rest of the CARISMA team for data. CARISMA is operated by the University of Alberta, funded by the Canadian Space Agency.

Keywords: EMIC waves, energetic electron precipitation

## Polarization characteristics of Pc1 pearl structure observed at Kawatabi, Osaki, Miyagi prefecture

Shinya Miyake<sup>1</sup>, Seiya Yamakawa<sup>1</sup>, Yuma Doi<sup>1</sup>, \*Tomoko Nakagawa<sup>1</sup>

1. Information and Communication Engineering, Tohoku Institute of Technology

Pc1 geomagnetic pulsations are often found in the ELF magnetic field data obtained by two sets of induction magnetometers EL-12 constructed by Tierra Technica Co. Ltd, placed in North-South and East-West directions at Kawatabi, Osaki, Miyagi prefecture Japan. The magnetic latitude of the observation site is N30 and the L value is about 1.3.

Although the data coverage was not very good, we have found 7 examples of pearl structures, within the frequency range of 1 to 5 Hz. They showed temporal variation of bandwidth such as 0.3 to 1.2Hz, or 0.9 to 1.8 Hz, forming pearl structures in dynamic spectra. The frequency itself also varied with time: they rose in 3 cases found in pre-midnight, and fall in 4 cases in the pre-midnight region.

Polarization of the magnetic variation was examined by using Fourier components of N-S and E-W magnetic field components. We have 4 events for which E-W and N-S observations were available. The polarization was steady and right-handed for one event, but for the rest, it was variable. One event showed left-handed polarization at higher frequency and right-handed in the lower. Other two events showed alternative polarizations pearl to pearl.

Keywords: Pc1, pearl, pulsation, ELF, magnetic variation, polarization

## Formation of butterfly pitch angle distributions of relativistic electrons in the outer radiation belt due to the drift resonance with a monochromatic Pc5 wave

\*Kei Kamiya<sup>1</sup>, Kanako Seki<sup>2</sup>, Shinji Saito<sup>1</sup>, Takanobu Amano<sup>2</sup>, Yoshizumi Miyoshi<sup>3</sup>

1. Graduate School of Science, Nagoya University, 2. Department of Earth and Planetary Science, Graduate School of Science, University of Tokyo, 3. Institute for Space-Earth Environmental Research, Nagoya University

Radial transport of relativistic electrons in the inner magnetosphere can be driven by drift resonance with Pc5 Ultra Low Frequency (ULF) waves. The radial transport due to the drift resonance has been considered as one of important acceleration mechanisms of the outer radiation belt electrons. In the course of the radial transport, the energy and equatorial pitch angle of electron change under conservation of the first and second adiabatic invariants. The change in the drift period in the course of radial transport thus depends on the adiabatic process, and it can affect the radial transport rate of the electrons. In other words, the radial transport rate due to the drift resonance depends on the equatorial pitch angle and can form the characteristic pitch angle distributions (PADs). In this study, we investigate the radial transport of relativistic electrons due to the drift resonance with a monochromatic Pc5 wave and focus on formation of PADs of the outer radiation belt electrons.

We use two simulation models of the inner magnetosphere: GEMSIS-Ring Current (RC) and GEMSIS-Radiation Belt (RB). The RC simulation, which is a self-consistent and kinetic numerical simulation code, solves the five-dimensional Boltzmann equation for the ring-current ions coupled with Maxwell equations. The RB simulation calculates trajectories of guiding center of test-particles in arbitrary magnetic and electric field. We used electric and magnetic fields of a monochromatic Pc5 wave in the inner magnetosphere obtained from the RC simulation as background fields in the RB simulations. We traced an order of  $10^7$  of radiation belt electrons to calculate phase space density of the electrons at each position in the equatorial plane.

Simulation results show formation of characteristic PADs depending on the energy and location (L value), which can be explicable of the pitch angle dependence of resonance conditions. At some fixed location and energy range, the PADs can change from pancake-like to butterfly-like distributions, as the transport by the monochromatic Pc5 wave progresses. These butterfly distributions can be seen when electrons with small (oblique) pitch angles satisfy the resonance condition. It is also found that the small pitch angle electrons can be transported further inward because PA change to larger value through the adiabatic transport enables them to satisfy resonance condition in wider L range compared to the 90-degrees PA electrons.

Keywords: Radiation belt electrons, Drift resonance, Pitch angle distributions

## Localtime Dependence of the Pc5 Wave associated with MeV Electron Flux Enhancement Observed by two GOES Satellites

\*Kentarou Kitamura<sup>1</sup>, Satoko Saita<sup>2</sup>, Yoshimasa Tanaka<sup>3</sup>, Akiko Fujimoto<sup>4</sup>

1. NIT, Tokuyama College, 2. NIT, Kitakyushu College, 3. National Institute of Polar Research, 4. Kyushu Univ.

It is well known that MeV electron flux efficiently increases during the recovery phase of magnetic storms. ULF wave propagating in the magnetosphere is recognized as one of the possible candidates which can accelerate the electron in the radiation belt while various acceleration processes have been widely proposed by many investigators.

In this study, total 20 electron flux enhancement events associated with the CIR (Corotating Interaction Region) driven storms in 2008 have been analyzed using the magnetic field vector data obtained by GOES 10 and 11 satellites. The GOES 10 and 11 were located at 60 deg. and 75 deg. West in geographical longitude, respectively, which corresponds to 1 hour separation in local time. We used the bandpass filtered (150-1000 sec) magnetic data in the ENP coordinate system to investigate the oscillation mode of the field line and the propagation characteristics of Pc5 pulsations in the GEO orbit (6.6 Re).

As a result, following features are observed, that is: both the P (compressional mode) and T (transverse mode) components of the Pc5 strongly enhances at the beginning of the electron flux decreasing in the night side sectors: the Pc5 power is relatively low at the morning sectors: the dominant frequencies vary from high to low during the electron flux decreasing, which is quite apparent at the afternoon-night sectors. These observational facts indicate that the source region of the Pc5 during the electron flux decreasing can be considered at the evening sectors. The particle injection from night side associated with substorms may generate the ULF wave in the evening sectors. The decreasing of the dominant frequencies also suggests the particle injection from night side associated with substorms from the night to evening sectors.

Keywords: ULF wave, MeV electron flux, Substorm

## ULF wave modulation on the generation process of whistler-mode chorus emissions

\*Yuto Katoh<sup>1</sup>, Chen Lunjin<sup>2</sup>

1. Graduate School of Science, Tohoku University, 2. University of Texas at Dallas

We study the modulation of the generation process of whistler-mode chorus emissions under the presence of ULF waves in the inner magnetosphere. Previous studies revealed properties of chorus generation depending on the number density of energetic electrons, temperature anisotropy of velocity distribution function, and spatial gradient of the background magnetic field [Katoh and Omura, 2011, 2013, 2017]. The properties of both energetic electrons and the background magnetic field are also varied by the presence of ULF waves in the inner magnetosphere [e.g., Xia et al., 2016]. The range of parameters controlling chorus generation should be examined by a self-consistent simulation reproducing the generation process of chorus emissions. By referring the range of variations of the background magnetic field for toroidal and poloidal mode ULF waves, we carry out a series of electron hybrid code simulations, changing number density and temperature anisotropy of energetic electrons. Simulation results clarify that the variation of the spatial gradient of the background magnetic field controls whether or not distinct chorus emissions are generated from the magnetic equator. The results of the present study serve useful information in understanding in-situ observation of both chorus and ULF waves and related wave-particle interactions occurring in the inner magnetosphere.

Keywords: whistler-mode chorus, wave-particle interaction, numerical experiments

## Van Allen Probes Observations of Modulation of Energetic Ion Fluxes by a Fundamental Poloidal Mode Standing Alfvén Wave

\*Kazue Takahashi<sup>1</sup>, Seth Claudepierre<sup>2</sup>, Craig Kletzing<sup>3</sup>, Charles W Smith<sup>4</sup>

1. The Johns Hopkins University Applied Physics Laboratory, USA, 2. The Aerospace Corporation, USA, 3. Department of Physics and Astronomy, University of Iowa, USA, 4. Physics Department and Space Science Center, University of New Hampshire, USA

Magnetospheric ULF waves are known to cause periodic modulations of the flux of ring current ions (energy range  $\sim 10$ -300 keV). Many previous studies reported ion flux modulation events associated with second harmonic poloidal mode standing Alfvén waves. In this study, we report Van Allen Probes observations of an ion flux modulation event associated with a fundamental poloidal wave. The wave (period  $\sim 100$ s) was observed on 6 October 2012 in the prenoon and produced a giant pulsation on the ground. The standing wave mode was unambiguously determined from the relationship between the electric and magnetic field perturbations at the spacecraft. The field oscillations were accompanied by oscillations of the flux of ions over an energy range of 100-200 keV. Contrary to previously reported similar ULF wave events with strong modulation of equatorial protons, the flux oscillations in the present event were strongest at pitch angles around 30 degrees. The amplitude and phase of the ion flux oscillations exhibited signatures of drift resonance at 150 keV, from which the wave is inferred to be propagating westward with an azimuthal wave number of 35. This wave number is consistent with the ion finite Larmor radius effects seen in ion fluxes measured at different phases of the spacecraft spin. The ion phase space density exhibits a radial gradient that is consistent with theoretical prediction of an instability involving drift resonance of ring current ions with fundamental poloidal waves.

Keywords: Fundamental standing Alfvén wave, Van Allen Probes, Ring current ions

## Landau resonance between electrons and lower-hybrid waves in the inner magnetosphere

\*Fumiko Otsuka<sup>1</sup>, Kaiti Wang<sup>2</sup>, Shuichi Matsukiyo<sup>1</sup>, Tohru Hada<sup>1</sup>

1. Interdisciplinary Graduate School of Engineering Sciences Kyushu University, 2. Dept. Aerospace Engineering, Tamkang University

Lower-hybrid waves are frequently observed near the geomagnetic equator in the inner magnetosphere (i.e., equatorial noise). They are in the frequency range between the proton gyrofrequency and the LH frequency, and were found to propagate approximately perpendicular to the background magnetic field with almost linear polarization. We have focused on the capability of the LH waves to scatter electrons, and showed that the diffusions could occur via both cyclotron and Landau resonances. To have the cyclotron resonance to occur, the electron energies should be higher than 1.56 MeV. On the contrary, the Landau resonance occurs even for relatively lower energies from 1.4 keV. Here, the linear resonance condition is assumed under the observed LH wave parameters such as the propagation angle of 85 degree and the frequency of 130 Hz in a plasma environment with the Alfvén velocity of 1150 km/s.

In this presentation, we discuss the Landau resonance between electrons and LH waves, by performing test particle simulation. The LH waves are given as a superposition of sinusoidal waves with different frequencies propagating highly perpendicular to the background magnetic field. The given waves obey the cold plasma dispersion relation. We evaluate the pitch-angle diffusion coefficient of electrons with energies from a few eV to 1 MeV. We discuss changes in pitch-angle distributions related to the diffusion processes.

Keywords: pitch-angle diffusion, electron, lower-hybrid wave, inner magnetosphere

# Nonlinear pitch angle scattering of low pitch angle electrons caused by whistler mode chorus emissions

\*Masahiro Kitahara<sup>1</sup>, Yuto Katoh<sup>1</sup>

1. Department of Geophysics, Graduate School of Science, Tohoku University

Whistler-mode chorus emissions are defined as coherent emissions with frequency sweep, and are frequently observed by various satellites in the Earth's inner magnetosphere. Chorus emissions are generated by energetic electrons in the kinetic energy range from a few to tens of keV through nonlinear wave-particle interactions, and energetic electrons are scattered their pitch angle by generated chorus emissions. The pitch angle scattering is closely related to energetic electron precipitation into the ionosphere, contributing to diffuse and pulsating aurora. Conventionally, it has been considered that particles satisfying the cyclotron resonance condition are scattered toward the loss cone by whistler mode waves. Li et al. (2015) indicated, however, that low pitch angle particles tend to be scattered away from the loss cone by coherent whistler mode waves. Omura et al. (1991) reviewed the study of the motion of particles under the presence of coherent waves and represented the equation of motion of particle near the resonance condition as a pendulum equation. They assumed in the derivation of the equation that the pitch angle of particles is not small (Nunn, 1974).

In this study, we derive the equation of the motion of particles without the assumption of small pitch angle to consider pitch angle scattering near the loss cone in the velocity phase space. We clarify that electrons near the loss cone satisfying the cyclotron resonant condition are scattered away from the loss cone due to the Lorentz force caused by the wave magnetic field and the parallel velocity component of electrons. In order to reproduce the pitch angle scattering caused by chorus emissions, we carry out a test particle simulation using the simulation system along a dipole magnetic field line and a whistler mode wave model. Results of the test particle simulation are consistently explained by the nonlinear theory we derived, and the pitch angle variation due to the nonlinear effect strongly depends on the wave amplitude. In particular, for the case of the large amplitude wave, most of resonant electrons are trapped by the coherent wave and are efficiently scattered away from the loss cone, resulting in less precipitating electrons. Furthermore, assuming 20 keV electrons uniformly distributed in the pitch angle range from 0 to 90 degrees, we estimate the modulation of pitch angle distribution while electrons encounter one wave packet of chorus emissions. Our results indicate that most of low pitch angle electrons scattered away from the loss cone and build a bump of distribution at the moderate pitch angle satisfying the cyclotron resonant condition. These results suggest that the relation between chorus wave intensity and the flux of auroral electron precipitation is not straightforward and that the nonlinear effect newly proposed by the present study should be taken into account.

Keywords: chorus emissions, pitch angle scattering, wave-particle interaction

# Three-Dimensional Forward Modeling of Lightning-Induced Electron Precipitation from the Radiation Belts

\*Austin P Sousa<sup>1</sup>, Robert A Marshall<sup>2</sup>

1. Stanford University, 2. University of Colorado, Boulder

Pitch-angle scattering by radio waves in the VLF (3-30kHz) band is thought to be a major loss mechanism for energetic radiation-belt electrons. Resonant interactions with Whistler-mode VLF waves can alter the reflection altitude of trapped electrons 100keV - 1MeV; when a particle reflects at a low enough altitude, it can be removed from the magnetosphere through collisions with ionospheric constituents. Terrestrial lightning provides a natural and constantly-occurring source of VLF waves. Here we present a three-dimensional forward model of lightning-induced electron precipitation (LEP) due to resonant pitch-angle scattering from a single lightning stroke.

Previous efforts (Lauben 1998, Bortnik 2006) have used two-dimensional raytracing combined with analytical expressions of pitch-angle scattering to forward model precipitation from a single stroke as a function of input and output latitude. However these models are limited in geospatial accuracy by their use of ideal plasmasphere and magnetic field models. We expand on these techniques by incorporating three-dimensional raytracing through a realistic plasmasphere and magnetic field model, to better capture the spatial dependence of LEP.

We then combine our end-to-end model of the LEP process with terrestrial lightning activity data from the GLD360 sensor network to construct a realtime geospatial model of LEP-driven energy deposition into the ionosphere. We explore global and seasonal statistics, provide precipitation estimates across a variety of magnetospheric conditions, and compare the total impact to other magnetospheric loss processes. The completed model is well-suited for comparison with satellite electron flux measurements, such as those from the Arase mission.

Keywords: Wave-Particle Interactions, Radiation Belts, Lightning-Induced Electron Precipitation

## Acceleration of energetic electrons by oblique whistler-mode chorus in the radiation belt

\*Yikai HSIEH<sup>1</sup>, Yoshiharu Omura<sup>1</sup>

1. Reserach Institute for Sustainable Humanosphere, Kyoto University

We perform test particle simulations for relativistic electrons interacting with a whistler-mode chorus packet propagating at oblique angles. The properties of group velocity of obliquely propagating whistler mode waves are analyzed. The group velocity of lower band chorus is nearly parallel to the magnetic field, which justify the gyroaveraging method. In the gyroaveraging method, we calculate the equations of motion of electrons averaging the cyclotron motion at gyrocenter and reducing the simulation from two-dimensional system to one-dimensional system. In the simulations, we found that multiple resonances are essential in electron accelerations. We trace evolution of a delta function of relativistic electrons in a phase space of kinetic energy and equatorial pitch angle, and then obtain numerical Green's functions of the chorus wave-particle interactions. The efficiency of the MeV electron acceleration by the Landau resonance is confirmed by examining the Green's functions in a wide range of kinetic energies. We investigate the rate of energy gain of the cyclotron resonance acceleration and the Landau resonance acceleration, and find that the perpendicular component of wave electric field dominates both accelerations for MeV electrons. Furthermore, we find that the efficient acceleration of MeV electrons is contributed by the proximity between the parallel components of  $V_p$  and  $V_g$  of oblique whistler mode waves. The proximity makes the interaction time of the Landau resonance much longer than that of the cyclotron resonance, resulting in efficient acceleration of MeV electrons.

Keywords: Whistler mode waves, Relativistic electrons, Landau resonance

# Nonlinear dynamics of resonant particles interacting with coherent waves and comparison with quasi-linear theory

\*Miwa Tobita<sup>1</sup>, Yoshiharu Omura<sup>1</sup>

1. Research Institute for Sustainable Humanosphere, Kyoto University

We perform test particle simulations of charged particles in an external constant electric field and electrostatic wave fields. We solve equations of motion, which take the same form as the pendulum equation, and we study nonlinear dynamics of particles with different values of inhomogeneity factor  $S$  defined as a ratio of the wave amplitude to the background electrostatic field. The target of the present study is to understand nonlinear dynamics of resonant particles interacting with coherent waves in space plasmas. Electromagnetic waves such as whistler-mode chorus, hiss emissions, and electromagnetic ion cyclotron (EMIC) waves in the inner magnetosphere contain structures of coherent waves with various discrete frequencies. Their interaction with resonant particles can be approximated by the nonlinear pendulum equation, or the equations of motion for a charged particle in a one-dimensional electrostatic wave potential. Conventionally, particle scattering and energization are studied as particle diffusion by incoherent waves without any background field causing adiabatic variation of the drift velocity. However, this model, called quasi-linear diffusion model, cannot explain “super diffusion”, referring to MeV energization and precipitation of particles in a few minutes, found in observations of chorus, hiss, and EMIC waves. To describe the super diffusion with the simplified electrostatic model, we introduce the external constant field, which corresponds to an inhomogeneous magnetic field causing adiabatic motion such as mirror motion. All particles are accelerated uniformly and constantly as an adiabatic motion, and they are dramatically heated by scattering and/or nonlinear trapping through resonant interaction with the waves. The present simulation model demonstrates the super diffusion due to coherent waves and the external field.

Keywords: wave-particle interaction, nonlinear wave trapping, particle acceleration

# Dynamics of energetic electrons interacting with sub-packet chorus emissions in the outer radiation belt

\*RYOKO HIRAGA<sup>1</sup>

1. Kyoto University Electricity Engineering Research Institute for Sustainable Humanosphere

Our preceding study examined the efficient electrons acceleration processes called Relativistic Turning Acceleration (RTA) and Ultra-relativistic Turning Acceleration (URA), and the mechanism how the outer radiation belt is formed. This time, we undertake an update of the chorus wave model used in the preceding simulation in order to reflect the observational data more precisely. By referring to the latest observations by Van Allen Probes (Foster et al. 2017), we update the chorus wave source model excited around the equator whose amplitude was assumed to grow monotonically in the preceding simulation. The new wave model in the current simulation represents the sub-packet structure in its amplitude variation. Sub-packet amplitude structure is such that when the wave amplitude nonlinearly grows to reach the optimum amplitude, it starts decreasing until crossing the threshold. Once it crosses the threshold, the wave dissipates and a new wave arises to repeat the nonlinear growth and damping in the same manner. The multiple occurrence of these wave generation to dissipation processes forms a saw tooth-like amplitude variation called sub-packet. This sub-packet structure is one of the most distinctive features of chorus waves we can find from the observations and hence should be carefully included in the simulation wave model. Due to the rapid variation of amplitude sub-packet structure, however, the wave frequency as a function of amplitude also undergoes a fluctuation in time variation. This fluctuation is assumed to decrease the duration and efficiency of wave-electron resonance and resultant electron acceleration. We examine the electrons acceleration processes including RTA and URA by the sub-packets and analyze the formation mechanism of a highly energized radiation belt. First we insert 36 test particles assigned with different gyro-phases for 3 different initial energy levels: 500keV, 1MeV and 2MeV with the pitch angle of 85 degrees. The simulation results here are compared with those from the preceding study to well understand the acceleration mechanism of individual electrons. Based on this, we next conduct a statistical analysis how these accelerated electrons collectively form the outer radiation belt. We apply the Green's function method covering a sufficient number of electrons with the initial energies from 10keV to 2MeV and the initial pitch angles from 10 to 90 degrees. By the overall simulations, we reach a conclusion on how in detail the individual electron is sufficiently accelerated by the sub-packet chorus waves, and how the accelerated electrons collectively form the outer radiation belt.

## Two-dimensional Particle Simulations of Oblique Whistler-mode Instability

\*Takeshi Nogi<sup>1</sup>, Yoshiharu Omura<sup>1</sup>

1. Reserch Institute for Sustainable Humansphere, Kyoto University

We perform two-dimensional electromagnetic particle simulations to study basic characteristics of whistler-mode

wave particle interaction involved in chorus emissions propagating oblique to the static magnetic field.

We assume

a simple periodic  $(x, y)$  system with the magnetic field taken in the  $x$ -direction. Assuming energetic electrons

with an anisotropic bi-Maxwellian velocity distribution function, we first test the linear whistler-mode instability

driven by temperature anisotropy to confirm the numerical property of the simulation code. With the electrostatic

components parallel to the magnetic field, which have been neglected in the previous simulation studies on chorus

emissions, we find the linear phase of the instability is much affected by the Electrostatic thermal fluctuations.

It is necessary to put many super-particles in a grid cell to suppress the thermal fluctuation. With 30,000 particles

per cell, we have confirmed a good agreement of the wave growth in the parallel direction with the linear growth

rate. We next put an array of antennas with obliquely aligned to uniform magnetic field, and oscillate the antenna

current with a variable frequency below the electron cyclotron frequency to excite a large amplitude whistler-mode

wave obliquely propagating to the static magnetic field. In addition to the nonlinear trapping of energetic electrons

through the cyclotron resonance, another nonlinear trapping of electrons by the Landau resonance takes place.

Structures of the nonlinear trapping potentials changes with a varying frequency, affecting the efficiency of

energy transfer between the wave and energetic electrons. We study nonlinear evolution of the wave packet, and

competing processes of both resonances in accelerating the energetic electrons to higher energies.

## A statistical investigation of plasmaspheric plasma by using global GPS TEC data

\*Qing-He Zhang<sup>1</sup>, Gun Li<sup>2</sup>, Yan-Qiu Ren<sup>1,2</sup>, Yong Wang<sup>1</sup>, John Foster<sup>3</sup>, Michael Lockwood<sup>4</sup>, Shun-Rong Zhang<sup>3</sup>

1. Institute of Space Sciences, Shandong University, Weihai, China, 2. School of Aeronautics and Astronautics, University of Electronic Science and Technology of China, Chengdu, China., 3. Haystack Observatory, Massachusetts Institute of Technology, Westford, Massachusetts, USA, 4. Department of Meteorology, University of Reading, Earley Gate, Post Office Box 243, RG6 6BB, UK.

A statistical analysis of the daily, monthly, yearly and solar cycle variations of plasmaspheric plasma from near earth to the dayside magnetopause is investigated by using the equatorial mapping data of the ground-based global position system (GPS) total electron content (TEC) from the noon meridian for the years of 2003-2015. During the geomagnetic storms, the mapped TEC data often showed clear plume structures in the afternoon sectors, which had been detached from plasmasphere towards sunward and reached the dayside magnetopause. However, the plasma in the plume was exhausted after the main phase of the storms and the plasmasphere needed one day or a few days to be refilled. This exhaustion may be because of the plasmaspheric plasma escaping through the dayside magnetopause where the magnetic reconnection or magnetopause shadowing occurred. The mapped TEC plumes and plasmasphere refillings preferred to appear around every afternoon and often reached the dayside magnetopause with different TEC value depending on the solar and geomagnetic conditions, which also preferred to appear in the months from March to May and from October to December and in the years during solar maximum. These results may suggest that the plasma in the plumes escaped away through the magnetopause due to the dayside magnetic reconnection or magnetopause shadowing during the recover phase of the geomagnetic storms, and the polar ionosphere continuously refills plasma into the plasmasphere during the quiet days, which can be stored and partially convected sunward to the dayside magnetopause for forming plumes.

Keywords: Plasmaspheric plasma, Plasmaspheric plume, plasmaspheric refilling

# Statistical Analysis of Substorm-Associated SAR arcs at Subauroral Latitudes Based on All-sky Imaging Observations at Athabasca, Canada

\*Yuuki Takagi<sup>1</sup>, Kazuo Shiokawa<sup>1</sup>, Yuichi Otsuka<sup>1</sup>, Martin Connors<sup>2</sup>

1. Institute for Space-Earth Environmental Research, Nagoya University, Japan, 2. Athabasca University, Canada

SAR arcs are the optical phenomenon caused by low-energy electron precipitation into the ionospheric F layer from the interaction region between the ring current and the plasmasphere. In the recovery phase of geomagnetic storms, low-energy electrons in the plasmasphere are heated by high-energy plasma in the ring current, and these electrons precipitate into the F layer at subauroral latitude where oxygen atoms are excited at altitudes about 400 km. Thus, SAR arcs have been observed at subauroral latitudes during geomagnetic storms. However, Shiokawa et al. (2009) reported an event of SAR arcs detached from the main oval after substorms, based on observation at Athabasca, Canada (54.7N, 246.7E, magnetic latitude = 61.7N). However, statistical analysis of such substorm-associated SAR arcs have not been done yet. Thus, in this study, we do a statistical analysis of substorm-associated SAR arcs observed at Athabasca. We analyzed all-sky images at wavelengths of 630.0 nm obtained at Athabasca from 3 September, 2005 to 31 December, 2009, and found 98 events. This result indicates that the SAR arcs are often detached from the main oval after substorms at Athabasca. We investigated dependences of these SAR arc appearances and their latitudes and durations on AU/AL indices, SYM-H, X component of magnetic field variation at Yellowknife (YKC), north of Athabasca in the auroral zone, solar wind pressure, and IMF-Bz. We found that when SAR arcs occur, AL and YKC-X component tend to decrease, indicating substorm association of these SAR arcs. We found that the SAR arc occurrence peak is around midnight with a peak rate of ~5 % with decreasing rates in both pre-midnight and post-midnight sectors. We then classified these SAR arcs into 3 types by using simultaneous 557.7-nm images as: 1) 557.7-nm images show weak structures similar to the 630.0-nm SAR arcs (30 %), 2) 557.7-nm images does not show structures similar to the 630.0-nm SAR arcs (55 %), 3) 557.7-nm images show structures different from the 630.0-nm SAR arcs (15 %). In the presentation, we will discuss possible cause of the detachment of SAR arcs from the main oval associated with substorms.

Keywords: SAR arc, substorm, MI coupling

## North-south asymmetry of vortex evolution in conjugate aurora

\*Herbert Akihito Uchida<sup>1</sup>, Ryuho Kataoka<sup>2,1</sup>, Akira Kadokura<sup>2,1</sup>, Yasunobu Ogawa<sup>2,1</sup>, Yoko Fukuda<sup>3</sup>, Yoshizumi Miyoshi<sup>4</sup>

1. Department of Polar Science, SOKENDAI (The Graduate University for Advanced Studies), 2. National Institute of Polar Research, 3. Department of Earth and Planetary Science, University of Tokyo, 4. Institute for Space-Earth Environmental Research, Nagoya University

The influence of the ionosphere on the aurora can be evaluated by examining the asymmetry of conjugate aurora at various time-space scales. We conducted a high-speed imaging observation of aurora at Tjornes/Iceland and Syowa/Antarctica for the time interval from 2 September 2016 to 7 September 2016 when high aurora activity continued for several days, which was driven by high-speed solar wind from a large coronal hole. It is found that an overall conjugacy is not good as we originally expected, even considering the modeled conjugate points. For example, vortex evolution from small scale (so called folds, a few ten km) to large scale (spirals, a few hundred km) occurred over whole field of view at Syowa, while such vortex structures themselves are hard to recognize at Tjornes at the same time. In this talk we discuss the conjugate morphology in more detail. Further we present the current situation and future development of the high-speed conjugate imaging observations.

Keywords: Aurora, Conjugacy, High-speed imaging